(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 25 May 2001 (25.05.2001)

PCT

(10) International Publication Number WO 01/35726 A1

(51) International Patent Classification7: 5/00, C12N 5/14, 15/82

A01H 1/00.

(21) International Application Number: PCT/US00/31418

Helen Place, Castro Valley, CA 94546 (US). REUBER, Lynne [US/US]; 2000 Walnut Avenue, Fremont, CA 94538 (US). ADAM, Luc [CA/US]; 25800 Industrial Boulevard, L403, Hayward, CA 94545 (US).

(22) International Filing Date:

ne: 14 November 2000 (14.11.2000)

(25) Filing Language:

;

English

(26) Publication Language:

English

(30) Priority Data:

60/166,228 17 November 1999 (17.11.1999) US 60/197,899 17 April 2000 (17.04.2000) US 60/227,439 22 August 2000 (22.08.2000) US

(71) Applicant (for all designated States except US): MENDEL BIOTECHNOLOGY, INC. [US/US]; 21375 Cabot Boulevard, Hayward, CA 94541 (US).

(71) Applicants and

(72) Inventors: HEARD, Jacqueline [US/US]; 810 Guildford Avenue, San Mateo, CA 94402 (US). RATCLIFFE, Oliver [GB/US]; 814 East 21st Street, Oakland, CA 94606 (US). CREELMAN, Robert [US/US]; 2801 Jennifer Drive, Castro Valley, CA 94546 (US). JIANG, Cai-Zhong [CN/US]; 34495 Heathrow Terrace, Fremont, CA 94555 (US). PINEDA, Omaira [CO/US]; 19563

- (74) Agent: GUERRERO, Karen; Mendel Biotechnology, Inc., 21375 Cabot Boulevard, Hayward, CA 94545 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

1/35726 A

(54) Title: PATHOGEN TOLERANCE GENES

(57) Abstract: The present invention relates to transgenic plants and methods of making transgenic plant using punitive transcription factors that modulate the transgenic plant's susceptibility to disease.

PCT/US00/31418

5

10

15

20

25

30

35

PATHOGEN TOLERANCE GENES

RELATED APPLICATION INFORMATION

The present invention claims the benefit from US Provisional Patent Application Serial Nos. 60/166,228 filed November 17, 1999 and 60/197,899 filed April 17, 2000 and "Plant Trait Modification III" filed August 22, 2000.

FIELD OF THE INVENTION

This invention relates to the field of plant biology. More particularly, the present invention pertains to compositions and methods for phenotypically modifying a plant.

BACKGROUND OF THE INVENTION

Transcription factors can modulate gene expression, either increasing or decreasing (inducing or repressing) the rate of transcription. This modulation results in differential levels of gene expression at various developmental stages, in different tissues and cell types, and in response to different exogenous (e.g., environmental) and endogenous stimuli throughout the life cycle of the organism.

Because transcription factors are key controlling elements of biological pathways, altering the expression levels of one or more transcription factors can change entire biological pathways in an organism. For example, manipulation of the levels of selected transcription factors may result in increased expression of economically useful proteins or metabolic chemicals in plants or to improve other agriculturally relevant characteristics. Conversely, blocked or reduced expression of a transcription factor may reduce biosynthesis of unwanted compounds or remove an undesirable trait. Therefore, manipulating transcription factor levels in a plant offers tremendous potential in agricultural biotechnology for modifying a plant's traits.

The present invention provides novel transcription factors useful for modifying a plant's phenotype in desirable ways, such as modifying a plant's pathogen tolerance.

SUMMARY OF THE INVENTION

In a first aspect, the invention relates to a recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29, or a complementary nucleotide sequence thereof; (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a); (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-29, or a complementary nucleotide sequence thereof; (d) a nucleotide sequence

comprising silent substitutions in a nucleotide sequence of (c); (e) a nucleotide sequence which hybridizes under stringent conditions over substantially the entire length of a nucleotide sequence of one or more of: (a), (b), (c), or (d); (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e); (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide having a biological activity that modifies a plant's pathogen tolerance; (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g); (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g); (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; and (1) a nucleotide sequence which encodes a conserved domain of a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-29. The recombinant polynucleotide may further comprise a constitutive, inducible, or tissue-active promoter operably linked to the nucleotide sequence. The invention also relates to compositions comprising at least two of the above described polynucleotides.

5

10

15

20

25

30

35

In a second aspect, the invention is an isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide described above.

In another aspect, the invention is a transgenic plant comprising one or more of the above described recombinant polynucleotides. In yet another aspect, the invention is a plant with altered expression levels of a polynucleotide described above or a plant with altered expression or activity levels of an above described polypeptide. Further, the invention may be a plant lacking a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29.

The plant may be a soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, or vegetable brassicas plant.

In a further aspect, the invention relates to a cloning or expression vector comprising the isolated or recombinant polynucleotide described above or cells comprising the cloning or expression vector.

In yet a further aspect, the invention relates to a composition produced by incubating a polynucleotide of the invention with a nuclease, a restriction enzyme, a polymerase; a polymerase and a primer; a cloning vector, or with a cell.

Furthermore, the invention relates to a method for producing a plant having improved pathogen tolerance. The method comprises altering the expression of an isolated or recombinant polynucleotide of the invention or altering the expression or activity of a polypeptide of the invention in a plant to produce a modified plant, and selecting the modified plant for modified pathogen tolerance.

5

10

15

20

25

30

35

In another aspect, the invention relates to a method of identifying a factor that is modulated by or interacts with a polypeptide encoded by a polynucleotide of the invention. The method comprises expressing a polypeptide encoded by the polynucleotide in a plant; and identifying at least one factor that is modulated by or interacts with the polypeptide. In one embodiment the method for identifying modulating or interacting factors is by detecting binding by the polypeptide to a promoter sequence, or by detecting interactions between an additional protein and the polypeptide in a yeast two hybrid system, or by detecting expression of a factor by hybridization to a microarray, subtractive hybridization or differential display.

In yet another aspect, the invention is a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest. The method comprises placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of the invention and monitoring one or more of the expression level of the polynucleotide in the plant, the expression level of the polypeptide in the plant, and modulation of an activity of the polypeptide in the plant.

In yet another aspect, the invention relates to an integrated system, computer or computer readable medium comprising one or more character strings corresponding to a polynucleotide of the invention, or to a polypeptide encoded by the polynucleotide. The integrated system, computer or computer readable medium may comprise a link between one or more sequence strings to a modified plant pathogen tolerance phenotype.

In yet another aspect, the invention is a method for identifying a sequence similar or homologous to one or more polynucleotides of the invention, or one or more polypeptides encoded by the polynucleotides. The method comprises providing a sequence database; and, querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.

The method may further comprise of linking the one or more of the polynucleotides of the invention, or encoded polypeptides, to a modified plant pathogen tolerance phenotype.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a table of exemplary polynucleotide and polypeptide sequences of the invention. The table includes from left to right for each sequence: the SEQ ID No., the internal code reference number (GID), whether the sequence is a polynucleotide or polypeptide sequence, and identification of any conserved domains for the polypeptide sequences.

Figure 2 provides a table of exemplary sequences that are homologous to other sequences provided in the Sequence Listing and that are derived from Arabidopsis thaliana. The table includes from left to right: the SEQ ID No., the internal code reference number (GID), identification of the homologous sequence, whether the sequence is a polynucleotide or polypeptide sequence, and identification of any conserved domains for the polypeptide sequences.

10

15

20

25

30

35

Figure 3 provides a table of exemplary sequences that are homologous to the sequences provided in Figures 1 and 2 and that are derived from plants other than *Arabidopsis thaliana*. The table includes from left to right: the SEQ ID No., the internal code reference number (GID), the unique GenBank sequence ID No. (NID), the probability that the comparison was generated by chance (P-value), and the species from which the homologous gene was identified.

DETAILED DESCRIPTION

The present invention relates to polynucleotides and polypeptides, e.g. for modifying phenotypes of plants.

In particular, the polynucleotides or polypeptides are useful for modifying traits associated with a plant's pathogen tolerance when the expression levels of the polynucleotides or expression levels or activity levels of the polypeptides are altered. Specifically, the polynucleotides and polypeptides are useful for modifying traits associated with a plant's pathogen tolerance, such as alterations in cell wall composition, trichome number or structure, callose induction, phytoalexin induction, alterations in the cell death response, or the like. Transgenic plants employing the polynucleotides or polypeptides of the invention are more tolerant to biotrophic or necrotrophic pathogens such as fungi, bacteria, mollicutes, viruses, nematodes, parasitic higher plants or the like.

The polynucleotides of the invention encode plant transcription factors. The plant transcription factors are derived, e.g., from *Arabidopsis thaliana* and can belong, e.g., to one or more of the following transcription factor families: the AP2 (APETALA2) domain transcription factor family (Riechmann and Meyerowitz (1998) <u>J. Biol. Chem.</u> 379:633-646); the MYB transcription factor family (Martin and Paz-Ares (1997) <u>Trends Genet.</u> 13:67-73); the MADS domain transcription factor family (Riechmann and Meyerowitz (1997) <u>J. Biol.</u>

Chem. 378:1079-1101); the WRKY protein family (Ishiguro and Nakamura (1994) Mol. Gen. Genet. 244:563-571); the ankyrin-repeat protein family (Zhang et al. (1992) Plant Cell 4:1575-1588); the miscellaneous protein (MISC) family (Kim et al. (1997) Plant J. 11:1237-1251); the zinc finger protein (Z) family (Klug and Schwabe (1995) FASEB J. 9: 597-604); the homeobox (HB) protein family (Duboule (1994) Guidebook to the Homeobox Genes. Oxford University Press); the CAAT-element binding proteins (Forsburg and Guarente (1989) Genes Dev. 3:1166-1178); the squamosa promoter binding proteins (SPB) (Klein et al. (1996) Mol. Gen. Genet. 1996 250:7-16); the NAM protein family; the IAA/AUX proteins (Rouse et al. (1998) Science 279:1371-1373); the HLH/MYC protein family (Littlewood et al. (1994) Prot. Profile 1:639-709); the DNA-binding protein (DBP) family (Tucker et al. (1994) EMBO J. 13:2994-3002); the bZIP family of transcription factors (Foster et al. (1994) FASEB J. 8:192-200); the BPF-1 protein (Box P-binding factor) family (da Costa e Silva et al. (1993) Plant J. 4:125-135); and the golden protein (GLD) family (Hall et al. (1998) Plant Cell 10:925-936).

In addition to methods for modifying a plant phenotype by employing one or more polynucleotides and polypeptides of the invention described herein, the polynucleotides and polypeptides of the invention have a variety of additional uses. These uses include their use in the recombinant production (i.e, expression) of proteins; as regulators of plant gene expression, as diagnostic probes for the presence of complementary or partially complementary nucleic acids (including for detection of natural coding nucleic acids); as substrates for further reactions, e.g., mutation reactions, PCR reactions, or the like, of as substrates for cloning e.g., including digestion or ligation reactions, and for identifying exogenous or endogenous modulators of the transcription factors.

DEFINITIONS

10

15

20

25

30

35

A "polynucleotide" is a nucleic acid sequence comprising a plurality of polymerized nucleotide residues, e.g., at least about 15 consecutive polymerized nucleotide residues, optionally at least about 30 consecutive nucleotides, at least about 50 consecutive nucleotides. In many instances, a polynucleotide comprises a nucleotide sequence encoding a polypeptide (or protein) or a domain or fragment thereof. Additionally, the polynucleotide may comprise a promoter, an intron, an enhancer region, a polyadenylation site, a translation initiation site, 5' or 3' untranslated regions, a reporter gene, a selectable marker, or the like. The polynucleotide can be single stranded or double stranded DNA or RNA. The polynucleotide optionally comprises modified bases or a modified backbone. The polynucleotide can be, e.g., genomic DNA or RNA, a transcript (such as an mRNA), a cDNA, a PCR product, a cloned DNA, a synthetic DNA or RNA, or the like. The polynucleotide can comprise a sequence in either sense or antisense orientations.

A "recombinant polynucleotide" is a polynucleotide that is not in its native state, e.g., the polynucleotide comprises a nucleotide sequence not found in nature, or the polynucleotide is in a context other than that in which it is naturally found, e.g., separated from nucleotide sequences with which it typically is in proximity in nature, or adjacent (or contiguous with) nucleotide sequences with which it typically is not in proximity. For example, the sequence at issue can be cloned into a vector, or otherwise recombined with one or more additional nucleic acid.

5

10

15

20

25

30

35

An "isolated polynucleotide" is a polynucleotide whether naturally occurring or recombinant, that is present outside the cell in which it is typically found in nature, whether purified or not. Optionally, an isolated polynucleotide is subject to one or more enrichment or purification procedures, e.g., cell lysis, extraction, centrifugation, precipitation, or the like.

A "recombinant polypeptide" is a polypeptide produced by translation of a recombinant polypucleotide. An "isolated polypeptide," whether a naturally occurring or a recombinant polypeptide, is more enriched in (or out of) a cell than the polypeptide in its natural state in a wild type cell, e.g., more than about 5% enriched, more than about 10% enriched, or more than about 20%, or more than about 50%, or more, enriched, i.e., alternatively denoted: 105%, 110%, 120%, 150% or more, enriched relative to wild type standardized at 100%. Such an enrichment is not the result of a natural response of a wild type plant. Alternatively, or additionally, the isolated polypeptide is separated from other cellular components with which it is typically associated, e.g., by any of the various protein purification methods herein.

The term "transgenic plant" refers to a plant that contains genetic material, not found in a wild type plant of the same species, variety or cultivar. The genetic material may include a transgene, an insertional mutagenesis event (such as by transposon or T-DNA insertional mutagenesis), an activation tagging sequence, a mutated sequence, a homologous recombination event or a sequence modified by chimeraplasty. Typically, the foreign genetic material has been introduced into the plant by human manipulation.

A transgenic plant may contain an expression vector or cassette. The expression cassette typically comprises a polypeptide-encoding sequence operably linked (i.e., under regulatory control of) to appropriate inducible or constitutive regulatory sequences that allow for the expression of polypeptide. The expression cassette can be introduced into a plant by transformation or by breeding after transformation of a parent plant. A plant refers to a whole plant as well as to a plant part, such as seed, fruit, leaf, or root, plant tissue, plant cells or any other plant material, e.g., a plant explant, as well as to progeny thereof, and to *in vitro* systems that mimic biochemical or cellular components or processes in a cell.

The phrase "ectopically expression or altered expression" in reference to a polynucleotide indicates that the pattern of expression in, e.g., a transgenic plant or plant

6

tissue, is different from the expression pattern in a wild type plant or a reference plant of the same species. For example, the polynucleotide or polypeptide is expressed in a cell or tissue type other than a cell or tissue type in which the sequence is expressed in the wild type plant, or by expression at a time other than at the time the sequence is expressed in the wild type plant, or by a response to different inducible agents, such as hormones or environmental signals, or at different expression levels (either higher or lower) compared with those found in a wild type plant. The term also refers to altered expression patterns that are produced by lowering the levels of expression to below the detection level or completely abolishing expression. The resulting expression pattern can be transient or stable, constitutive or inducible. In reference to a polypeptide, the term "ectopic expression or altered expression" further may relate to altered activity levels resulting from the interactions of the polypeptides with exogenous or endogenous modulators or from interactions with factors or as a result of the chemical modification of the polypeptides.

10

15

20

25

30

35

The term "fragment" or "domain," with respect to a polypeptide, refers to a subsequence of the polypeptide. In some cases, the fragment or domain, is a subsequence of the polypeptide which performs at least one biological function of the intact polypeptide in substantially the same manner, or to a similar extent, as does the intact polypeptide. For example, a polypeptide fragment can comprise a recognizable structural motif or functional domain such as a DNA binding domain that binds to a DNA promoter region, an activation domain or a domain for protein-protein interactions. Fragments can vary in size from as few as 6 amino acids to the full length of the intact polypeptide, but are preferably at least about 30 amino acids in length and more preferably at least about 60 amino acids in length. In reference to a nucleotide sequence, "a fragment" refers to any subsequence of a polynucleotide, typically, of at least consecutive about 15 nucleotides, preferably at least about 30 nucleotides, more preferably at least about 50, of any of the sequences provided herein.

The term "trait" refers to a physiological, morphological, biochemical or physical characteristic of a plant or particular plant material or cell. In some instances, this characteristic is visible to the human eye, such as seed or plant size, or can be measured by available biochemical techniques, such as the protein, starch or oil content of seed or leaves or by the observation of the expression level of genes, e.g., by employing Northern analysis, RT-PCR, microarray gene expression assays or reporter gene expression systems, or by agricultural observations such as stress tolerance, yield or pathogen tolerance.

"Trait modification" refers to a detectable difference in a characteristic in a plant ectopically expressing a polynucleotide or polypeptide of the present invention relative to a plant not doing so, such as a wild type plant. In some cases, the trait modification can be evaluated quantitatively. For example, the trait modification can entail at least about a 2%

increase or decrease in an observed trait (difference), at least a 5% difference, at least about a 10% difference, at least about a 20% difference, at least about a 30%, at least about a 50%, at least about a 70%, or at least about a 100%, or an even greater difference. It is known that there can be a natural variation in the modified trait. Therefore, the trait modification observed entails a change of the normal distribution of the trait in the plants compared with the distribution observed in wild type plant.

5

10

15

20

25

30

35

Trait modifications of particular interest include those to seed (such as embryo or endosperm), fruit, root, flower, leaf, stem, shoot, seedling or the like, including: enhanced tolerance to environmental conditions including freezing, chilling, heat, drought, water saturation, radiation and ozone; improved tolerance to microbial, fungal or viral diseases; improved tolerance to pest infestations, including nematodes, mollicutes, parasitic higher plants or the like; decreased herbicide sensitivity; improved tolerance of heavy metals or enhanced ability to take up heavy metals; improved growth under poor photoconditions (e.g., low light and/or short day length), or changes in expression levels of genes of interest. Other phenotype that can be modified relate to the production of plant metabolites, such as variations in the production of taxol, tocopherol, tocotrienol, sterols, phytosterols, vitamins, wax monomers, anti-oxidants, amino acids, lignins, cellulose, tannins, prenyllipids (such as chlorophylls and carotenoids), glucosinolates, and terpenoids, enhanced or compositionally altered protein or oil production (especially in seeds), or modified sugar (insoluble or soluble) and/or starch composition. Physical plant characteristics that can be modified include cell development (such as the number of trichomes), fruit and seed size and number, yields of plant parts such as stems, leaves and roots, the stability of the seeds during storage, characteristics of the seed pod (e.g., susceptibility to shattering), root hair length and quantity, internode distances, or the quality of seed coat. Plant growth characteristics that can be modified include growth rate, germination rate of seeds, vigor of plants and seedlings, leaf and flower senescence, male sterility, apomixis, flowering time, flower abscission, rate of nitrogen uptake, biomass or transpiration characteristics, as well as plant architecture characteristics such as apical dominance, branching patterns, number of organs, organ identity, organ shape or size.

POLYPEPTIDES AND POLYNUCLEOTIDES OF THE INVENTION

The present invention provides, among other things, transcription factors (TFs), and transcription factor homologue polypeptides, and isolated or recombinant polynucleotides encoding the polypeptides. These polypeptides and polynucleotides may be employed to modify a plant's pathogen tolerance.

Exemplary polynucleotides encoding the polypeptides of the invention were identified in the *Arabidopsis thaliana* GenBank database using publicly available sequence

analysis programs and parameters. Sequences initially identified were then further characterized to identify sequences comprising specified sequence strings corresponding to sequence motifs present in families of known transcription factors. Polynucleotide sequences meeting such criteria were confirmed as transcription factors.

Additional polynucleotides of the invention were identified by screening Arabidopsis thaliana and/or other plant cDNA libraries with probes corresponding to known transcription factors under low stringency hybridization conditions. Additional sequences, including full length coding sequences were subsequently recovered by the rapid amplification of cDNA ends (RACE) procedure, using a commercially available kit according to the manufacturer's instructions. Where necessary, multiple rounds of RACE are performed to isolate 5' and 3' ends. The full length cDNA was then recovered by a routine end-to-end polymerase chain reaction (PCR) using primers specific to the isolated 5' and 3' ends. Exemplary sequences are provided in the Sequence Listing.

The polynucleotides of the invention were ectopically expressed in overexpressor or knockout plants and changes in the pathogen tolerance of the plants was observed. Therefore, the polynucleotides and polypeptides can be employed to improve the pathogen resistance of plants.

Making polynucleotides

5

10

15

20

25

30

35

The polynucleotides of the invention include sequences that encode transcription factors and transcription factor homologue polypeptides and sequences complementary thereto, as well as unique fragments of coding sequence, or sequence complementary thereto. Such polynucleotides can be, e.g., DNA or RNA, e.g., mRNA, cRNA, synthetic RNA, genomic DNA, cDNA synthetic DNA, oligonucleotides, etc. The polynucleotides are either double-stranded or single-stranded, and include either, or both sense (i.e., coding) sequences and antisense (i.e., non-coding, complementary) sequences. The polynucleotides include the coding sequence of a transcription factor, or transcription factor homologue polypeptide, in isolation, in combination with additional coding sequences (e.g., a purification tag, a localization signal, as a fusion-protein, as a pre-protein, or the like), in combination with non-coding sequences (e.g., introns or inteins, regulatory elements such as promoters, enhancers, terminators, and the like), and/or in a vector or host environment in which the polynucleotide encoding a transcription factor or transcription factor homologue polypeptide is an endogenous or exogenous gene.

A variety of methods exist for producing the polynucleotides of the invention. Procedures for identifying and isolating DNA clones are well known to those of skill in the art, and are described in, e.g., Berger and Kimmel, <u>Guide to Molecular Cloning Techniques</u>, <u>Methods in Enzymology</u> volume 152 Academic Press, Inc., San Diego, CA ("Berger");

Sambrook et al., Molecular Cloning - A Laboratory Manual (2nd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989 ("Sambrook") and Current Protocols in Molecular Biology, F.M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2000) ("Ausubel").

Alternatively, polynucleotides of the invention, can be produced by a variety of in vitro amplification methods adapted to the present invention by appropriate selection of specific or degenerate primers. Examples of protocols sufficient to direct persons of skill through in vitro amplification methods, including the polymerase chain reaction (PCR) the ligase chain reaction (LCR), Qbeta-replicase amplification and other RNA polymerase mediated techniques (e.g., NASBA), e.g., for the production of the homologous nucleic acids of the invention are found in Berger, Sambrook, and Ausubel, as well as Mullis et al., (1987) PCR Protocols A Guide to Methods and Applications (Innis et al. eds) Academic Press Inc. San Diego, CA (1990) (Innis). Improved methods for cloning in vitro amplified nucleic acids are described in Wallace et al., U.S. Pat. No. 5,426,039. Improved methods for amplifying large nucleic acids by PCR are summarized in Cheng et al. (1994) Nature 369: 684-685 and the references cited therein, in which PCR amplicons of up to 40kb are generated. One of skill will appreciate that essentially any RNA can be converted into a double stranded DNA suitable for restriction digestion, PCR expansion and sequencing using reverse transcriptase and a polymerase. See, e.g., Ausubel, Sambrook and Berger, all supra.

Alternatively, polynucleotides and oligonucleotides of the invention can be assembled from fragments produced by solid-phase synthesis methods. Typically, fragments of up to approximately 100 bases are individually synthesized and then enzymatically or chemically ligated to produce a desired sequence, e.g., a polynucleotide encoding all or part of a transcription factor. For example, chemical synthesis using the phosphoramidite method is described, e.g., by Beaucage et al. (1981) Tetrahedron Letters 22:1859-69; and Matthes et al. (1984) EMBO J. 3:801-5. According to such methods, oligonucleotides are synthesized, purified, annealed to their complementary strand, ligated and then optionally cloned into suitable vectors. And if so desired, the polynucleotides and polypeptides of the invention can be custom ordered from any of a number of commercial suppliers.

HOMOLOGOUS SEQUENCES

5

10

15

20

25

30

35

Sequences homologous, i.e., that share significant sequence identity or similarity, to those provided in the Sequence Listing, derived from Arabidopsis thaliana or from other plants of choice are also an aspect of the invention. Homologous sequences can be derived from any plant including monocots and dicots and in particular agriculturally important plant species, including but not limited to, crops such as soybean, wheat, corn,

potato, cotton, rice, oilseed rape (including canola), sunflower, alfalfa, sugarcane and turf; or fruits and vegetables, such as banana, blackberry, blueberry, strawberry, and raspberry, cantaloupe, carrot, cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits (such as apple, peach, pear, cherry and plum) and vegetable brassicas (such as broccoli, cabbage, cauliflower, brussel sprouts and kohlrabi). Other crops, fruits and vegetables whose phenotype can be changed include barley, rye, millet, sorghum, currant, avocado, citrus fruits such as oranges, lemons, grapefruit and tangerines, artichoke, cherries, nuts such as the walnut and peanut, endive, leek, roots, such as arrowroot, beet, cassava, turnip, radish, yam, and sweet potato, and beans. The homologous sequences may also be derived from woody species, such pine, poplar and eucalyptus.

10

15

20

25

30

35

Transcription factors that are homologous to the listed sequences will typically share at least about 31% amino acid sequence identity. More closely related transcription factors can share at least about 50%, about 60%, about 65%, about 70%, about 75% or about 80% or about 90% or about 95% or about 98% or more sequence identity with the listed sequences. Factors that are most closely related to the listed sequences share, e.g., at least about 85%, about 90% or about 95% or more % sequence identity to the listed sequences. At the nucleotide level, the sequences will typically share at least about 40% nucleotide sequence identity, preferably at least about 50%, about 60%, about 70% or about 80% sequence identity, and more preferably about 85%, about 90%, about 95% or about 97% or more sequence identity to one or more of the listed sequences. The degeneracy of the genetic code enables major variations in the nucleotide sequence of a polynucleotide while maintaining the amino acid sequence of the encoded protein. Conserved domains within a transcription factor family may exhibit a higher degree of sequence homology, such as at least 65% sequence identity including conservative substitutions, and preferably at least 80% sequence identity.

Identifying Nucleic Acids by Hybridization

Polynucleotides homologous to the sequences illustrated in the Sequence Listing can be identified, e.g., by hybridization to each other under stringent or under highly stringent conditions. Single stranded polynucleotides hybridize when they associate based on a variety of well characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. The stringency of a hybridization reflects the degree of sequence identity of the nucleic acids involved, such that the higher the stringency, the more similar are the two polynucleotide strands. Stringency is influenced by a variety of factors, including temperature, salt concentration and composition, organic and non-organic additives, solvents, etc. present in both the hybridization and wash solutions and incubations (and number), as described in more detail in the references cited above.

An example of stringent hybridization conditions for hybridization of complementary nucleic acids which have more than 100 complementary residues on a filter in a Southern or northern blot is about 5°C to 20°C lower than the thermal melting point (Tm) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Nucleic acid molecules that hybridize under stringent conditions will typically hybridize to a probe based on either the entire cDNA or selected portions, e.g., to a unique subsequence, of the cDNA under wash conditions of 0.2x SSC to 2.0 x SSC, 0.1% SDS at 50-65° C, for example 0.2 x SSC, 0.1% SDS at 65° C. For identification of less closely related homologues washes can be performed at a lower temperature, e.g., 50° C. In general, stringency is increased by raising the wash temperature and/or decreasing the concentration of SSC.

As another example, stringent conditions can be selected such that an oligonucleotide that is perfectly complementary to the coding oligonucleotide hybridizes to the coding oligonucleotide with at least about a 5-10x higher signal to noise ratio than the ratio for hybridization of the perfectly complementary oligonucleotide to a nucleic acid encoding a transcription factor known as of the filing date of the application. Conditions can be selected such that a higher signal to noise ratio is observed in the particular assay which is used, e.g., about 15x, 25x, 35x, 50x or more. Accordingly, the subject nucleic acid hybridizes to the unique coding oligonucleotide with at least a 2x higher signal to noise ratio as compared to hybridization of the coding oligonucleotide to a nucleic acid encoding known polypeptide. Again, higher signal to noise ratios can be selected, e.g., about 5x, 10x, 25x, 35x, 50x or more. The particular signal will depend on the label used in the relevant assay, e.g., a fluorescent label, a colorimetric label, a radio active label, or the like.

Alternatively, transcription factor homologue polypeptides can be obtained by screening an expression library using antibodies specific for one or more transcription factors. With the provision herein of the disclosed transcription factor, and transcription factor homologue nucleic acid sequences, the encoded polypeptide(s) can be expressed and purified in a heterologous expression system (e.g., E. coli) and used to raise antibodies (monoclonal or polyclonal) specific for the polypeptide(s) in question. Antibodies can also be raised against synthetic peptides derived from transcription factor, or transcription factor homologue, amino acid sequences. Methods of raising antibodies are well known in the art and are described in Harlow and Lane (1988) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York. Such antibodies can then be used to screen an expression library produced from the plant from which it is desired to clone additional transcription

factor homologues, using the methods described above. The selected cDNAs can be confirmed by sequencing and enzymatic activity.

SEQUENCE VARIATIONS

5

10

It will readily be appreciated by those of skill in the art, that any of a variety of polynucleotide sequences are capable of encoding the transcription factors and transcription factor homologue polypeptides of the invention. Due to the degeneracy of the genetic code, many different polynucleotides can encode identical and/or substantially similar polypeptides in addition to those sequences illustrated in the Sequence Listing.

For example, Table 1 illustrates, e.g., that the codons AGC, AGT, TCA, TCC, TCG, and TCT all encode the same amino acid: serine. Accordingly, at each position in the sequence where there is a codon encoding serine, any of the above trinucleotide sequences can be used without altering the encoded polypeptide.

Table 1

Amino acids	•		Codon			<u>.</u>		
Alanine	Ala	A	GCA	GCC	GCG	GCU		
Cysteine	Cys	С	TGC	TGT				
Aspartic acid	Asp	D	GAC	GAT				
Glutamic acid	Glu	E	GAA	GAG			•	
Phenylalanine	Phe	F	TTC	TTT				
Glycine	Gly	G	GGA	GGC	GGG	GGT		
Histidine	His	H	CAC	CAT				
Isoleucine	Ile	I	ATA	ATC	ATT			
Lysine	Lys	K	AAA -	AAG				
Leucine	Leu	L	TTA	TTG	CTA	CTC	CTG	CTT
Methionine	Met	M	ATG		•			
Asparagine	Asn	N	AAC	AAT			•	
Proline	Pro	P	CCA	CCC	CCG	CCT		
Glutamine	Gln	Q	CAA	CAG				
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGT
Serine	Ser	S	AGC	AGT	TCA	TCC	TCG	TCT
Threonine	Thr	T	ACA	ACC	ACG	ACT]
Valine	Val	V	GTA	GTC	GTG	GTT		Ì
Tryptophan	Тгр	W	TGG					ĺ
Tyrosine	Tyr	Y	TAC	TAT				

15

20

Sequence alterations that do not change the amino acid sequence encoded by the polynucleotide are termed "silent" variations. With the exception of the codons ATG and TGG, encoding methionine and tryptophan, respectively, any of the possible codons for the same amino acid can be substituted by a variety of techniques, e.g., site-directed mutagenesis, available in the art. Accordingly, any and all such variations of a sequence selected from the above table are a feature of the invention.

In addition to silent variations, other conservative variations that alter one, or a few amino acids in the encoded polypeptide, can be made without altering the function of the polypeptide, these conservative variants are, likewise, a feature of the invention.

For example, substitutions, deletions and insertions introduced into the sequences provided in the Sequence Listing are also envisioned by the invention. Such sequence modifications can be engineered into a sequence by site-directed mutagenesis (Wu (ed.) Meth. Enzymol. (1993) vol. 217, Academic Press) or the other methods noted below. Amino acid substitutions are typically of single residues; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. In preferred embodiments, deletions or insertions are made in adjacent pairs, e.g., a deletion of two residues or insertion of two residues. Substitutions, deletions, insertions or any combination thereof can be combined to arrive at a sequence. The mutations that are made in the polynucleotide encoding the transcription factor should not place the sequence out of reading frame and should not create complementary regions that could produce secondary mRNA structure. Preferably, the polypeptide encoded by the DNA performs the desired function.

Conservative substitutions are those in which at least one residue in the amino acid sequence has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the Table 2 when it is desired to maintain the activity of the protein. Table 2 shows amino acids which can be substituted for an amino acid in a protein and which are typically regarded as conservative substitutions.

25

5

10

15

20

30

35

Table 2

Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln; His
Asp	Glu
Gln	Asn
Cys	Ser
Glu	Asp
Gly	Pro
His	Asn; Gln
Пе	Leu, Val
Leu	Пе; Val
Lys	Arg; Gln
Met	Leu; Ile
Phe	Met; Leu; Tyr
Ser	Thr; Gly
Thr	Ser;Val
Ттр	Tyr
Тут	Trp; Phe
Val	Ile; Leu

Substitutions that are less conservative than those in Table 2 can be selected by picking residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in protein properties will be those in which (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

5

10

FURTHER MODIFYING SEQUENCES OF THE INVENTION—MUTATION/ FORCED EVOLUTION

In addition to generating silent or conservative substitutions as noted, above, the present invention optionally includes methods of modifying the sequences of the Sequence Listing. In the methods, nucleic acid or protein modification methods are used to alter the given sequences to produce new sequences and/or to chemically or enzymatically modify given sequences to change the properties of the nucleic acids or proteins.

5

10

15

20

. 30

35

Thus, in one embodiment, given nucleic acid sequences are modified, e.g., according to standard mutagenesis or artificial evolution methods to produce modified sequences. For example, Ausubel, *supra*, provides additional details on mutagenesis methods. Artificial forced evolution methods are described, e.g., by Stemmer (1994) Nature 370:389-391, and Stemmer (1994) Proc. Natl. Acad. Sci. USA 91:10747-10751. Many other mutation and evolution methods are also available and expected to be within the skill of the practitioner.

Similarly, chemical or enzymatic alteration of expressed nucleic acids and polypeptides can be performed by standard methods. For example, sequence can be modified by addition of lipids, sugars, peptides, organic or inorganic compounds, by the inclusion of modified nucleotides or amino acids, or the like. For example, protein modification techniques are illustrated in Ausubel, *supra*. Further details on chemical and enzymatic modifications can be found herein. These modification methods can be used to modify any given sequence, or to modify any sequence produced by the various mutation and artificial evolution modification methods noted herein.

Accordingly, the invention provides for modification of any given nucleic acid by mutation, evolution, chemical or enzymatic modification, or other available methods, as well as for the products produced by practicing such methods, e.g., using the sequences herein as a starting substrate for the various modification approaches.

For example, optimized coding sequence containing codons preferred by a particular prokaryotic or eukaryotic host can be used e.g., to increase the rate of translation or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, as compared with transcripts produced using a non-optimized sequence. Translation stop codons can also be modified to reflect host preference. For example, preferred stop codons for S. cerevisiae and mammals are TAA and TGA, respectively. The preferred stop codon for monocotyledonous plants is TGA, whereas insects and E. coli prefer to use TAA as the stop codon.

The polynucleotide sequences of the present invention can also be engineered in order to alter a coding sequence for a variety of reasons, including but not limited to, alterations which modify the sequence to facilitate cloning, processing and/or expression of

the gene product. For example, alterations are optionally introduced using techniques which are well known in the art, e.g., site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to introduce splice sites, etc.

Furthermore, a fragment or domain derived from any of the polypeptides of the invention can be combined with domains derived from other transcription factors or synthetic domains to modify the biological activity of a transcription factor. For instance, a DNA binding domain derived from a transcription factor of the invention can be combined with the activation domain of another transcription factor or with a synthetic activation domain. A transcription activation domain assists in initiating transcription from a DNA binding site. Examples include the transcription activation region of VP16 or GAL4 (Moore et al. (1998) Proc. Natl. Acad. Sci. USA 95: 376-381; and Aoyama et al. (1995) Plant Cell 7:1773-1785), peptides derived from bacterial sequences (Ma and Ptashne (1987) Cell 51; 113-119) and synthetic peptides (Giniger and Ptashne, (1987) Nature 330:670-672).

EXPRESSION AND MODIFICATION OF POLYPEPTIDES

5

10

15

20

25

30

35

Typically, polynucleotide sequences of the invention are incorporated into recombinant DNA (or RNA) molecules that direct expression of polypeptides of the invention in appropriate host cells, transgenic plants, in vitro translation systems, or the like. Due to the inherent degeneracy of the genetic code, nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence can be substituted for any listed sequence to provide for cloning and expressing the relevant homologue.

Vectors, Promoters and Expression Systems

The present invention includes recombinant constructs comprising one or more of the nucleic acid sequences herein. The constructs typically comprise a vector, such as a plasmid, a cosmid, a phage, a virus (e.g., a plant virus), a bacterial artificial chromosome (BAC), a yeast artificial chromosome (YAC), or the like, into which a nucleic acid sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available.

General texts which describe molecular biological techniques useful herein, including the use and production of vectors, promoters and many other relevant topics, include Berger, Sambrook and Ausubel, *supra*. Any of the identified sequences can be incorporated into a cassette or vector, e.g., for expression in plants. A number of expression vectors suitable for stable transformation of plant cells or for the establishment of transgenic plants have been described including those described in Weissbach and Weissbach, (1989) Methods for Plant Molecular Biology, Academic Press, and Gelvin et al., (1990) Plant

Molecular Biology Manual, Kluwer Academic Publishers. Specific examples include those derived from a Ti plasmid of Agrobacterium tumefaciens, as well as those disclosed by Herrera-Estrella et al. (1983) Nature 303: 209, Bevan (1984) Nucl Acid Res. 12: 8711-8721, Klee (1985) Bio/Technology 3: 637-642, for dicotyledonous plants.

5

15

20

25

30

Alternatively, non-Ti vectors can be used to transfer the DNA into monocotyledonous plants and cells by using free DNA delivery techniques. Such methods can involve, for example, the use of liposomes, electroporation, microprojectile bombardment, silicon carbide whiskers, and viruses. By using these methods transgenic plants such as wheat, rice (Christou (1991) Bio/Technology 9: 957-962) and corn (Gordon-10 Kamm (1990) Plant Cell 2: 603-618) can be produced. An immature embryo can also be a good target tissue for monocots for direct DNA delivery techniques by using the particle gun (Weeks et al. (1993) Plant Physiol 102: 1077-1084; Vasil (1993) Bio/Technology 10: 667-674; Wan and Lemeaux (1994) Plant Physiol 104: 37-48, and for Agrobacterium-mediated DNA transfer (Ishida et al. (1996) Nature Biotech 14: 745-750).

Typically, plant transformation vectors include one or more cloned plant coding sequence (genomic or cDNA) under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant transformation vectors typically also contain a promoter (e.g., a regulatory region controlling inducible or constitutive, environmentally-or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, an RNA processing signal (such as intron splice sites), a transcription termination site, and/or a polyadenylation signal.

Examples of constitutive plant promoters which can be useful for expressing the TF sequence include: the cauliflower mosaic virus (CaMV) 35S promoter, which confers constitutive, high-level expression in most plant tissues (see, e.g., Odel et al. (1985) Nature 313:810); the nopaline synthase promoter (An et al. (1988) Plant Physiol 88:547); and the octopine synthase promoter (Fromm et al. (1989) Plant Cell 1: 977).

A variety of plant gene promoters that regulate gene expression in response to environmental, hormonal, chemical, developmental signals, and in a tissue-active manner can be used for expression of a TF sequence in plants. Choice of a promoter is based largely on the phenotype of interest and is determined by such factors as tissue (e.g., seed, fruit, root, pollen, vascular tissue, flower, carpel, etc.), inducibility (e.g., in response to wounding, heat, cold, drought, light, pathogens, etc.), timing, developmental stage, and the like. Numerous known promoters have been characterized and can favorable be employed to promote expression of a polynucleotide of the invention in a transgenic plant or cell of interest. For example, tissue specific promoters include: seed-specific promoters (such as the napin, phaseolin or DC3 promoter described in US Pat. No. 5,773,697), fruit-specific promoters that are active during fruit ripening (such as the dru 1 promoter (US Pat. No. 5,783,393), or the

2A11 promoter (US Pat. No. 4,943,674) and the tomato polygalacturonase promoter (Bird et al. (1988) Plant Mol Biol 11:651), root-specific promoters, such as those disclosed in US Patent Nos. 5,618,988, 5,837,848 and 5,905,186, pollen-active promoters such as PTA29, PTA26 and PTA13 (US Pat. No. 5,792,929), promoters active in vascular tissue (Ringli and Keller (1998) Plant Mol Biol 37:977-988), flower-specific (Kaiser et al, (1995) Plant Mol Biol 28:231-243), pollen (Baerson et al. (1994) Plant Mol Biol 26:1947-1959), carpels (Ohl et al. (1990) Plant Cell 2:837-848), pollen and ovules (Baerson et al. (1993) Plant Mol Biol 22:255-267), auxin-inducible promoters (such as that described in van der Kop et al. (1999) Plant Mol Biol 39:979-990 or Baumann et al. (1999) Plant Cell 11:323-334), cytokinininducible promoter (Guevara-Garcia (1998) Plant Mol Biol 38:743-753), promoters responsive to gibberellin (Shi et al. (1998) Plant Mol Biol 38:1053-1060, Willmott et al. (1998) 38:817-825) and the like. Additional promoters are those that elicit expression in response to heat (Ainley et al. (1993) Plant Mol Biol 22: 13-23), light (e.g., the pea rbcS-3A promoter, Kuhlemeier et al. (1989) Plant Cell 1:471, and the maize rbcS promoter, Schaffner and Sheen (1991) Plant Cell 3: 997); wounding (e.g., wunI, Siebertz et al. (1989) Plant Cell 1: 961); pathogens (such as the PR-1 promoter described in Buchel et al. (1999) Plant Mol. Biol. 40:387-396, and the PDF1.2 promoter described in Manners et al. (1998) Plant Mol. Biol. 38:1071-80), and chemicals such as methyl jasmonate or salicylic acid (Gatz et al. (1997) Plant Mol Biol 48: 89-108). In addition, the timing of the expression can be controlled by using promoters such as those acting at senescence (An and Amazon (1995) Science 270: 1986-1988); or late seed development (Odell et al. (1994) Plant Physiol 106:447-458).

10

15

20

25

30

35

Plant expression vectors can also include RNA processing signals that can be positioned within, upstream or downstream of the coding sequence. In addition, the expression vectors can include additional regulatory sequences from the 3'-untranslated region of plant genes, e.g., a 3' terminator region to increase mRNA stability of the mRNA, such as the PI-II terminator region of potato or the octopine or nopaline synthase 3' terminator regions.

Additional Expression Elements

Specific initiation signals can aid in efficient translation of coding sequences. These signals can include, e.g., the ATG initiation codon and adjacent sequences. In cases where a coding sequence, its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only coding sequence (e.g., a mature protein coding sequence), or a portion thereof, is inserted, exogenous transcriptional control signals including the ATG initiation codon can be separately provided. The initiation codon is provided in the correct reading frame to facilitate transcription. Exogenous transcriptional elements and initiation

codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers appropriate to the cell system in use.

Expression Hosts

10

15

20

25

30

35

The present invention also relates to host cells which are transduced with vectors of the invention, and the production of polypeptides of the invention (including fragments thereof) by recombinant techniques. Host cells are genetically engineered (i.e, nucleic acids are introduced, e.g., transduced, transformed or transfected) with the vectors of this invention, which may be, for example, a cloning vector or an expression vector comprising the relevant nucleic acids herein. The vector is optionally a plasmid, a viral particle, a phage, a naked nucleic acids, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants, or amplifying the relevant gene. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art and in the references cited herein, including, Sambrook and Ausubel.

The host cell can be a eukaryotic cell, such as a yeast cell, or a plant cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Plant protoplasts are also suitable for some applications. For example, the DNA fragments are introduced into plant tissues, cultured plant cells or plant protoplasts by standard methods including electroporation (Fromm et al., (1985) Proc. Natl. Acad. Sci. USA 82, 5824, infection by viral vectors such as cauliflower mosaic virus (CaMV) (Hohn et al., (1982) Molecular Biology of Plant Tumors, (Academic Press, New York) pp. 549-560; US 4,407,956), high velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface (Klein et al., (1987) Nature 327, 70-73), use of pollen as vector (WO 85/01856), or use of Agrobacterium tumefaciens or A. rhizogenes carrying a T-DNA plasmid in which DNA fragments are cloned. The T-DNA plasmid is transmitted to plant cells upon infection by Agrobacterium tumefaciens, and a portion is stably integrated into the plant genome (Horsch et al. (1984) Science 233:496-498; Fraley et al. (1983) Proc. Natl. Acad. Sci. USA 80, 4803).

The cell can include a nucleic acid of the invention which encodes a polypeptide, wherein the cells expresses a polypeptide of the invention. The cell can also include vector sequences, or the like. Furthermore, cells and transgenic plants which include any polypeptide or nucleic acid above or throughout this specification, e.g., produced by transduction of a vector of the invention, are an additional feature of the invention.

For long-term, high-yield production of recombinant proteins, stable expression can be used. Host cells transformed with a nucleotide sequence encoding a polypeptide of the invention are optionally cultured under conditions suitable for the

expression and recovery of the encoded protein from cell culture. The protein or fragment thereof produced by a recombinant cell may be secreted, membrane-bound, or contained intracellularly, depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides encoding mature proteins of the invention can be designed with signal sequences which direct secretion of the mature polypeptides through a prokaryotic or eukaryotic cell membrane.

Modified Amino Acids

10

15

25

30

35

Polypeptides of the invention may contain one or more modified amino acids. The presence of modified amino acids may be advantageous in, for example, increasing polypeptide half-life, reducing polypeptide antigenicity or toxicity, increasing polypeptide storage stability, or the like. Amino acid(s) are modified, for example, co-translationally or post-translationally during recombinant production or modified by synthetic or chemical means.

Non-limiting examples of a modified amino acid include incorporation or other use of acetylated amino acids, glycosylated amino acids, sulfated amino acids, prenylated (e.g., farnesylated, geranylgeranylated) amino acids, PEG modified (e.g., "PEGylated") amino acids, biotinylated amino acids, carboxylated amino acids, phosphorylated amino acids, etc. References adequate to guide one of skill in the modification of amino acids are replete throughout the literature.

20 IDENTIFICATION OF ADDITIONAL FACTORS

A transcription factor provided by the present invention can also be used to identify additional endogenous or exogenous molecules that can affect a phentoype or trait of interest. On the one hand, such molecules include organic (small or large molecules) and/or inorganic compounds that affect expression of (i.e., regulate) a particular transcription factor. Alternatively, such molecules include endogenous molecules that are acted upon either at a transcriptional level by a transcription factor of the invention to modify a phenotype as desired. For example, the transcription factors can be employed to identify one or more downstream gene with which is subject to a regulatory effect of the transcription factor. In one approach, a transcription factor or transcription factor homologue of the invention is expressed in a host cell, e.g, a transgenic plant cell, tissue or explant, and expression products, either RNA or protein, of likely or random targets are monitored, e.g., by hybridization to a microarray of nucleic acid probes corresponding to genes expressed in a tissue or cell type of interest, by two-dimensional gel electrophoresis of protein products, or by any other method known in the art for assessing expression of gene products at the level of RNA or protein. Alternatively, a transcription factor of the invention can be used to identify promoter sequences (i.e., binding sites) involved in the regulation of a downstream target. After

identifying a promoter sequence, interactions between the transcription factor and the promoter sequence can be modified by changing specific nucleotides in the promoter sequence or specific amino acids in the transcription factor that interact with the promoter sequence to alter a plant trait. Typically, transcription factor DNA binding sites are identified by gel shift assays. After identifying the promoter regions, the promoter region sequences can be employed in double-stranded DNA arrays to identify molecules that affect the interactions of the transcription factors with their promoters (Bulyk et al. (1999) Nature Biotechnology 17:573-577).

The identified transcription factors are also useful to identify proteins that modify the activity of the transcription factor. Such modification can occur by covalent modification, such as by phosphorylation, or by protein-protein (homo or-heteropolymer) interactions. Any method suitable for detecting protein-protein interactions can be employed. Among the methods that can be employed are co-immunoprecipitation, cross-linking and co-purification through gradients or chromatographic columns, and the two-hybrid yeast system.

The two-hybrid system detects protein interactions in vivo and is described in Chien, et al., (1991), Proc. Natl. Acad. Sci. USA 88, 9578-9582 and is commercially available from Clontech (Palo Alto, Calif.). In such a system, plasmids are constructed that encode two hybrid proteins: one consists of the DNA-binding domain of a transcription activator protein fused to the TF polypeptide and the other consists of the transcription activator protein's activation domain fused to an unknown protein that is encoded by a cDNA that has been recombined into the plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the cDNA library are transformed into a strain of the yeast Saccharomyces cerevisiae that contains a reporter gene (e.g., lacZ) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene. Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product. Then, the library plasmids responsible for reporter gene expression are isolated and sequenced to identify the proteins encoded by the library plasmids. After identifying proteins that interact with the transcription factors, assays for compounds that interfere with the TF protein-protein interactions can be preformed.

IDENTIFICATION OF MODULATORS

10

15

20

25

30

35

In addition to the intracellular molecules described above, extracellular molecules that alter activity or expression of a transcription factor, either directly or indirectly, can be identified. For example, the methods can entail first placing a candidate molecule in contact with a plant or plant cell. The molecule can be introduced by topical administration, such as spraying or soaking of a plant, and then the molecule's effect on the

22

expression or activity of the TF polypeptide or the expression of the polynucleotide monitored. Changes in the expression of the TF polypeptide can be monitored by use of polyclonal or monoclonal antibodies, gel electrophoresis or the like. Changes in the expression of the corresponding polynucleotide sequence can be detected by use of microarrays, Northerns, quantitative PCR, or any other technique for monitoring changes in mRNA expression. These techniques are exemplified in Ausubel et al. (eds) <u>Current Protocols in Molecular Biology</u>, John Wiley & Sons (1998). Such changes in the expression levels can be correlated with modified plant traits and thus identified molecules can be useful for soaking or spraying on fruit, vegetable and grain crops to modify traits in plants.

Essentially any available composition can be tested for modulatory activity of expression or activity of any nucleic acid or polypeptide herein. Thus, available libraries of compounds such as chemicals, polypeptides, nucleic acids and the like can be tested for modulatory activity. Often, potential modulator compounds can be dissolved in aqueous or organic (e.g., DMSO-based) solutions for easy delivery to the cell or plant of interest in which the activity of the modulator is to be tested. Optionally, the assays are designed to screen large modulator composition libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays).

In one embodiment, high throughput screening methods involve providing a combinatorial library containing a large number of potential compounds (potential modulator compounds). Such "combinatorial chemical libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as target compounds.

A combinatorial chemical library can be, e.g., a collection of diverse chemical compounds generated by chemical synthesis or biological synthesis. For example, a combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (e.g., in one example, amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound of a set length). Exemplary libraries include peptide libraries, nucleic acid libraries, antibody libraries (see, e.g., Vaughn et al. (1996) Nature Biotechnology, 14(3):309-314 and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al. Science (1996) 274:1520-1522 and U.S. Patent 5,593,853), peptide nucleic acid libraries (see, e.g., U.S. Patent 5,539,083), and small organic molecule libraries (see, e.g., benzodiazepines, Baum C&EN Jan 18, page 33 (1993); isoprenoids, U.S. Patent 5,569,588; thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pyrrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337) and the like.

Preparation and screening of combinatorial or other libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Patent 5,010,175, Furka, <u>Int. J. Pept. Prot. Res.</u> 37:487-493 (1991) and Houghton et al. <u>Nature 354:84-88 (1991)</u>). Other chemistries for generating chemical diversity libraries can also be used.

5

10

15

20

25

30

35

In addition, as noted, compound screening equipment for high-throughput screening is generally available, e.g., using any of a number of well known robotic systems that have also been developed for solution phase chemistries useful in assay systems. These systems include automated workstations including an automated synthesis apparatus and robotic systems utilizing robotic arms. Any of the above devices are suitable for use with the present invention, e.g., for high-throughput screening of potential modulators. The nature and implementation of modifications to these devices (if any) so that they can operate as discussed herein will be apparent to persons skilled in the relevant art.

Indeed, entire high throughput screening systems are commercially available. These systems typically automate entire procedures including all sample and reagent pipetting, liquid dispensing, timed incubations, and final readings of the microplate in detector(s) appropriate for the assay. These configurable systems provide high throughput and rapid start up as well as a high degree of flexibility and customization. Similarly, microfluidic implementations of screening are also commercially available.

The manufacturers of such systems provide detailed protocols the various high throughput. Thus, for example, Zymark Corp. provides technical bulletins describing screening systems for detecting the modulation of gene transcription, ligand binding, and the like. The integrated systems herein, in addition to providing for sequence alignment and, optionally, synthesis of relevant nucleic acids, can include such screening apparatus to identify modulators that have an effect on one or more polynucleotides or polypeptides according to the present invention.

In some assays it is desirable to have positive controls to ensure that the components of the assays are working properly. At least two types of positive controls are appropriate. That is, known transcriptional activators or inhibitors can be incubated with cells/plants/ etc. in one sample of the assay, and the resulting increase/decrease in transcription can be detected by measuring the resulting increase in RNA/ protein expression, etc., according to the methods herein. It will be appreciated that modulators can also be combined with transcriptional activators or inhibitors to find modulators which inhibit transcriptional activation or transcriptional repression. Either expression of the nucleic acids and proteins herein or any additional nucleic acids or proteins activated by the nucleic acids or proteins herein, or both, can be monitored.

In an embodiment, the invention provides a method for identifying compositions that modulate the activity or expression of a polynucleotide or polypeptide of the invention. For example, a test compound, whether a small or large molecule, is placed in contact with a cell, plant (or plant tissue or explant), or composition comprising the polynucleotide or polypeptide of interest and a resulting effect on the cell, plant, (or tissue or explant) or composition is evaluated by monitoring, either directly or indirectly, one or more of: expression level of the polynucleotide or polypeptide, activity (or modulation of the activity) of the polynucleotide or polypeptide. In some cases, an alteration in a plant phenotype can be detected following contact of a plant (or plant cell, or tissue or explant) with the putative modulator, e.g., by modulation of expression or activity of a polynucleotide or polypeptide of the invention.

SUBSEQUENCES

10

15

20

25

30

35

Also contemplated are uses of polynucleotides, also referred to herein as oligonucleotides, typically having at least 12 bases, preferably at least 15, more preferably at least 20, 30, or 50 bases, which hybridize under at least highly stringent (or ultra-high stringent or ultra-ultra- high stringent conditions) conditions to a polynucleotide sequence described above. The polynucleotides may be used as probes, primers, sense and antisense agents, and the like, according to methods as noted *supra*.

Subsequences of the polynucleotides of the invention, including polynucleotide fragments and oligonucleotides are useful as nucleic acid probes and primers. An oligonucleotide suitable for use as a probe or primer is at least about 15 nucleotides in length, more often at least about 18 nucleotides, often at least about 21 nucleotides, frequently at least about 30 nucleotides, or about 40 nucleotides, or more in length. A nucleic acid probe is useful in hybridization protocols, e.g., to identify additional polypeptide homologues of the invention, including protocols for microarray experiments. Primers can be annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, and then extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other nucleic-acid amplification methods. See Sambrook and Ausubel, *supra*.

In addition, the invention includes an isolated or recombinant polypeptide including a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotides of the invention. For example, such polypeptides, or domains or fragments thereof, can be used as immunogens, e.g., to produce antibodies specific for the polypeptide sequence, or as probes for detecting a sequence of interest. A

subsequence can range in size from about 15 amino acids in length up to and including the full length of the polypeptide.

PRODUCTION OF TRANSGENIC PLANTS

5

10

15

20

25

30

35

Modification of Traits

The polynucleotides of the invention are favorably employed to produce transgenic plants with various traits, or characteristics, that have been modified in a desirable manner, e.g., to improve the pathogen resistance of a plant. For example, alteration of expression levels or patterns (e.g., spatial or temporal expression patterns) of one or more of the transcription factors (or transcription factor homologues) of the invention, as compared with the levels of the same protein found in a wild type plant, can be used to modify a plant's traits. An illustrative example of trait modification, improved pathogen tolerance, by altering expression levels of a particular transcription factor is described further in the Examples and the Sequence Listing.

Antisense and Cosuppression Approaches

In addition to expression of the nucleic acids of the invention as gene replacement or plant phenotype modification nucleic acids, the nucleic acids are also useful for sense and anti-sense suppression of expression, e.g., to down-regulate expression of a nucleic acid of the invention, e.g., as a further mechanism for modulating plant phenotype. That is, the nucleic acids of the invention, or subsequences or anti-sense sequences thereof, can be used to block expression of naturally occurring homologous nucleic acids. A variety of sense and anti-sense technologies are known in the art, e.g., as set forth in Lichtenstein and Nellen (1997) Antisense Technology: A Practical Approach IRL Press at Oxford University, Oxford, England. In general, sense or anti-sense sequences are introduced into a cell, where they are optionally amplified, e.g., by transcription. Such sequences include both simple oligonucleotide sequences and catalytic sequences such as ribozymes.

For example, a reduction or elimination of expression (i.e., a "knock-out") of a transcription factor or transcription factor homologue polypeptide in a transgenic plant, e.g., to modify a plant trait, can be obtained by introducing an antisense construct corresponding to the polypeptide of interest as a cDNA. For antisense suppression, the transcription factor or homologue cDNA is arranged in reverse orientation (with respect to the coding sequence) relative to the promoter sequence in the expression vector. The introduced sequence need not be the full length cDNA or gene, and need not be identical to the cDNA or gene found in the plant type to be transformed. Typically, the antisense sequence need only be capable of hybridizing to the target gene or RNA of interest. Thus, where the introduced sequence is of shorter length, a higher degree of homology to the endogenous transcription factor sequence will be needed for effective antisense suppression. While antisense sequences of various

lengths can be utilized, preferably, the introduced antisense sequence in the vector will be at least 30 nucleotides in length, and improved antisense suppression will typically be observed as the length of the antisense sequence increases. Preferably, the length of the antisense sequence in the vector will be greater than 100 nucleotides. Transcription of an antisense construct as described results in the production of RNA molecules that are the reverse complement of mRNA molecules transcribed from the endogenous transcription factor gene in the plant cell.

Suppression of endogenous transcription factor gene expression can also be achieved using a ribozyme. Ribozymes are RNA molecules that possess highly specific endoribonuclease activity. The production and use of ribozymes are disclosed in U.S. Patent No. 4,987,071 and U.S. Patent No. 5,543,508. Synthetic ribozyme sequences including antisense RNAs can be used to confer RNA cleaving activity on the antisense RNA, such that endogenous mRNA molecules that hybridize to the antisense RNA are cleaved, which in turn leads to an enhanced antisense inhibition of endogenous gene expression.

15

20

10

Vectors in which RNA encoded by a transcription factor or transcription factor homologue cDNA is over-expressed can also be used to obtain co-suppression of a corresponding endogenous gene, e.g., in the manner described in U.S. Patent No. 5,231,020 to Jorgensen. Such co-suppression (also termed sense suppression) does not require that the entire transcription factor cDNA be introduced into the plant cells, nor does it require that the introduced sequence be exactly identical to the endogenous transcription factor gene of interest. However, as with antisense suppression, the suppressive efficiency will be enhanced as specificity of hybridization is increased, e.g., as the introduced sequence is lengthened, and/or as the sequence similarity between the introduced sequence and the endogenous transcription factor gene is increased.

25

30

35

Vectors expressing an untranslatable form of the transcription factor mRNA, e.g., sequences comprising one or more stop codon, or nonsense mutation) can also be used to suppress expression of an endogenous transcription factor, thereby reducing or eliminating it's activity and modifying one or more traits. Methods for producing such constructs are described in U.S. Patent No. 5,583,021. Preferably, such constructs are made by introducing a premature stop codon into the transcription factor gene. Alternatively, a plant trait can be modified by gene silencing using double-strand RNA (Sharp (1999) Genes and Development 13: 139-141).

Another method for abolishing the expression of a gene is by insertion mutagenesis using the T-DNA of *Agrobacterium tumefaciens*. After generating the insertion mutants, the mutants can be screened to identify those containing the insertion in a transcription factor or transcription factor homologue gene. Plants containing a single

transgene insertion event at the desired gene can be crossed to generate homozygous plants for the mutation (Koncz et al. (1992) Methods in Arabidopsis Research, World Scientific).

Alternatively, a plant phenotype can be altered by eliminating an endogenous gene, such as a transcription factor or transcription factor homologue, e.g., by homologous recombination (Kempin et al. (1997) <u>Nature</u> 389:802).

5

10

15

20

25

30

35

A plant trait can also be modified by using the cre-lox system (for example, as described in US Paent No. 5,658,772). A plant genome can be modified to include first and second lox sites that are then contacted with a Cre recombinase. If the lox sites are in the same orientation, the intervening DNA sequence between the two sites is excised. If the lox sites are in the opposite orientation, the intervening sequence is inverted.

The polynucleotides and polypeptides of this invention can also be expressed in a plant in the absence of an expression cassette by manipulating the activity or expression level of the endogenous gene by other means. For example, by ectopically expressing a gene by T-DNA activation tagging (Ichikawa et al. (1997) Nature 390 698-701; Kakimoto et al. (1996) Science 274: 982-985). This method entails transforming a plant with a gene tag containing multiple transcriptional enhancers and once the tag has inserted into the genome, expression of a flanking gene coding sequence becomes deregulated. In another example, the transcriptional machinery in a plant can be modified so as to increase transcription levels of a polynucleotide of the invention (See, e.g., PCT Publications WO 96/06166 and WO 98/53057 which describe the modification of the DNA binding specificity of zinc finger proteins by changing particular amino acids in the DNA binding motif).

The transgenic plant can also include the machinery necessary for expressing or altering the activity of a polypeptide encoded by an endogenous gene, for example by altering the phosphorylation state of the polypeptide to maintain it in an activated state.

Transgenic plants (or plant cells, or plant explants, or plant tissues) incorporating the polynucleotides of the invention and/or expressing the polypeptides of the invention can be produced by a variety of well established techniques as described above. Following construction of a vector, most typically an expression cassette, including a polynucleotide, e.g., encoding a transcription factor or transcription factor homologue, of the invention, standard techniques can be used to introduce the polynucleotide into a plant, a plant cell, a plant explant or a plant tissue of interest. Optionally, the plant cell, explant or tissue can be regenerated to produce a transgenic plant.

The plant can be any higher plant, including gymnosperms, monocotyledonous and dicotyledenous plants. Suitable protocols are available for *Leguminosae* (alfalfa, soybean, clover, etc.), *Umbelliferae* (carrot, celery, parsnip), *Cruciferae* (cabbage, radish, rapeseed, broccoli, etc.), *Curcurbitaceae* (melons and cucumber), *Gramineae* (wheat, corn, rice, barley, millet, etc.), *Solanaceae* (potato, tomato, tobacco,

peppers, etc.), and various other crops. See protocols described in Ammirato et al. (1984) Handbook of Plant Cell Culture -Crop Species. Macmillan Publ. Co. Shimamoto et al. (1989) Nature 338:274-276; Fromm et al. (1990) Bio/Technology 8:833-839; and Vasil et al. (1990) Bio/Technology 8:429-434.

5

10

15

20

25

30

35

Transformation and regeneration of both monocotyledonous and dicotyledonous plant cells is now routine, and the selection of the most appropriate transformation technique will be determined by the practitioner. The choice of method will vary with the type of plant to be transformed; those skilled in the art will recognize the suitability of particular methods for given plant types. Suitable methods can include, but are not limited to: electroporation of plant protoplasts; liposome-mediated transformation; polyethylene glycol (PEG) mediated transformation; transformation using viruses; microinjection of plant cells; micro-projectile bombardment of plant cells; vacuum infiltration; and Agrobacterium tumeficiens mediated transformation. Transformation means introducing a nucleotide sequence in a plant in a manner to cause stable or transient expression of the sequence.

Successful examples of the modification of plant characteristics by transformation with cloned sequences which serve to illustrate the current knowledge in this field of technology, and which are herein incorporated by reference, include: U.S. Patent Nos. 5,571,706; 5,677,175; 5,510,471; 5,750,386; 5,597,945; 5,589,615; 5,750,871; 5,268,526; 5,780,708; 5,538,880; 5,773,269; 5,736,369 and 5,610,042.

Following transformation, plants are preferably selected using a dominant selectable marker incorporated into the transformation vector. Typically, such a marker will confer antibiotic or herbicide resistance on the transformed plants, and selection of transformants can be accomplished by exposing the plants to appropriate concentrations of the antibiotic or herbicide.

After transformed plants are selected and grown to maturity, those plants showing a modified trait are identified. The modified trait can be any of those traits described above. Additionally, to confirm that the modified trait is due to changes in expression levels or activity of the polypeptide or polynucleotide of the invention can be determined by analyzing mRNA expression using Northern blots, RT-PCR or microarrays, or protein expression using immunoblots or Western blots or gel shift assays.

INTEGRATED SYSTEMS—SEQUENCE IDENTITY

Additionally, the present invention may be an integrated system, computer or computer readable medium that comprises an instruction set for determining the identity of one or more sequences in a database. In addition, the instruction set can be used to generate or identify sequences that meet any specified criteria. Furthermore, the instruction set may

be used to associate or link certain functional benefits, such improved pathogen tolerance, with one or more identified sequence.

For example, the instruction set can include, e.g., a sequence comparison or other alignment program, e.g., an available program such as, for example, the Wisconsin Package Version 10.0, such as BLAST, FASTA, PILEUP, FINDPATTERNS or the like (GCG, Madision, WI). Public sequence databases such as GenBank, EMBL, Swiss-Prot and PIR or private sequence databases such as PhytoSeq (Incyte Pharmaceuticals, Palo Alto, CA) can be searched.

Alignment of sequences for comparison can be conducted by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity method of Pearson and Lipman (1988) Proc. Natl. Acad. Sci. U.S.A. 85: 2444, by computerized implementations of these algorithms. After alignment, sequence comparisons between two (or more) polynucleotides or polypeptides are typically performed by comparing sequences of the two sequences over a comparison window to identify and compare local regions of sequence similarity. The comparison window can be a segment of at least about 20 contiguous positions, usually about 50 to about 200, more usually about 100 to about 150 contiguous positions. A description of the method is provided in Ausubel et al., supra.

10

15

20

25

30

35

A variety of methods of determining sequence relationships can be used, including manual alignment and computer assisted sequence alignment and analysis. This later approach is a preferred approach in the present invention, due to the increased throughput afforded by computer assisted methods. As noted above, a variety of computer programs for performing sequence alignment are available, or can be produced by one of skill.

One example algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul et al. J. Mol. Biol 215:403-410 (1990). Software for performing BLAST analyses is publicly available, e.g., through the National Center for Biotechnology Information (http://www.ncbi.nlm.nih.gov/). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters

M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, a cutoff of 100, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff (1989) Proc. Natl. Acad. Sci. USA 89:10915).

10

15

20

25

30

35

In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence (and, therefore, in this context, homologous) if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, or less than about 0.01, and or even less than about 0.001. An additional example of a useful sequence alignment algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. The program can align, e.g., up to 300 sequences of a maximum length of 5,000 letters.

The integrated system, or computer typically includes a user input interface allowing a user to selectively view one or more sequence records corresponding to the one or more character strings, as well as an instruction set which aligns the one or more character strings with each other or with an additional character string to identify one or more region of sequence similarity. The system may include a link of one or more character strings with a particular phenotype or gene function. Typically, the system includes a user readable output element which displays an alignment produced by the alignment instruction set.

The methods of this invention can be implemented in a localized or distributed computing environment. In a distributed environment, the methods may implemented on a single computer comprising multiple processors or on a multiplicity of computers. The computers can be linked, e.g. through a common bus, but more preferably the computer(s) are nodes on a network. The network can be a generalized or a dedicated local or

wide-area network and, in certain preferred embodiments, the computers may be components of an intra-net or an internet.

Thus, the invention provides methods for identifying a sequence similar or homologous to one or more polynucleotides as noted herein, or one or more target polypeptides encoded by the polynucleotides, or otherwise noted herein and may include linking or associating a given plant phenotype or gene function with a sequence. In the methods, a sequence database is provided (locally or across an inter or intra net) and a query is made against the sequence database using the relevant sequences herein and associated plant phenotypes or gene functions.

Any sequence herein can be entered into the database, before or after querying the database. This provides for both expansion of the database and, if done before the querying step, for insertion of control sequences into the database. The control sequences can be detected by the query to ensure the general integrity of both the database and the query. As noted, the query can be performed using a web browser based interface. For example, the database can be a centralized public database such as those noted herein, and the querying can be done from a remote terminal or computer across an internet or intranet.

EXAMPLES

The following examples are intended to illustrate but not limit the present invention.

20 EXAMPLE I, FULL LENGTH GENE IDENTIFICATION AND CLONING

10

15

25

30

Putative transcription factor sequences (genomic or ESTs) related to known transcription factors were identified in the *Arabidopsis thaliana* GenBank database using the tblastn sequence analysis program using default parameters and a P-value cutoff threshold of -4 or -5 or lower, depending on the length of the query sequence. Putative transcription factor sequence hits were then screened to identify those containing particular sequence strings. If the sequence hits contained such sequence strings, the sequences were confirmed as transcription factors.

Alternatively, Arabidopsis thaliana cDNA libraries derived from different tissues or treatments, or genomic libraries were screened to identify novel members of a transcription family using a low stringency hybridization approach. Probes were synthesized using gene specific primers in a standard PCR reaction (annealing temperature 60° C) and labeled with ³²P dCTP using the High Prime DNA Labeling Kit (Boehringer Mannheim). Purified radiolabelled probes were added to filters immersed in Church hybridization medium (0.5 M NaPO₄ pH 7.0, 7% SDS, 1 % w/v bovine serum albumin) and hybridized overnight at 60 °C with shaking. Filters were washed two times for 45 to 60 minutes with 1xSCC, 1% SDS at 60° C.

To identify additional sequence 5' or 3' of a partial cDNA sequence in a cDNA library, 5' and 3' rapid amplification of cDNA ends (RACE) was performed using the MarathonTM cDNA amplification kit (Clontech, Palo Alto, CA). Generally, the method entailed first isolating poly(A) mRNA, performing first and second strand cDNA synthesis to generate double stranded cDNA, blunting cDNA ends, followed by ligation of the MarathonTM Adaptor to the cDNA to form a library of adaptor-ligated ds cDNA.

Gene-specific primers were designed to be used along with adaptor specific primers for both 5' and 3' RACE reactions. Nested primers, rather than single primers, were used to increase PCR specificity. Using 5' and 3' RACE reactions, 5' and 3' RACE fragments were obtained, sequenced and cloned. The process can be repeated until 5' and 3' ends of the full-length gene were identified. Then the full-length cDNA was generated by PCR using primers specific to 5' and 3' ends of the gene by end-to-end PCR.

EXAMPLE II. CONSTRUCTION OF EXPRESSION VECTORS

5

10

30

35

The sequence was amplified from a genomic or cDNA library using primers specific to sequences upstream and downstream of the coding region. The expression vector 15 was pMEN20 or pMEN65, which are both derived from pMON316 (Sanders et al, (1987) Nucleic Acids Research 15:1543-58) and contain the CaMV 35S promoter to express transgenes. To clone the sequence into the vector, both pMEN20 and the amplified DNA fragment were digested separately with SalI and NotI restriction enzymes at 37° C for 2 hours. 20 The digestion products were subject to electrophoresis in a 0.8% agarose gel and visualized by ethidium bromide staining. The DNA fragments containing the sequence and the linearized plasmid were excised and purified by using a Qiaquick gel extraction kit (Qiagen, CA). The fragments of interest were ligated at a ratio of 3:1 (vector to insert). Ligation reactions using T4 DNA ligase (New England Biolabs, MA) were carried out at 16° C for 16 25 hours. The ligated DNAs were transformed into competent cells of the E. coli strain DH5alpha by using the heat shock method. The transformations were plated on LB plates containing 50 mg/l kanamycin (Sigma).

Individual colonies were grown overnight in five milliliters of LB broth containing 50 mg/l kanamycin at 37° C. Plasmid DNA was purified by using Qiaquick Mini Prep kits (Qiagen, CA).

EXAMPLE III. TRANSFORMATION OF AGROBACTERIUM WITH THE EXPRESSION VECTOR

After the plasmid vector containing the gene was constructed, the vector was used to transform Agrobacterium tumefaciens cells expressing the gene products. The stock of Agrobacterium tumefaciens cells for transformation were made as described by Nagel et al. (1990) FEMS Microbiol Letts. 67: 325-328. Agrobacterium strain ABI was grown in 250 ml

LB medium (Sigma) overnight at 28°C with shaking until an absorbance (A_{600}) of 0.5 – 1.0 was reached. Cells were harvested by centrifugation at 4,000 x g for 15 min at 4° C. Cells were then resuspended in 250 μ l chilled buffer (1 mM HEPES, pH adjusted to 7.0 with KOH). Cells were centrifuged again as described above and resuspended in 125 μ l chilled buffer. Cells were then centrifuged and resuspended two more times in the same HEPES buffer as described above at a volume of 100 μ l and 750 μ l, respectively. Resuspended cells were then distributed into 40 μ l aliquots, quickly frozen in liquid nitrogen, and stored at -80° C.

Agrobacterium cells were transformed with plasmids prepared as described above following the protocol described by Nagel et al. For each DNA construct to be transformed, 50 – 100 ng DNA (generally resuspended in 10 mM Tris-HCl, 1 mM EDTA, pH 8.0) was mixed with 40 μl of Agrobacterium cells. The DNA/cell mixture was then transferred to a chilled cuvette with a 2mm electrode gap and subject to a 2.5 kV charge dissipated at 25 μF and 200 μF using a Gene Pulser II apparatus (Bio-Rad). After electroporation, cells were immediately resuspended in 1.0 ml LB and allowed to recover without antibiotic selection for 2 – 4 hours at 28° C in a shaking incubator. After recovery, cells were plated onto selective medium of LB broth containing 100 μg/ml spectinomycin (Sigma) and incubated for 24-48 hours at 28° C. Single colonies were then picked and inoculated in fresh medium. The presence of the plasmid construct was verified by PCR amplification and sequence analysis.

10

15

20

25

30

35

EXAMPLE IV. TRANSFORMATION OF ARABIDOPSIS PLANTS WITH AGROBACTERIUM TUMEFACIENS WITH EXPRESSION VECTOR

After transformation of Agrobacterium tumefaciens with plasmid vectors containing the gene, single Agrobacterium colonies were identified, propagated, and used to transform Arabidopsis plants. Briefly, 500 ml cultures of LB medium containing 50 mg/l kanamycin were inoculated with the colonies and grown at 28° C with shaking for 2 days until an absorbance (A_{600}) of > 2.0 is reached. Cells were then harvested by centrifugation at 4,000 x g for 10 min, and resuspended in infiltration medium (1/2 X Murashige and Skoog salts (Sigma), 1 X Gamborg's B-5 vitamins (Sigma), 5.0% (w/v) sucrose (Sigma), 0.044 μ M benzylamino purine (Sigma), 200 μ l/L Silwet L-77 (Lehle Seeds) until an absorbance (A_{600}) of 0.8 was reached.

Prior to transformation, Arabidopsis thaliana seeds (ecotype Columbia) were sown at a density of ~10 plants per 4" pot onto Pro-Mix BX potting medium (Hummert International) covered with fiberglass mesh (18 mm X 16 mm). Plants were grown under continuous illumination (50-75 μ E/m²/sec) at 22-23° C with 65-70% relative humidity. After about 4 weeks, primary inflorescence stems (bolts) are cut off to encourage growth of

multiple secondary bolts. After flowering of the mature secondary bolts, plants were prepared for transformation by removal of all siliques and opened flowers.

The pots were then immersed upside down in the mixture of Agrobacterium infiltration medium as described above for 30 sec, and placed on their sides to allow draining into a 1' x 2' flat surface covered with plastic wrap. After 24 h, the plastic wrap was removed and pots are turned upright. The immersion procedure was repeated one week later, for a total of two immersions per pot. Seeds were then collected from each transformation pot and analyzed following the protocol described below.

EXAMPLE V. IDENTIFICATION OF ARABIDOPSIS PRIMARY TRANSFORMANTS

Seeds collected from the transformation pots were sterilized essentially as follows. Seeds were dispersed into in a solution containing 0.1% (v/v) Triton X-100 (Sigma) and sterile H₂O and washed by shaking the suspension for 20 min. The wash solution was then drained and replaced with fresh wash solution to wash the seeds for 20 min with shaking. After removal of the second wash solution, a solution containing 0.1% (v/v) Triton X-100 and 70% ethanol (Equistar) was added to the seeds and the suspension was shaken for 5 min. After removal of the ethanol/detergent solution, a solution containing 0.1% (v/v) Triton X-100 and 30% (v/v) bleach (Clorox) was added to the seeds, and the suspension was shaken for 10 min. After removal of the bleach/detergent solution, seeds were then washed five times in sterile distilled H₂O. The seeds were stored in the last wash water at 4° C for 2 days in the dark before being plated onto antibiotic selection medium (1 X Murashige and Skoog salts (pH adjusted to 5.7 with 1M KOH), 1 X Gamborg's B-5 vitamins, 0.9% phytagar (Life Technologies), and 50 mg/l kanamycin). Seeds were germinated under continuous illumination (50-75 µE/m²/sec) at 22-23° C. After 7-10 days of growth under these conditions, kanamycin resistant primary transformants (T₁ generation) were visible and obtained. These seedlings were transferred first to fresh selection plates where the seedlings continued to grow for 3-5 more days, and then to soil (Pro-Mix BX potting medium).

Primary transformants were crossed and progeny seeds (T₂) collected; kanamycin resistant seedlings were selected and analyzed. The expression levels of the recombinant polynucleotides in the transformants varies from about a 5% expression level increase to a least a 100% expression level increase. Similar observations are made with respect to polypeptide level expression.

EXAMPLE VI. IDENTIFICATION OF ARABIDOPSIS PLANTS WITH TRANSCRIPTION FACTOR GENE KNOCKOUTS

30

5

10

15

20

25

The screening of insertion mutagenized Arabidopsis collections for null mutants in a known target gene was essentially as described in Krysan et al (1999) Plant Cell 11:2283-2290. Briefly, gene-specific primers, nested by 5-250 bases to each other, were designed from the 5' and 3' regions of a known target gene. Similarly, nested sets of primers were also created specific to each of the T-DNA or transposon ends (the "right" and "left" borders). All possible combinations of gene specific and T-DNA/transposon primers were used to detect by PCR an insertion event within or close to the target gene. The amplified DNA fragments were then sequenced which allows the precise determination of the T-DNA/transposon insertion point relative to the target gene. Insertion events within the coding or intervening sequence of the genes were deconvoluted from a pool comprising a plurality of insertion events to a single unique mutant plant for functional characterization. The method is described in more detail in Yu and Adam, US Application Serial No. 09/177,733 filed October 23, 1998.

15 EXAMPLE VII. IDENTIFICATION OF PATHOGEN INDUCED GENES

10

20

25

30

In some instances, expression patterns of the pathogen induced genes (such as defense genes) was monitored by microarray experiments. cDNAs were generated by PCR and resuspended at a final concentration of ~ 100 ng/ul in 3X SSC or 150mM Na-phosphate (Eisen and Brown (1999) *Meth. in Enzymol.* 303:179-205). The cDNAs were spotted on microscope glass slides coated with polylysine. The prepared cDNAs were aliquoted into 384 well plates and spotted on the slides using an x-y-z gantry (OmniGrid) purchased from GeneMachines (Menlo Park, CA) outfitted with quill type pins purchased from Telechem International (Sunnyvale, CA). After spotting, the arrays were cured for a minimum of one week at room temperature, rehydrated and blocked following the protocol recommended by Eisen and Brown (1999).

Sample total RNA (10 ug) samples were labeled using fluorescent Cy3 and Cy5 dyes. Labeled samples were resuspended in 4X SSC/0.03% SDS/4 ug salmon sperm DNA/2 ug tRNA/50mM Na-pyrophosphate, heated for 95°C for 2.5 minutes, spun down and placed on the array. The array was then covered with a glass coverslip and placed in a sealed chamber. The chamber was then kept in a water bath at 62°C overnight. The arrays were washed as described in Eisen and Brown (1999) and scanned on a General Scanning 3000 laser scanner. The resulting files are subsequently quantified using Imagene a software purchased from BioDiscovery (Los Angeles, CA).

EXAMPLE VIII. IDENTIFICATION OF PATHOGEN TOLERANCE PHENOTYPE IN OVEREXPRESSOR OR GENE KNOCKOUT PLANTS

5

10

15

20

25

30

35

Experiments were performed to identify those transformants or knockouts that exhibited an improved pathogen tolerance. For such studies, the transformants were exposed to biotropic fungal pathogens, such as Erisyphe orontii, and necrotropic fungal pathogens, such as Fusarium oxysporum. Fusarium oxysporum isolates cause vascular wilts and damping off of various annual vegetables, perennials and weeds (Mauch-Mani and Slusarenko (1994) Molecular Plant-Microbe Interactions 7: 378-383). For Fusarium oxysporum experiments, plants grown on petri dishes were sprayed with a fresh spore suspension of F. oxysporum. The spore suspension was prepared as follows: A plug of fungal hyphae from a plate culture was placed on a fresh potato dextrose agar plate and allowed to spread for one week. 5 ml sterile water was then added to the plate, swirled, and pipetted into 50 ml Armstrong Fusarium medium. Spores were grown overnight in Fusarium medium and then sprayed onto plants using a Preval paint sprayer. Plant tissue was harvested and frozen in liquid nitrogen 48 hours post infection.

Erysiphe orontii is a causal agent of powdery mildew. For Erysiphe orontii experiments, plants were grown approximately 4 weeks in a greenhouse under 12 hour light (20 C, ~30% relative humidity (rh)). Individual leaves were infected with E. orontii spores from infected plants using a camel's hair brush, and the plants were transferred to a Percival growth chamber (20 C, 80% rh.). Plant tissue was harvested and frozen in liquid nitrogen 7 days post infection.

Botrytis cinerea is a necrotrophic pathogen. Botrytis cinerea was grown on potato dextrose agar in the light. A spore culture was made by spreading 10 ml of sterile water on the fungus plate, swirling and transferring spores to 10 ml of sterile water. The spore inoculum (approx. 105 spores/ml) was used to spray 10 day-old seedlings grown under sterile conditions on MS (-sucrose) media. Symptoms were evaluated every day up to approximately 1 week.

Infection with bacterial pathogens Pseudomonas syringae pv maculicola strain 4326 and pv maculicola strain 4326 was performed by hand inoculation at two doses. Two inoculation doses allows the differentiation between plants with enhanced susceptibility and plants with enhanced resistance to the pathogen. Plants were grown for 3 weeks in the greenhouse, then transferred to the growth chamber for the remainder of their growth. Psm ES4326 was hand inoculated with 1 ml syringe on 3 fully-expanded leaves per plant (4 1/2 wk old), using at least 9 plants per overexpressing line at two inoculation doses, OD=0.005 and OD=0.0005. Disease scoring occurred at day 3 post-inoculation with pictures of the plants and leaves taken in parallel.

Table 3 shows the phenotypes observed for particular overexpressor or knockout plants and provides the SEQ ID No., the internal reference code (GID), whether a knockout or overexpressor plant was analyzed and the observed phenotype.

Table 3

SEQ ID No.	GID	Knockout (KO) or overexpressor (OE)	Phenotype
1	G188	КО	Increased susceptibility to Fusarium
3	G616	OE	Increased tolerance to Erysiphe
5	G19	OE	Increased tolerance to Erysiphe
7.	G261	OE	Increased susceptibility to Botrytis
9	G28	OE	Increased resistance to Erysiphe
11	G869	OE	Increased susceptibility to Fusarium
13	G237	OE	Increased tolerance to Erysiphe
15	G409	OE	Increased tolerance to Erysiphe
17	G418	OE	Increased tolerance to Pseudomonas
19	G591	OE	Increased tolerance to Erysiphe
21	G525	OE	Increased tolerance to Pseudomonas
23	G545	OE	Increased susceptibility to Pseudomonas, Erysiphe and Fusarium
25	G865	OE	Increased susceptibility to Erysiphe and Botrytis
27	G881	OE	Increased susceptibility to Erysiphe and Botrytis
29	G896	KO	Increased susceptibility to Fusarium
31	G378	OE	Increased resistance to Erysiphe
33	G569	OE	Decreased expression of defense genes
35	G558	OE	Increased expression of defense genes

For a particular overexpressor that shows an increased susceptibility to a pathogen, it may be more useful to select a plant with a decreased expression of the particular transcription factor. For a particular knockout that shows an increased susceptibility to a pathogen, it may be more useful to select a plant with an increased expression of the particular transcription factor.

Other than Fusarium oxysporum, Erysyphe orontii, the transgenic plants are more tolerant to Sclerotinia spp., soil-borne oomycetes, foliar oomycetes, Botrytis spp., Rhizoctonia spp, Verticillium dahliae/albo-atrum, Alternaria spp., rusts, Mycosphaerella spp, Fusarium solani, or the like. The transgenic plants are more resistant to fungal diseases such as rusts, smuts, wilts, yellows, root rot, leaf drop, ergot, leaf blight of potato, brown spot of rice, leaf blight, late blight, powdery mildew, downy mildew, and the like; viral diseases such as sugarcane mosaic, cassava mosaic, sugar beet yellows, plum pox, barley yellow dwarf, tomato yellow leaf curl, tomato spotted wilt virus, and the like; bacterial diseases such as citrus canker, bacterial leaf blight, bacterial will, soft rot of vegetables, and the like; nematode diseases such as root knot, sugar beet cyst nematode or the like.

5

10

EXAMPLE IX. IDENTIFICATION OF HOMOLOGOUS SEQUENCES

5

10

15

20

25

30

35

Homologous sequences from *Arabidopsis* and plant species other than *Arabidopsis* were identified using database sequence search tools, such as the Basic Local Alignment Search Tool (BLAST) (Altschul et al. (1990) <u>J. Mol. Biol.</u> 215:403-410; and Altschul et al. (1997) <u>Nucl. Acid Res.</u> 25: 3389-3402). The tblastx sequence analysis programs were employed using the BLOSUM-62 scoring matrix (Henikoff, S. and Henikoff, J. G. (1992) <u>Proc. Natl. Acad. Sci. USA</u> 89: 10915-10919).

Identified Arabidopsis homologous sequences are provided in Figure 2 and included in the Sequence Listing. The percent sequence identity among these sequences is as low as 47% sequence identity. Additionally, the entire NCBI GenBank database was filtered for sequences from all plants except Arabidopsis thaliana by selecting all entries in the NCBI GenBank database associated with NCBI taxonomic ID 33090 (Viridiplantae; all plants) and excluding entries associated with taxonomic ID 3701 (Arabidopsis thaliana). These sequences were compared to sequences representing genes of SEQ IDs Nos. 1-58 on 9/26/2000 using the Washington University TBLASTX algorithm (version 2.0a19MP). For each gene of SEQ IDs Nos. 1-58, individual comparisons were ordered by probability score (P-value), where the score reflects the probability that a particular alignment occurred by chance. For example, a score of 3.6e-40 is 3.6 x 10⁻⁴⁰. For up to ten species, the gene with the lowest P-value (and therefore the most likely homolog) is listed in Figure 3.

In addition to P-values, comparisons were also scored by percentage identity. Percentage identity reflects the degree to which two segments of DNA or protein are identical over a particular length. The ranges of percent identity between the non-Arabidopsis genes shown in Figure 3 and the Arabidopsis genes in the sequence listing are: SEQ ID No. 1: 38%-76%; SEQ ID No. 3: 36%-72%; SEQ ID No. 5: 51%-75%; SEQ ID No. 7: 37%-76%; SEQ ID No. 9: 48%-75%; SEQ ID No. 11: 31%-68%; SEQ ID No. 13: 59%-81%; SEQ ID No. 15: 49%-81%; SEQ ID No. 17: 53%-87%; SEQ ID No. 19: 48%-84%; SEQ ID No. 21: 73%-89%; SEQ ID No. 23: 52%-64%; SEQ ID No. 25: 48%-83%; SEQ ID No. 27: 35%-92%; SEQ ID No. 29: 56%-89%; SEQ ID No. 31: 50%-90%; SEQ ID No. 33: 50%-93%; SEQ ID No. 35: 52%-81%; SEQ ID No. 37: 75%-81%; SEQ ID No. 39: 35%-72%; SEQ ID No. 41: 55%-89%; SEQ ID No. 43: 56%-77%; SEQ ID No. 45: 34%-72%; SEQ ID No. 47: 51%-86%; SEQ ID No. 49: 46%-86%; SEQ ID No. 51: 58%-80%; SEQ ID No. 53: 46%-55%; SEQ ID No. 55: 84%-89%; and SEQ ID No. 57: 43%-71%.

The polynucleotides and polypeptides in the Sequence Listing and the identified homologous sequences may be stored in a computer system and have associated or linked with the sequences a function, such as that the polynucleotides and polypeptides are useful for modifying the pathogen tolerance of a plant.

All references, publications, patents and other documents herein are incorporated by reference in their entirety for all purposes. Although the invention has been described with reference to the embodiments and examples above, it should be understood that various modifications can be made without departing from the spirit of the invention.

What is claimed is:

5

15

20

1. A transgenic plant with modified pathogen tolerance, which plant comprises a recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29, or a complementary nucleotide sequence thereof:
 - (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);
- (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos.2N-1, where N=1-29, or a complementary nucleotide sequence thereof;
 - (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of(c);
 - (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
 - (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e);
 - (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide that modifies a plant's pathogen tolerance;
 - (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
 - (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g);
- 25 (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29;
 - (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; and
- (1) a nucleotide sequence which encodes a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-29.
 - 2. The transgenic plant of claim 1, further comprising a constitutive, inducible, or tissue-active promoter operably linked to said nucleotide sequence.
- 35 3. The transgenic plant of claim 1, wherein the plant is selected from the group consisting of: soybean, wheat, corn, potato, cotton, rice, oilseed rape, sunflower, alfalfa, sugarcane, turf, banana, blackberry, blueberry, strawberry, raspberry, cantaloupe, carrot,

cauliflower, coffee, cucumber, eggplant, grapes, honeydew, lettuce, mango, melon, onion, papaya, peas, peppers, pineapple, spinach, squash, sweet corn, tobacco, tomato, watermelon, rosaceous fruits, and vegetable brassicas.

- 5 4. An isolated or recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) a nucleotide sequence encoding a polypeptide comprising a sequence selected from SEQ ID Nos. 2N, where N=1-29, or a complementary nucleotide sequence thereof;
- 10 (b) a nucleotide sequence encoding a polypeptide comprising a conservatively substituted variant of a polypeptide of (a);

15

- (c) a nucleotide sequence comprising a sequence selected from those of SEQ ID Nos. 2N-1, where N=1-29, or a complementary nucleotide sequence thereof;
- (d) a nucleotide sequence comprising silent substitutions in a nucleotide sequence of (c);
- (e) a nucleotide sequence which hybridizes under stringent conditions to a nucleotide sequence of one or more of: (a), (b), (c), or (d);
- (f) a nucleotide sequence comprising at least 15 consecutive nucleotides of a sequence of any of (a)-(e);
- 20 (g) a nucleotide sequence comprising a subsequence or fragment of any of (a)-(f), which subsequence or fragment encodes a polypeptide having a biological activity that modifies a plant's pathogen tolerance;
 - (h) a nucleotide sequence having at least 31% sequence identity to a nucleotide sequence of any of (a)-(g);
- 25 (i) a nucleotide sequence having at least 60% identity sequence identity to a nucleotide sequence of any of (a)-(g);
 - (j) a nucleotide sequence which encodes a polypeptide having at least 31% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29;
 - (k) a nucleotide sequence which encodes a polypeptide having at least 60% identity sequence identity to a polypeptide of SEQ ID Nos. 2N, where N=1-29; and
 - (1) a nucleotide sequence which encodes a conserved domain of a polypeptide having at least 65% sequence identity to a conserved domain of a polypeptide of SEQ ID Nos. 2N, where N=1-29.
- 35 5. The isolated or recombinant polynucleotide of claim 4, further comprising a constitutive, inducible, or tissue-active promoter operably linked to the nucleotide sequence.

6. A cloning or expression vector comprising the isolated or recombinant polynucleotide of claim 4.

- 7. A cell comprising the cloning or expression vector of claim 6.
- 8. A transgenic plant comprising the isolated or recombinant polynucleotide of claim 4.
- 9. A composition produced by one or more of:
 - (a) incubating one or more polynucleotide of claim 4 with a nuclease;
- 10 (b) incubating one or more polynucleotide of claim 4 with a restriction enzyme;
 - (c) incubating one or more polynucleotide of claim 4 with a polymerase;
 - (d) incubating one or more polynucleotide of claim 4 with a polymerase and a primer;
 - (e) incubating one or more polynucleotide of claim 4 with a cloning vector, or
 - (f) incubating one or more polynucleotide of claim 4 with a cell.

15

5

- 10. A composition comprising two or more different polynucleotides of claim 4.
- 11. An isolated or recombinant polypeptide comprising a subsequence of at least about 15 contiguous amino acids encoded by the recombinant or isolated polynucleotide of claim 4.

20

25

- 12. A plant comprising an isolated polypeptide of claim 11.
- 13. A method for producing a plant having a modified pathogen tolerance, the method comprising altering the expression of the isolated or recombinant polynucleotide of claim 4 or the expression levels or activity of a polypeptide of claim 11 in a plant, thereby producing a modified plant, and selecting the modified plant for improved pathogen tolerance thereby providing the modified plant with a modified pathogen tolerance.
 - 14. The method of claim 13, wherein the polynucleotide is a polynucleotide of claim 4.

- 15. A method of identifying a factor that is modulated by or interacts with a polypeptide encoded by a polynucleotide of claim 4, the method comprising:
 - (a) expressing a polypeptide encoded by the polynucleotide in a plant; and
- (b) identifying at least one factor that is modulated by or interacts with the polypeptide.

16. The method of claim 15, wherein the identifying is performed by detecting binding by the polypeptide to a promoter sequence, or detecting interactions between an additional protein and the polypeptide in a yeast two hybrid system.

- 5 17. The method of claim 15, wherein the identifying is performed by detecting expression of a factor by hybridization to a microarray, subtractive hybridization or differential display.
 - 18. A method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide of interest, the method comprising:
- (a) placing the molecule in contact with a plant comprising the polynucleotide or polypeptide encoded by the polynucleotide of claim 4; and,
 - (b) monitoring one or more of:
 - (i) expression level of the polynucleotide in the plant;
 - (ii) expression level of the polypeptide in the plant;
 - (iii) modulation of an activity of the polypeptide in the plant; or
 - (iv) modulation of an activity of the polynucleotide in the plant.
 - 19. An integrated system, computer or computer readable medium comprising one or more character strings corresponding to a polynucleotide of claim 4, or to a polypeptide encoded by the polynucleotide.
 - 20. The integrated system, computer or computer readable medium of claim 19, further comprising a link between said one or more sequence strings to a modified plant pathogen tolerance phenotype.

25

15

- 21. A method of identifying a sequence similar or homologous to one or more polynucleotides of claim 4, or one or more polypeptides encoded by the polynucleotides, the method comprising:
 - (a) providing a sequence database; and,
- (b) querying the sequence database with one or more target sequences corresponding to the one or more polynucleotides or to the one or more polypeptides to identify one or more sequence members of the database that display sequence similarity or homology to one or more of the one or more target sequences.
- 35 22. The method of claim 21, wherein the querying comprises aligning one or more of the target sequences with one or more of the one or more sequence members in the sequence database.

23. The method of claim 21, wherein the querying comprises identifying one or more of the one or more sequence members of the database that meet a user-selected identity criteria with one or more of the target sequences.

5

- 24. The method of claim 21, further comprising linking the one or more of the polynucleotides of claim 4, or encoded polypeptides, to a modified plant pathogen tolerance phenotype.
- 10 25. A plant comprising altered expression levels of an isolated or recombinant polynucleotide of claim 4.
 - 26. A plant comprising altered expression levels or the activity of an isolated or recombinant polypeptide of claim 11.

15

27. A plant lacking a nucleotide sequence encoding a polypeptide of claim 11.

Figure 1

SEQ ID No.	GID	cDNA or protein	conserved domain
1	G188	cDNA	
2	G188	protein	175-222
3	G616	cDNA	
4	G616	protein	39-95
5	G19	cDNA	
6	G19	protein	76-145
7	G261	cDNA	
8	G261	protein-	16-104
9	G28	cDNA	
10	G28	protein	145-213
11	G869	cDNA	
12	G869	protein	109-177
13	G237	cDNA	
14	G237	protein	11-113
15	G409	cDNA	
16	G409	protein	64-124
17	G418	cDNA	
18	G418	protein	500-560
19	G591	cDNA	
20	G591	protein	143-240
21	G525	cDNA	
22	G525	protein	23-167
23	G545	cDNA	
24	G545	protein	82-102, 136-154
25	G865	cDNA	
26	G865	protein	36-103
27	G881	cDNA	
28	G881	protein	176-233
29	G896	cDNA	
30	G896	protein	18-39
31	G378	cDNA	
32_	G378	protein	196-237
33	G569	cDNA	
34	G569	protein	90-153
35	G558	cDNA	
36	G558	protein	45-105

Figure 2

SEQ ID No.	GID	homolog	cDNA or protein	conserved domain
37	G1396	homolog of G1394	cDNA	
38	G1396	homolog of G1394	protein	entire protein
39	G265	homolog of G261	cDNA	
40	G265	homolog of G261	protein	14-105
41	G1006	homolog of G28	cDNA	
42	G1006	homolog of G28	protein	114-182
43	G1309	homolog of G237	cDNA	
44	G1309	homolog of G237	protein	9-114
45	G2550	homolog of G418	cDNA	
46	G2550	homolog of G418	protein	348-408
47	G965	homolog of G418	cDNA	
48	G965	homolog of G418	protein	423-486
49	G793	homolog of G591	cDNA	
50	G793	homolog of G591	protein	151-206
51	G764	homolog of G525	cDNA	
52	G764	homolog of G525	protein	27-171
53	G350	homolog of G545	cDNA	
54	G350	homolog of G545	protein	91-113,150-170
55	G986	homolog of G881	cDNA	
56	G986	homolog of G881	protein	146-203
57	G1349	homolog of G896	cDNA	
58	G1349	homolog of G896	protein	13-63

Figure 3A

SEQ ID No.	GID	Genbank NID	P-value	Species
1	G188	7779802	5.20E-36	Lotus japonicus
1	G188	7284340	2.10E-34	Glycine max
1	G188	9361307	1.20E-27	Triticum aestivum
1	G188	7340336	1.10E-22	Oryza sativa
1	G188	6529152	3.60E-22	Lycopersicon esculentum
1	G188	8748477	7.70E-21	Medicago truncatula
1	G188	5456433	7.10É-14	Zea mays
1	G188	9302479	1.60E-12	Sorghum bicolor
1	G188	6696287	4.10E-12	Pinus taeda
1	G188	562242	9.00E-12	Brassica rapa
3	G616	7719440	8.30E-37	Lotus japonicus
3	G616	7692230	5.90E-33	Glycine max
3	G616	7501307	1.10E-21	Gossypium arboreum
3	G616	8071090	1.50E-21	Solanum tuberosum
3	G616	8858771	1.50E-21	Oryza sativa
3	G616	5047315	1.50E-21	Gossypium hirsutum
3	G616	6358532	5.80E-20	Antirrhinum graniticum
3	G616	2826867	7.00E-20	Antirrhinum majus
3	G616	6358535	7.40E-20	
3	G616	6358538	7.40E-20 7.50E-20	Antirrhinum majus subsp. linkianum
5	G19			Antimhinum braun-blanquetii
5		8789223	2.80E-34	Citrus x paradisi
5	G19 G19	9434234	4.50E-34	Lycopersicon esculentum
5		7478682	1.30E-30	Glycine max
5	G19	6654934	1.20E-28	Medicago truncatula
5	G19	3264766	5.50E-26	Prunus armeniaca
5	G19	7624302	8.30E-26	Gossypium arboreum
5	G19 G19	9425363	2.90E-25	Triticum aestivum
5		688579	3.60E-25	Ricinus communis
5	G19	9419304	6.00E-25	Hordeum vulgare
7	G19	7720316	8.80E-25	Lotus japonicus
	G261	5821137	5.10E-93	Nicotiana tabacum
7	G261	7158881	8.80E-86	Medicago sativa
7	G261	886741	1.00E-73	Zea mays
7	G261	5900449	5.20E-47	Lycopersicon esculentum
7	G261	7561318	1.20E-46	Medicago truncatula
7	G261	19491	1.70E-42	Lycopersicon peruvianum
	G261	7233914	3.50E-41	Glycine max
7	G261	4528238	9.00E-41	Citrus unshiu
7 7	G261	8903922	4.00E-39	Hordeum vulgare
	G261	9251913	1.90E-36	Solanum tuberosum
9	G28	7528275	4.20E-62	Mesembryanthernum crystallinum
9	G28	6654776	1.20E-57	Medicago truncatula
9	G28	790362	2.30E-54	Nicotiana tabacum
9	G28	8809570	8.00E-54	Nicotiana sylvestris
9	G28	3342210	8.40E-54	Lycopersicon esculentum
9	G28	6566281	8.40E-47	Glycine max
9	G28	7627061	8.40E-47	Gossypium arboreum
9	G28	7324479	2.00E-44	Lycopersicon pennellii
9	G28	6478844	1.80E-35	Matricaria chamomilla
9	G28	7273972	7.80E-29	Oryza sativa
11	G869	2213784	1.30E-19	Lycopersicon esculentum
11	G869	3065894	7.30E-19	Nicotiana tabacum

Figure 3B

SEQ ID No.	GID	Genbank NID	P-value	Species
				Oryza sativa
11	G869	8570080	4.20E-18	Medicago truncatula
11	G869	7560260	1.50E-17	
11	G869	7534890	5.20E-14	Sorghum bicolor
11	G869	6455322	1.10E-13	Glycine max Triticum aestivum
11	G869	9362061	2.70E-13	
11	G869	7788764	5.70E-13	Lotus japonicus
11	G869	7624302	2.50E-12	Gossypium arboreum
11	G869	3858036	2.80E-12	Populus balsamifera subsp. trichocarpa
13	G237	8283916	4.70E-42	Glycine max
13	G237	9361969	8.30E-41	Triticum aestivum
13	G237	4753385	4.10E-39	Zea mays
13	G237	7535969	4.10E-33	Sorghum bicolor
13	G237	7566043	9.30E-33	Medicago truncatula
13	G237	7339127	2.00E-32	Lycopersicon esculentum
13	G237	5860031	1.10E-28	Pinus taeda
13	G237	7776223	2.20E-28	Lotus japonicus
13	G237	6850206	5.10E-28	Oryza sativa
13	G237	5048991	8.50E-28	Gossypium hirsutum
15	G409	6654773	6.10E-57	Medicago truncatula
15	G409	6531235	2.00E-56	Lycopersicon esculentum
15	G409	7924152	1.10E-47	Glycine max
15	G409	. 5006854	6.50E-43.	Oryza sativa
15	G409	8098529	2.10E-41	Hordeum vulgare
15	G409	767697	1.40E-37	Daucus carota
15	G409	8328991	3.30E-37	Mesembryanthemum crystallinum
15	G409	7415613	1.40E-32	Physcomitrella patens
15	G409	7785121	2.80E-32	Lotus japonicus
15	G409	6916941	4.80E-32	Lycopersicon pennellii
17	G418	7239156 ⁻	1.90E-123	Malus x domestica
17	G418	5892190	2.00E-62	Lycopersicon esculentum
17	G418	7628137	8.70E-58	Gossypium arboreum
17	G418	9205496	3.90E-51	Glycine max
17	G418	6069643	1.50E-45	Oryza sativa
17	G418	7562931	6.90E-45	Medicago truncatula
17	G418	7781695	5.50E-40	Lotus japonicus
17	G418	9298824	7.80E-34	Sorghum bicolor
17	G418	9428023	3.90E-32	Triticum aestivum
17	G418	7244366	1.30E-31	Mentha x piperita
19	G591	7646333	1.90E-55	Lycopersicon esculentum
19	G591	7924288	4.10E-53	Glycine max
19	G591	7722838	1.10E-41	Lotus japonicus
19	G591	5804781	1.40E-24	Nicotiana tabacum
19	G591	9198126	2.50E-23	Medicago truncatula
19	G591	427677	9.50E-15	Oryza sativa
19	G591	7624745	1.80E-14	Gossypium arboreum
19	G591	7535578	8.70E-14	Sorghum bicolor
19	G591	5915205	1.30E-11	Zea mays
19	G591	9249806	2.60E-11	Solanum tuberosum
21	G525	4384535	5.60E-61	Lycopersicon esculentum
21	G525	6454868	2.00E-58	Glycine max
21	G525	6066594	9.30E-54	Petunia x hybrida
21	G525	4977542	8.60E-51	Oryza sativa

Figure 3C

21 G525 9361647 2.50E-50 Triticum aestivum 21 G525 4218536 5.20E-50 Triticum sp. 21 G525 6732159 5.20E-50 Triticum monococcum 21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 7228328 3.70E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 438462 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula	
21 G525 4218536 5.20E-50 Triticum sp. 21 G525 6732159 5.20E-50 Triticum monococcum 21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum <td></td>	
21 G525 6732159 5.20E-50 Triticum monococcum 21 G525 5343151 2.70E-49 Zea mays 21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
21 G525 5049217 4.20E-48 Gossypium hirsutum 21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
21 G525 8708684 8.90E-48 Hordeum vulgare 23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 4666359 8.30E-55 Datisca glomerata 23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 7228328 3.70E-52 Medicago sativa 23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 1763062 1.30E-51 Glycine max 23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 7206360 3.10E-44 Medicago truncatula 23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 7626808 9.60E-40 Gossypium arboreum 23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 439492 3.90E-39 Petunia x hybrida 23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 4382658 1.70E-38 Lycopersicon esculentum 23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 8486215 8.70E-38 Euphorbia esula 23 G545 7322653 6.80E-37 Lycopersicon hirsutum	· ·
23 G545 7322653 6.80E-37 Lycopersicon hirsutum	
23 G545 7785845 1.10E-33 Lotus japonicus	<u> </u>
25 G865 9417297 1.70E-32 Triticum aestivum	
25 G865 7206394 4.90E-29 Medicago truncatula	
25 G865 7796858 5.70E-27 Glycine max	
25 G865 4387560 9.20E-25 Lycopersicon esculentum	1
25 G865 569065 1.50E-23 Oryza sativa	<u> </u>
25 G865 7788764 4.10E-23 Lotus japonicus	
25 G865 790362 8.40E-22 Nicotiana tabacum	
25 G865 7528275 5.90E-21 Mesembryanthemum crys	stallinum
25 G865 3264766 8.80E-20 Prunus armeniaca	
25 G865 8098026 2.00E-19 Hordeum vulgare	
27 G881 5820418 9.80E-29 Glycine max	
27 G881 8440065 1.00E-27 Gossypium hirsutum	
27 G881 4380578 1.50E-27 Lycopersicon esculentum	
27 G881 9199620 2.70E-27 Medicago truncatula	
27 G881 6472584 2.20E-24 Nicotiana tabacum	· · · · · · · · · · · · · · · · · · ·
27 G881 9250698 3.20E-24 Solanum tuberosum	
27 G881 8205146 5.20E-21 Oryza sativa	
27 G881 1159878 8.20E-17 Avena fatua	
27 G881 9299778 2.70E-16 Sorghum bicolor	
27 G881 9444636 1.10E-14 Triticum aestivum	······
29 G896 9410462 1.90E-101 Hordeum vulgare	
29 G896 7628908 3.60E-82 Gossypium arboreum	
29 G896 7244408 1.80E-79 Mentha x piperita	
29 G896 5046180 2.10E-73 Gossypium hirsutum	
29 G896 7678652 1.10E-63 Lotus japonicus	
29 G896 8286031 1.40E-60 Glycine max	
29 G896 5888938 4.50E-58 Lycopersicon esculentum	
29 G896 9298238 9.20E-54 Sorghum bicolor	
29 G896 7566414 8.00E-52 Medicago truncatula	
29 G896 8845076 1.00E-46 Triticum aestivum	
31 G378 5270028 5.10E-73 Lycopersicon esculentum	
31 G378 5048335 4.10E-58 Gossypium hirsutum	
31 G378 7239521 5.90E-42 Oryza sativa	
31 G378 5606120 6.80E-36 Glycine max	
31 G378 3853800 3.20E-30 Populus tremula x Populu	s tremuloides
31 G378 7659983 1.70E-23 Sorghum bicolor	

Figure 3D

SEQ ID No.	GID	Genbank NID	P-value	Species
31	G378	6626305	1.10E-21	Zea mays
31	G378	9412941	9.40E-19	Triticum aestivum
31	G378	3242033	4.30E-17	Mesembryanthemum crystallinum
31	G378	7626259	7.70E-13	Gossypium arboreum
33	G229	7337390	6.60E-51	Lycopersicon esculentum
33	G229	9823237	3.60E-50	Hordeum vulgare
33	G229	7244424	4.90E-50	Mentha x piperita
33	G229	7776053	1.70E-49	Lotus japonicus
33	G229	2921335	5.80E-48	Gossypium hirsutum
33	G229	1491932	4.50E-47	Zea mays
33	G229	6455590	2.80E-44	Glycine max
33	G229	6020191	2.00E-41	Pinus taeda
33	G229	10697236	4.20E-41	Oryza sativa
33	G229	7765706	5.10E-41	Medicago truncatula
35	G663	7673087	5.10E-43	Petunia integrifolia
35	G663	9508051	3.00E-41	Lycopersicon esculentum
35	G663	7673091	3.30E-41	Petunia x hybrida
35	G663	7673097	2.40E-36	Petunia axillaris
35	G663	5048991	1.20E-33	Gossypium hirsutum
	G663	6455590	2.50E-31	Glycine max
35 35	G663	7560175	1.90E-27	Medicago truncatula
	G663	7244424	4.10E-26	Mentha x piperita
35	G663	9954117	3.40E-25	Solanum tuberosum
35		6020191	3.60E-25	Pinus taeda
35	G663		5.20E-22	Spinacia oleracea
37	G1396	498704		Gossypium arboreum
37	G1396	7502400	1.20E-21	Populus balsamifera subsp. trichocarpa
37	G1396	3857536	3.40E-21 1.20E-20	Lycopersicon esculentum
37	G1396	4385300	1.50E-20	Lycopersicon pennellii
37	G1396	6917249	1.70E-20	Glycine max
37	G1396	6915979		Medicago truncatula
37	G1396	7674530	2.70E-20	Sorghum bicolor
37	G1396	8090319	3.40E-20	
37	G1396	3592182	9.10E-20	Oryza sativa
37	G1396	6654124	1.10E-19	Zea mays
39	G265	5821137	6.50E-83	Nicotiana tabacum
39	G265	7158881	3.80E-79	Medicago sativa
39	G265	886741	1.60E-70	Zea mays
39	G265	5900449	5.60E-43	Lycopersicon esculentum
39	G265	8903922	8.20E-43	Hordeum vulgare
39	G265	7561318	2.10E-41	Medicago truncatula
39	G265	9204445	5.30E-36	Glycine max
. 39	G265	4528238	5.40E-36	Citrus unshiu
39	G265	19489	2.10E-35	Lycopersicon peruvianum
39	G265	9251913	2.00E-32	Solanum tuberosum
41	G1006	7528275	2.70E-51	Mesembryanthemum crystallinum
41	G1006	3342210	4.90E-49	Lycopersicon esculentum
41	G1006	6654776	1.90E-48	Medicago truncatula
41	G1006	790362	2.30E-47	Nicotiana tabacum
41	G1006	8809570	2.00E-46	Nicotiana sylvestris
41	G1006	7627061	6.40E-41	Gossypium arboreum
41	G1006	7324479	1.20E-35	Lycopersicon pennellii
41	G1006	6478844	1.80E-35	Matricaria chamomilla

Figure 3E

SEQ ID No.	GID	Genbank NID	P-value	Species
41	G1006	6566281	1.30E-34	Glycine max
41	G1006	4716624	3.80E-28	Oryza sativa
43	G1309	9361969	2.40E-45	Triticum aestivum
43	G1309	7566043	9.60E-35	Medicago truncatula
43	G1309	5891104	2.20E-31	Lycopersicon esculentum
43	G1309	5860031	2.10E-30	Pinus taeda
43	G1309	5049507	6.20E-30	Gossypium hirsutum
43	G1309	5139805	1.30E-29	Glycine max
43	G1309	6850206	2.50E-29	Oryza sativa
43	G1309	7721017	3.40E-29	Lotus japonicus
43	G1309	8368245	5.20E-28	Zea mays
43	G1309	20560	9.50E-27	Petunia x hybrida
45	G2550	4380729	2.80E-51	Lycopersicon esculentum
45	G2550	5667196	2.20E-49	Oryza sativa
45	G2550	8669454	1.40E-48	Glycine max
45	G2550	9298824	1.50E-48	Sorghum bicolor
45	G2550	7239156	9.90E-46	Malus x domestica
45	G2550	7570704	5.70E-45	Medicago truncatula
45	G2550	7628137	3.30E-42	Gossypium arboreum
. 45	G2550	7244366	6.00E-41	Mentha x piperita
45	G2550	9428023	4.70E-40	Triticum aestivum
45	G2550	9250642	3.50E-39	Solanum tuberosum
47	G965	7239156	3.10E-126	Malus x domestica
47	G965	5892190	2.00E-62	Lycopersicon esculentum
47	G965	7628137	1.60E-56	Gossypium arboreum
47	G965	9205496	2.60E-49	Glycine max
47	G965	6069643	1.70E-45	Oryza sativa
47	G965	7562931	2.50E-44	Medicago truncatula
47	G965	7781695	1.60E-41	Lotus japonicus
47	G965	9298824	6.30E-33	Sorghum bicolor
47	G965	9428023	1.50E-31	Triticum aestivum
47	G965	7244366	1.20E-29	Mentha x piperita
49	G793	6976712	3.60E-43	Lycopersicon esculentum
49	G793	7924288	2.00E-41	Glycine max
49	G793	7614163	3.90E-34	Lotus japonicus
49	G793	9198126	5.70E-23	Medicago truncatula
49	G793	5804781	1.10E-22	Nicotiana tabacum
49	G793	7535578	1.60E-14	Sorghum bicolor
49	G793	427677	6.10E-14	Oryza sativa
49	G793	5915205	2.90E-10	Zea mays
49	G793	9249806	4.20E-10	Solanum tuberosum
49	G793	7624745	1.30E-09	Gossypium arboreum
51	G764	4384535	7.00E-70	Lycopersicon esculentum
51	G764	5049217	1.80E-65	Gossypium hirsutum
51	G764	6454868	1.90E-64	Glycine max
51	G764	6066594	5.20E-59	Petunia x hybrida
51	G764	4218536	2.30E-52	Triticum sp.
51	G764	6732159	2.30E-52	Triticum monococcum
51	G764	9361647	7.50E-52	Triticum aestivum
51	G764	4977542	4.10E-49	Oryza sativa
51	G764	6799764	4.40E-49	Medicago truncatula
51	G764	9296257	1.00E-48	Sorghum bicolor

Figure 3F

SEQ ID No.	GID	Genbank NID	P-value	Species
53	G350	439492	5.20E-53	Petunia x hybrida
53	G350	7228328	8.90E-51	Medicago sativa
53	G350	4666359	3.10E-48	Datisca glomerata
53	G350	1763062	8.30E-48	Glycine max
53	G350	7626808	9.10E-44	Gossypium arboreum
53	G350	7206360	2.20E-43	Medicago truncatula
53	G350	2981168	2.10E-38	Nicotiana tabacum
53	G350	7322653	2.00E-37	Lycopersicon hirsutum
53	G350	5276755	2.40E-37	Lycopersicon esculentum
53	G350	2058503	1.10E-31	Brassica rapa
55	G986	6472584	1.00E-34	Nicotiana tabacum
55	G986	8440065	8.80E-33	Gossypium hirsutum
55	G986	4385167	1.50E-32	Lycopersicon esculentum
55	G986	8205146	5.50E-30	Oryza sativa
55	G986	5820418	8.80E-26	Glycine max
55	G986	1159878	2.30E-23	Avena fatua
55	G986	9250698	4.60E-22	Solanum tuberosum
55	G986	9413507	7.90E-21	Triticum aestivum
55	G986	7748539	2.30E-20	Lotus japonicus
55	G986	9199620	1.30E-16	Medicago truncatula
57	G1349	8904043	1.50E-47	Hordeum vulgare
57	G1349	7244408	2.40E-47	Mentha x piperita
57	G1349	8286031	3.60E-46	Glycine max
57	G1349	9298238	9.10E-36	Sorghum bicolor
57	G1349	7628908	4.70E-34	Gossypium arboreum
57	G1349	5046180	1.50E-33	Gossypium hirsutum
57	G1349	5888938	1.30E-30	Lycopersicon esculentum
57	G1349	5043924	6.20E-30	Pinus taeda
57	G1349	8845076	4.40E-29	Triticum aestivum
57	G1349	7678652	4.20E-27	Lotus japonicus

MBI15 Sequence Listing.ST25 SEQUENCE LISTING

<pre><110> Heard, Jacqueline Keddie, James Yu, Guo-Liang Ratcliffe, Oliver Creelman, Robert Jiang, Cai-zhong Pineda, Omaira Reuber, Lynne Adam, Luc</pre>	
<120> Pathogen Resistance Genes	
<130> MBI-0015	
<150> 60/166,228 <151> 1999-11-17	
<150> Plant Trait Modification III <151> 2000-08-22	
<150> 60/197,899 <151> 2000-04-17	
<160> 58	
<170> PatentIn version 3.0	
<210> 1 <211> 1187 <212> DNA <213> Arabidopsis thaliana	
<220> <221> CDS <222> (50)(1096) <223> G188	
<400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser 1	58 r
<400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser	
<pre><400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc</pre>	r
<pre> <400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc</pre>	106
<pre> <400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc</pre>	106 154
<pre> <400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc</pre>	106 154 202
<pre><400> 1 ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc</pre>	106 154 202 250
ctctcaccaa cataatcaaa gaagctttcc tcacgaattc aagatcgcc atg tcc tcc Met Ser Ser gag gat tgg gat ctc ttc gcc gtc gtc aga agc tgc agc tct tct gtt Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser Ser Ser Val 5 tcc acc acc act aat tct tgt gct ggt cat gaa gac gac ata gga aac tgt Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile Gly Asn Cys 20 aaa caa caa caa gat cct cct cct cct cct ctg ttt caa gct tct tct Lys Gln Gln Gln Asp Pro Pro Pro Pro Pro Leu Phe Gln Ala Ser Ser tct tgc aac gag tta caa gat tct tgc aaa cca ttt tta ccc gtt act Ser Cys Asn Glu Leu Gln Asp Ser Cys Lys Pro Phe Leu Pro Val Thr 65 act act act act act act act tgg tct cct cct cct cct cta ctt cct cct Thr Thr Thr Thr Thr Thr Thr Trp Ser Pro Pro Pro Leu Leu Pro Pro Pro 70 aaa gcc tca tca cca tct ccc aat atc tta cta aaa caa gaa caa gta Lys Ala Ser Ser Pro Ser Pro Asn Ile Leu Leu Lys Gln Glu Gln Val	106 154 202 250 298

MBI15 Sequence Listing.ST25	
cag ctt ctt caa caa tcc caa cct ccc ctt cga tct aga aaa aga Gln Leu Leu Gln Gln Ser Gln Pro Pro Leu Arg Ser Arg Lys Arg 135 140 145	490
aag aat cag caa aaa aga acc ata tgt cat gta acg caa gag aat ctt Lys Asn Gln Gln Lys Arg Thr Ile Cys His Val Thr Gln Glu Asn Leu 150 155	538
tet tet gat ttg tgg get tgg egt aaa tae ggt caa aaa eee ate aaa Ser Ser Asp Leu Trp Ala Trp Arg Lys Tyr Gly Gln Lys Pro Ile Lys 165 170 175	586
ggc tct cct tat cca agg aat tat tac aga tgt agt agc tca aaa gga Gly Ser Pro Tyr Pro Arg Asn Tyr Tyr Arg Cys Ser Ser Ser Lys Gly 180 185	634
tgt tta gca cga aaa caa gtt gaa aga agt aat tta gat cct aat atc Cys Leu Ala Arg Lys Gln Val Glu Arg Ser Asn Leu Asp Pro Asn Ile 200 205 210	682
ttc atc gtt act tac acc gga gaa cac act cat cca cgt cct act cac Phe Ile Val Thr Tyr Thr Gly Glu His Thr His Pro Arg Pro Thr His 225	730
cgg aac tot oto goo gga agt act ogt aac aaa tot cag oco gtt aac Arg Asn Ser Leu Ala Gly Ser Thr Arg Asn Lys Ser Gln Pro Val Asn 230 235 240	778
ccg gtt cct aaa ccg gac aca tct cct tta tcg gat aca gta aaa gaa Pro Val Pro Lys Pro Asp Thr Ser Pro Leu Ser Asp Thr Val Lys Glu 245 250 255	826
gag att cat ctt tct ccg acg aca ccg ttg aaa gga aac gat gac gtt Glu Ile His Leu Ser Pro Thr Thr Pro Leu Lys Gly Asn Asp Asp Val 270 275	874
caa gaa acg aat gga gat gaa gat atg gtt ggt caa gaa gtc aac atg Gln Glu Thr Asn Gly Asp Glu Asp Met Val Gly Gln Glu Val Asn Met 280 285 290	922
gaa gag gaa gag gaa gaa gaa gtg gaa gaa	970
gaa gat gat gac gtg gat gat ctt ttg ata cca aat tta gcg gtg Glu Asp Asp Asp Val Asp Asp Leu Leu Ile Pro Asn Leu Ala Val 310 315 320	1018
aga gat cga gat gat ttg ttc ttc gct gga agt ttt cca tct tgg tcc Arg Asp Arg Asp Asp Leu Phe Phe Ala Gly Ser Phe Pro Ser Trp Ser 325 330 335	1066
gcc gga tcc gcc ggt gac ggt gga tga tgaaaacgaa taaaatctca Ala Gly Ser Ala Gly Asp Gly Gly Gly 340 345	1116
atttacaatt tacaaaaaga aaaaagtcag tttttaatta ttatttttgt ttgttaaaac	1176
ttgacattta t	1187
<210> 2 <211> 348 <212> PRT <213> Arabidopsis thaliana	
<pre><400> 2 Met Ser Ser Glu Asp Trp Asp Leu Phe Ala Val Val Arg Ser Cys Ser 1</pre>	
Ser Ser Val Ser Thr Thr Asn Ser Cys Ala Gly His Glu Asp Asp Ile	

MBI15 Sequence Listing.ST25 25 30

20

Gly Asn Cys Lys Gln Gln Gln Asp Pro Pro Pro Pro Pro Leu Phe Gln 35 40 45

Ala Ser Ser Cys Asn Glu Leu Gln Asp Ser Cys Lys Pro Phe Leu 50 55 60

Pro Val Thr Thr Thr Thr Thr Thr Thr Trp Ser Pro Pro Pro Leu Leu 65 70 70 80

Pro Pro Pro Lys Ala Ser Ser Pro Ser Pro Asn Ile Leu Leu Lys Gln 85 90 95

Glu Gln Val Leu Leu Glu Ser Gln Asp Gln Lys Pro Pro Leu Ser Val 100 105 110

Arg Val Phe Pro Pro Ser Thr Ser Ser Ser Val Phe Val Phe Arg Gly
115 120 125

Gln Arg Asp Gln Leu Leu Gln Gln Gln Ser Gln Pro Pro Leu Arg Ser 130 135 140

Arg Lys Arg Lys Asn Gln Gln Lys Arg Thr Ile Cys His Val Thr Gln 145 150 155 160

Glu Asn Leu Ser Ser Asp Leu Trp Ala Trp Arg Lys Tyr Gly Gln Lys 165 170 175

Pro Ile Lys Gly Ser Pro Tyr Pro Arg Asn Tyr Tyr Arg Cys Ser Ser 180 185 190

Ser Lys Gly Cys Leu Ala Arg Lys Gln Val Glu Arg Ser Asn Leu Asp 195 200 205

Pro Asn Ile Phe Ile Val Thr Tyr Thr Gly Glu His Thr His Pro Arg 210 215 220

Pro Thr His Arg Asn Ser Leu Ala Gly Ser Thr Arg Asn Lys Ser Gln 225 230 235 240

Pro Val Asn Pro Val Pro Lys Pro Asp Thr Ser Pro Leu Ser Asp Thr 245 250 255

Val Lys Glu Glu Ile His Leu Ser Pro Thr Thr Pro Leu Lys Gly Asn 260 265 270

Asp Asp Val Glu Glu Thr Asn Gly Asp Glu Asp Met Val Gly Glu Glu 275 280 285

Glu Glu Glu Glu Asp Asp Asp Asp Val Asp Asp Leu Leu Ile Pro Asn 305 310 315 320

MBI15 Sequence Listing.ST25

Leu Ala Val Arg Asp Arg Asp Leu Phe Phe Ala Gly Ser Phe Pro 335

330

335

Ser Trp Ser Ala Gly Ser Ala Gly Asp Gly Gly Gly 345

<210> 3	
<210> 3 <211> 1431 <212> DNA <213> Arabidopsis thaliana	
<220> <221> CDS <222> (129)(1211) <223> G616	
<400> 3 ctttaaatcc caaaccaacc ctaaagtttt gatttttaat tttgggggta accaaaaaaa	60
aaacaaaacc ctaatttttt ttctttagtg atgagattat tggtgatgat gaaatgattg	120
gagatcta atg aag aat aac aac aat ggc gac gtt gtg gat aac gaa gtg Met Lys Asn Asn Asn Gly Asp Val Val Asp Asn Glu Val 1 5	170
aac aac cgg tta agc cgg tgg cat cac aat tct tcc cgg ata att agg Asn Asn Arg Leu Ser Arg Trp His His Asn Ser Ser Arg Ile Ile Arg 30	218
gtt tca cga gct tcc ggt ggt aaa gat cga cac agc aaa gtc ttg act Val Ser Arg Ala Ser Gly Gly Lys Asp Arg His Ser Lys Val Leu Thr 35 40 45	266
tct aaa gga cca cgt gac cgt cgt gtc cgg tta tca gtc tcc acc gct Ser Lys Gly Pro Arg Asp Arg Arg Val Arg Leu Ser Val Ser Thr Ala 50 55 60	314
ctt caa ttc tat gat ctt caa gat cgg tta ggt tat gat caa cct agc Leu Gln Phe Tyr Asp Leu Gln Asp Arg Leu Gly Tyr Asp Gln Pro Ser 65 70 75	362
aaa gct gtt gaa tgg tta atc aaa gct gct gaa gat tca atc tct gag Lys Ala Val Glu Trp Leu Ile Lys Ala Ala Glu Asp Ser Ile Ser Glu 80 85	410
ctt cct tca ctc aac aac act cat ttt ccg acc gat gac gag aat cac Leu Pro Ser Leu Asn Asn Thr His Phe Pro Thr Asp Asp Glu Asn His 100 105 110	458
cag aat cag aca tta aca aca gtt gct gct aat tcc ttg tct aaa tct Gln Asn Gln Thr Leu Thr Thr Val Ala Ala Asn Ser Leu Ser Lys Ser 115	506
gct tgt agt agc aat tca gac acg agc aag aac tct tct ggt ttg tct Ala Cys Ser Ser Asn Ser Asp Thr Ser Lys Asn Ser Ser Gly Leu Ser 130 135	554
tta tca aga tcg gag ctt aga gat aaa gct aga gag agg gct aga gag Leu Ser Arg Ser Glu Leu Arg Asp Lys Ala Arg Glu Arg Ala Arg Glu 145 150 155	602
aga aca gct aaa gag acc aag gaa aga gat cat aac cac act tcg ttt Arg Thr Ala Lys Glu Thr Lys Glu Arg Asp His Asn His Thr Ser Phe 160 165 170	650
acg gat ttg tta aat tcc ggt tca gat ccg gtt aac tca aac cgg caa Thr Asp Leu Leu Asn Ser Gly Ser Asp Pro Val Asn Ser Asn Arg Gln 180 185 190	698
tgg atg gct tca gct cct tct tca tct cca atg gag tat ttc agt tcg Trp Met Ala Ser Ala Pro Ser Ser Ser Pro Met Glu Tyr Phe Ser Ser Trp Met Ala Ser Ala Pro Ser Ser Ser 200	746
Page 4	

MBI15 Sequence Listing.ST25

ggt tta att ctc ggg tcg ggt caa caa acc cat ttc cct att tca aca Gly Leu Ile Leu Gly Ser Gly Gln Gln Thr His Phe Pro Ile Ser Thr 210 215 220	794
aat tot cat cot tto toa toa ato too gat cat cat cat cat cot Asn Ser His Pro Phe Ser Ser Ile Ser Asp His His His His Pro 225 230 235	842
cat cat cag cat caa gag ttt tca ttc gtt ccc gac cat ttg ata tca His His Gln His Gln Glu Phe Ser Phe Val Pro Asp His Leu Ile Ser 240 245 250	890
ccg gca gaa tcc aac ggc gga gca ttc aat ctt gat ttt aat atg tca Pro Ala Glu Ser Asn Gly Gly Ala Phe Asn Leu Asp Phe Asn Met Ser 255 260 265 270	938
aca ccc tcc ggc gcc gga gct gcc gtc tcc gcc gca tca ggt ggc ggc Thr Pro Ser Gly Ala Gly Ala Ala Val Ser Ala Ala Ser Gly Gly 275 280 285	986
ttc agt ggt ttc aac agg ggg acc ctt cag tcc aat tca aca aat cag Phe Ser Gly Phe Asn Arg Gly Thr Leu Gln Ser Asn Ser Thr Asn Gln 290 295 300	1034
cat cag tca ttc ctc gct aat cta cag agg ttt cca aca tca gaa agt His Gln Ser Phe Leu Ala Asn Leu Gln Arg Phe Pro Thr Ser Glu Ser 305 310 315	1082
gga gga ggt cca cag ttc tta ttc ggt gca ctg cct gca gag aat cac Gly Gly Pro Gln Phe Leu Phe Gly Ala Leu Pro Ala Glu Asn His 320 325 330	1130
cac cac aat cac cag ttt cag ctt tac tat gaa aat gga tgc aga aac His His Asn His Gln Phe Gln Leu Tyr Tyr Glu Asn Gly Cys Arg Asn 335 340 345 350	1178
tca tca gaa cat aag ggt aaa ggc aag aac tga tgatattaat tattgcatct Ser Ser Glu His Lys Gly Lys Gly Lys Asn 355 360	1231
ttggttttgt tcaaatgctc attttgtatg tttatctttg gtttatttca aaacaaatgt	1291
taatetettt egttgtetga tgtgtgttag ggttttgttt tatgtattga gggtetttgg	1351
aaatettttt geattgtget tgtaatgttg tatttgtgat aatageattt tgtttgtgag	1411
ttaaaaaaaa aaaaaaaaaa	1431

:210> 4

<211> 360 <212> PRT

<213> Arabidopsis thaliana

<400> 4

Met Lys Asn Asn Asn Asn Gly Asp Val Val Asp Asn Glu Val Asn Asn 1 5 10 15

Arg Ala Ser Gly Gly Lys Asp Arg His Ser Lys Val Leu Thr Ser Lys 35 40 45

Gly Pro Arg Asp Arg Val Arg Leu Ser Val Ser Thr Ala Leu Gln 50 55

Phe Tyr Asp Leu Gln Asp Arg Leu Gly Tyr Asp Gln Pro Ser Lys Ala 65 70 75 80

MBI15 Sequence Listing.ST25

- Val Glu Trp Leu Ile Lys Ala Ala Glu Asp Ser Ile Ser Glu Leu Pro 85 90 95
- Ser Leu Asn Asn Thr His Phe Pro Thr Asp Asp Glu Asn His Gln Asn 100 105 110
- Gln Thr Leu Thr Thr Val Ala Ala Asn Ser Leu Ser Lys Ser Ala Cys 115 120 125
- Ser Ser Asn Ser Asp Thr Ser Lys Asn Ser Ser Gly Leu Ser Leu Ser 130 135 140
- Arg Ser Glu Leu Arg Asp Lys Ala Arg Glu Arg Ala Arg Glu Arg Thr 145 150 155 160
- Ala Lys Glu Thr Lys Glu Arg Asp His Asn His Thr Ser Phe Thr Asp 165 170 175
- Leu Leu Asn Ser Gly Ser Asp Pro Val Asn Ser Asn Arg Gln Trp Met 180 185 190
- Ala Ser Ala Pro Ser Ser Ser Pro Met Glu Tyr Phe Ser Ser Gly Leu 195 200 205
- Ile Leu Gly Ser Gly Gln Gln Thr His Phe Pro Ile Ser Thr Asn Ser 210 215 220
- His Pro Phe Ser Ser Ile Ser Asp His His His His Bro His His 225 230 235 240
- Gln His Gln Glu Phe Ser Phe Val Pro Asp His Leu Ile Ser Pro Ala 245 250 250
- Glu Ser Asn Gly Gly Ala Phe Asn Leu Asp Phe Asn Met Ser Thr Pro 260 265
- Ser Gly Ala Gly Ala Ala Val Ser Ala Ala Ser Gly Gly Gly Phe Ser 275 280 285
- Gly Phe Asn Arg Gly Thr Leu Gln Ser Asn Ser Thr Asn Gln His Gln 290 295 300
- Ser Phe Leu Ala Asn Leu Gln Arg Phe Pro Thr Ser Glu Ser Gly Gly 305 310 315
- Gly Pro Gln Phe Leu Phe Gly Ala Leu Pro Ala Glu Asn His His His 325 330 335
- Asn His Gln Phe Gln Leu Tyr Tyr Glu Asn Gly Cys Arg Asn Ser Ser 340 345 350
- Glu His Lys Gly Lys Gly Lys Asn 355 Asn

<210> 5

WO 01/35726 PCT/US00/31418 \

MBI15 Sequence Listing.ST25

<211 <212 <213	> E	.055 NA Lrabi	.dops	is t	hali	ana	MBII	.5 56	equei	ice i	11561	ing.s	,123			
<220 <221 <222 <223	> (:DS (70). ;19	. (81	6)												
<400 ataa	> 5 aggo	at t	tcag	ctcc	a co	gtag	gaaa	ctt	tctc	ttg	aaag	aaac	cc a	cago	aacaa	60
acag	agaa	a at Me	g tg et Cy	rs Gl	y Gl	t go y Al 5	t at a Il	t at e Il	t to .e Se	c ga er As	t ta p Ty 10	r Al	c co a Pr	t ct o Le	c gtc u Val	111
acc Thr 15	aag Lys	gcc Ala	aag Lys	ggc Gly	cgt Arg 20	aaa Lys	ctc Leu	acg Thr	gct Ala	gag Glu 25	gaa Glu	ctc Leu	tgg Trp	tca Ser	gag Glu 30	159
ctc Leu	gat Asp	gct Ala	tcc Ser	gcc Ala 35	gcc Ala	gac Asp	gac Asp	ttc Phe	tgg Trp 40	ggt Gly	ttc Phe	tat Tyr	tcc Ser	acc Thr 45	tcc Ser	207
aaa Lys	ctc Leu	cat His	ccc Pro 50	acc Thr	aac Asn	caa Gln	gtt Val	aac Asn 55	gtg Val	aaa Lys	gag Glu	gag Glu	gca Ala 60	gtg Val	aag Lys	255
aag Lys	gag Glu	cag Gln 65	gca Ala	aca Thr	gag Glu	ccg Pro	999 Gly 70	aaa Lys	cgg Arg	agg Arg	aag Lys	agg Arg 75	aag Lys	aat Asn	gtt Val	303
tat Tyr	aga Arg 80	999 Gly	ata Ile	cgt Arg	aag Lys	cgt Arg 85	cca Pro	tgg Trp	gga Gly	aaa Lys	tgg Trp 90	gcg Ala	gct Ala	gag Glu	att Ile	351
cga Arg 95	gat Asp	cca Pro	cga Arg	aaa Lys	ggt Gly 100	gtt Val	aga Arg	gtt Val	tgg Trp	ctt Leu 105	ggt Gly	acg Thr	ttc Phe	aac Asn	acg Thr 110	399
gcg Ala	gag Glu	gaa Glu	gct Ala	gcc Ala 115	atg Met	gct Ala	tat Tyr	gat Asp	gtt Val 120	gcg Ala	gcc Ala	aag Lys	cag Gln	atc Ile 125	cgt Arg	447
ggt Gly	gat Asp	aaa Lys	gcc Ala 130	aag Lys	ctc Leu	aac Asn	ttc Phe	cca Pro 135	Asp	ctg Leu	cac His	cat His	cct Pro 140	cct Pro	cct . Pro	· 495
cct Pro	aat Asn	tat Tyr 145	act Thr	cct Pro	ccg Pro	ccg Pro	tca Ser 150	tcg Ser	cca Pro	cga Arg	tca Ser	acc Thr 155	gat Asp	cag Gln	cct Pro	543
ccg Pro	gcg Ala 160	aag Lys	aag Lys	gtc Val	tgc Cys	gtt Val 165	gtc Val	tct Ser	cag Gln	agt Ser	gag Glu 170	agc Ser	gag Glu	tta Leu	agt Ser	591
cag Gln 175	ccg Pro	agt Ser	ttc Phe	ccg Pro	gtg Val 180	gag Glu	tgt Cys	ata Ile	gga Gly	ttt Phe 185	gga Gly	aat Asn	999 Gly	gac Asp	gag Glu 190	639
ttt Phe	cag Gln	aac Asn	ctg Leu	agt Ser 195	tac Tyr	gga Gly	ttt Phe	gag Glu	ccg Pro 200	gat Asp	tat Tyr	gat Asp	ctg Leu	aaa Lys 205	cag Gln	687
cag Gln	ata Ile	tcg Ser	agc Ser 210	ttg Leu	gaa Glu	tcg Ser	ttc Phe	ctt Leu 215	gag Glu	ctg Leu	gac Asp	ggt Gly	aac Asn 220	acg Thr	gcg Ala	735
gag Glu	caa Gln	ccg Pro 225	Ser	cag Gln	ctt Leu	gat Asp	gag Glu 230	tcc Ser	gtt Val	tcc Ser	gag Glu	gtg Val 235	gat Asp	atg Met	tgg Trp	783
atg Met	ctt Leu	gat Asp	gat Asp	gtc Val	att Ile	gcg Ala	tcg Ser	tat Tyr	gag Glu	taa	aag	aaaa	aaa	ataa	gtttaa	836

	MBI15	Sequence	Listing.ST25
245		*	_

aaaaagttaa ataaagtctg taatatata gtaaccgccg ttacttttaa aaggtttta 896
ccgtcgcatt ggactgctga tgatgtctgt tgtgtaatgt gtagaatgtg accaaatgga 956
cgttatatta cggtttgtgg tattattagt ttcttagatg gaaaaactta catgtgtaaa 1016
taagatttgt aatgtaagac gaagtactta taacttctt 1055

<210> 6 <211> 248 <212> PRT

240

<213> Arabidopsis thaliana

<400> 6

Met Cys Gly Gly Ala Ile Ile Ser Asp Tyr Ala Pro Leu Val Thr Lys

1 10 15

Ala Lys Gly Arg Lys Leu Thr Ala Glu Glu Leu Trp Ser Glu Leu Asp 20 25 30

Ala Ser Ala Ala Asp Asp Phe Trp Gly Phe Trr Ser Thr Ser Lys Leu 35 40 45

His Pro Thr Asn Gln Val Asn Val Lys Glu Glu Ala Val Lys Lys Glu 50 55 60

Gln Ala Thr Glu Pro Gly Lys Arg Arg Lys Arg Lys Asn Val Tyr Arg 65 70 75 80

Gly Ile Arg Lys Arg Pro Trp Gly Lys Trp Ala Ala Glu Ile Arg Asp 85 90 95

Pro Arg Lys Gly Val Arg Val Trp Leu Gly Thr Phe Asn Thr Ala Glu 100 105 110

Glu Ala Ala Met Ala Tyr Asp Val Ala Ala Lys Gln Ile Arg Gly Asp 115 120 125

Lys Ala Lys Leu Asn Phe Pro Asp Leu His His Pro Pro Pro Pro Asn 130 135 140

Tyr Thr Pro Pro Pro Ser Ser Pro Arg Ser Thr Asp Gln Pro Pro Ala 145 150 155 160

Lys Lys Val Cys Val Val Ser Gln Ser Glu Ser Glu Leu Ser Gln Pro 165 170 175

Ser Phe Pro Val Glu Cys Ile Gly Phe Gly Asn Gly Asp Glu Phe Gln 180 185 190

Asn Leu Ser Tyr Gly Phe Glu Pro Asp Tyr Asp Leu Lys Gln Gln Ile 195 200 205

Ser Ser Leu Glu Ser Phe Leu Glu Leu Asp Gly Asn Thr Ala Glu Gln 210 215 220

Pro Ser Gln Leu Asp Glu Ser Val Ser Glu Val Asp Met Trp Met Leu Page 8

MBI15 Sequence Listing.ST25

225 230 235 240

Asp Asp Val Ile Ala Ser Tyr Glu 245

<210> <211> 1857 <212> DNA Arabidopsis thaliana <221> <222> (458) . . (1663) G261 <223> <400> 7 60 gtttaggttc gagaagcaga gagggttcga gaagctaata agggtttctt ctttttgatt ttaatgctaa aagggttcta gattcgttga attttacaag ggttttaggg gttcttagaa 120 gcttttgctt gattgtcttt tatttagaaa cagtggtgag tttttagtct ttcactttgt 180 tcaagttcga agctttttt ggagggaatt ttgggcttct gattttgatc gaaacttact 240 gatagtaagt tetttgagte eteettaact gtagtttetg tgtaetgaag ttattgaatt 300 gaaagttttt atcttttttg gttattgaaa ctttcatagt ttgatcaaaa gagtctcttg ctctgttttt ggctctgttt ttgtgagtgt gattgtaagc tttgttgtga gtagattgaa 420 tcaaggagtg tgagagttgt taaaagtgtt ttcagag atg gat gag aat aat cat Met Asp Glu Asn Asn His 475 gga gtt tca tca agc tca ctt cca cct ttc ctc acc aaa aca tat gag Gly Val Ser Ser Ser Leu Pro Pro Phe Leu Thr Lys Thr Tyr Glu 523 atg gtt gat gat tot toa too gat tot atc gtc tot tgg agt cag agc Met Val Asp Asp Ser Ser Ser Asp Ser Ile Val Ser Trp Ser Gln Ser 571 aat aag agt ttc atc gtt tgg aat ccg ccg gag ttt tct aga gat ctt Asn Lys Ser Phe Ile Val Trp Asn Pro Pro Glu Phe Ser Arg Asp Leu 619 ctt ccg aga ttc ttc aag cac aat aac ttc tct agc ttt atc cgc cag Leu Pro Arg Phe Phe Lys His Asn Asn Phe Ser Ser Phe Ile Arg Gln 667 ctt aac aca tat ggt ttt aga aaa gct gat cct gag caa tgg gaa ttt Leu Asn Thr Tyr Gly Phe Arg Lys Ala Asp Pro Glu Gln Trp Glu Phe 715 gcg aat gat gat ttt gtg aga ggt caa cct cat ctt atg aag aac att Ala Asn Asp Asp Phe Val Arg Gly Gln Pro His Leu Met Lys Asn Ile 763 cat aga cgc aaa cca gtt cat agc cac tct tta ccg aat ctt caa gct His Arg Arg Lys Pro Val His Ser His Ser Leu Pro Asn Leu Gln Ala 811 110 cag tta aac ccg ttg acg gat tca gaa cga gtg aga atg aat aat cag Gln Leu Asn Pro Leu Thr Asp Ser Glu Arg Val Arg Met Asn Asn Gln 859 att gag aga ttg aca aaa gag aaa gaa gga ttg ctt gaa gag tta cat lle Glu Arg Leu Thr Lys Glu Lys Glu Gly Leu Leu Glu Glu Leu His 907 145 aaa caa gac gag gaa cga gtg ttt gag atg caa gtg aaa gaa Ctt Lys Gln Asp Glu Glu Arg Glu Val Phe Glu Met Gln Val Lys Glu Leu 955

. 155	MBI15 Sequence 1	Listing.ST25 165	
aaa gaa cga tta caa ca	ac atg gag aag cgt cag	aaa aca atg gtt tcg	1003
Lys Glu Arg Leu Gln Hi	Ls Met Glu Lys Arg Gln	Lys Thr Met Val Ser	
170	175	180	
ttt gtt tct caa gta tt	ng gaa aag cca ggg ctt	gct ttg aac cta tcg	1051
Phe Val Ser Gln Val Le	eu Glu Lys Pro Gly Leu	Ala Leu Asn Leu Ser	
185	190	195	
ccg tgt gtt ccc gaa ac	ca aac gag agg aaa aga	agg ttc cct agg atc	1099
Pro Cys Val Pro Glu Th	nr Asn Glu Arg Lys Arg	Arg Phe Pro Arg Ile	
200	205	210	
gag ttc ttt ccc gat ga Glu Phe Phe Pro Asp Gl 215	lu Pro Met Leu Glu Glu	aac aaa act tgt gtt Asn Lys Thr Cys Val 230	1147
gtt gtg aga gag gaa gg	gt tot aca ago cot tot	tca cac aca aga gag	1195
Val Val Arg Glu Glu Gl	ly Ser Thr Ser Pro Ser	Ser His Thr Arg Glu	
235	240	245	
cat caa gtg gaa cag tt	ta gag tca tcg ata gcg	att tgg gag aat ctt	1243
His Gln Val Glu Gln Le	eu Glu Ser Ser Ile Ala	Ile Trp Glu Asn Leu	
250	255	260	
gta tcg gat tct tgt ga	ag agt atg tta caa tca	aga agt atg atg aca	1291
Val Ser Asp Ser Cys Gl	lu Ser Met Leu Gln Ser	Arg Ser Met Met Thr	
265	270	275	
ctt gat gtg gat gaa to	ca tct act ttt cca gag	agc cct cct ctt tct	1339
Leu Asp Val Asp Glu Se	er Ser Thr Phe Pro Glu	Ser Pro Pro Leu Ser	
280	285	290	
tgc ata cag tta agt gt Cys Ile Gln Leu Ser Va 295 30	al Asp Ser Arg Leu Lys	tct cct cct tct cca Ser Pro Pro Ser Pro 310	1387
agg atc atc gat atg aa	ac tgt gag ccc gat ggt	tcg aaa gaa cag aac	1435
Arg Ile Ile Asp Met As	sn Cys Glu Pro Asp Gly	Ser Lys Glu Gln Asn	
315	320	325	
act gtt gct gct cct cc	ct cct cct cca gta gca	gga gcg aat gat ggc	1483
Thr Val Ala Ala Pro Pr	ro Pro Pro Pro Val Ala	Gly Ala Asn Asp Gly	
330	335	340	
ttc tgg cag cag ttt tt	tc tca gag aat cct ggc	tca acc gag caa cgg	1531
Phe Trp Gln Gln Phe Ph	he Ser Glu Asn Pro Gly	Ser Thr Glu Gln Arg	
345	350	355	
gaa gtt caa tta gag ag	gg aaa gac gat aaa gat	aaa gcc gga gta cgt	1579
Glu Val Gln Leu Glu Ar	rg Lys Asp Asp Lys Asp	Lys Ala Gly Val Arg	
360	365	370	
Thr Glu Lys Cys Trp Tr	gg aat tcg aga aat gtt rp Asn Ser Arg Asn Val 80 385	aat gca att aca gaa Asn Ala Ile Thr Glu 390	1627
cag ctt gga cat ctg ac Gln Leu Gly His Leu Th 395	ct tct tca gag aga agt hr Ser Ser Glu Arg Ser 400	tga tatgtcaaag	1673
attaaatttc tagtctgttt	tagttacttg taaaataggg	tttctcagtt ttattgtttt	1733
		acttgtatga tctttcccag	1793
ttcattgtag cagacttcaa	tggtaatgat, aagctagagc	ttatggatag tattcataaa	1853
aaaa			1857

<210> 8 <211> 401 <212> PRT <213> Arabidopsis thaliana

MBI15 Sequence Listing.ST25

<400> 8

Met Asp Glu Asn Asn His Gly Val Ser Ser Ser Leu Pro Pro Phe

Leu Thr Lys Thr Tyr Glu Met Val Asp Asp Ser Ser Ser Asp Ser Ile 20 25 30

Val Ser Trp Ser Gln Ser Asn Lys Ser Phe Ile Val Trp Asn Pro Pro 35 40 45

Glu Phe Ser Arg Asp Leu Leu Pro Arg Phe Phe Lys His Asn Asn Phe 50 55 60

Ser Ser Phe Ile Arg Gln Leu Asn Thr Tyr Gly Phe Arg Lys Ala Asp 65 70 75 80

Pro Glu Gln Trp Glu Phe Ala Asn Asp Asp Phe Val Arg Gly Gln Pro

His Leu Met Lys Asn Ile His Arg Arg Lys Pro Val His Ser His Ser 100 105 110

Leu Pro Asn Leu Gln Ala Gln Leu Asn Pro Leu Thr Asp Ser Glu Arg 115 120 125

Val Arg Met Asn Asn Gln Ile Glu Arg Leu Thr Lys Glu Lys Glu Gly 130 135 140

Leu Leu Glu Glu Leu His Lys Gln Asp Glu Glu Arg Glu Val Phe Glu 145 150 155 160

Met Gln Val Lys Glu Leu Lys Glu Arg Leu Gln His Met Glu Lys Arg 165 170 175

Gln Lys Thr Met Val Ser Phe Val Ser Gln Val Leu Glu Lys Pro Gly 180 185 190

Leu Ala Leu Asn Leu Ser Pro Cys Val Pro Glu Thr Asn Glu Arg Lys 195 200 205

Arg Arg Phe Pro Arg Ile Glu Phe Phe Pro Asp Glu Pro Met Leu Glu 210 215 220

Glu Asn Lys Thr Cys Val Val Val Arg Glu Glu Gly Ser Thr Ser Pro 225 230 235 240

Ser Ser His Thr Arg Glu His Gln Val Glu Gln Leu Glu Ser Ser Ile 245 250 255

Ala Ile Trp Glu Asn Leu Val Ser Asp Ser Cys Glu Ser Met Leu Gln 260 265 270

Ser Arg Ser Met Met Thr Leu Asp Val Asp Glu Ser Ser Thr Phe Pro 275 280 285

MBI15 Sequence Listing.ST25 Glu Ser Pro Pro Leu Ser Cys Ile Gln Leu Ser Val Asp Ser Arg Leu 295 Lys Ser Pro Pro Ser Pro Arg Ile Ile Asp Met Asn Cys Glu Pro Asp Gly Ser Lys Glu Gln Asn Thr Val Ala Ala Pro Pro Pro Pro Pro Val Ala Gly Ala Asn Asp Gly Phe Trp Gln Gln Phe Phe Ser Glu Asn Pro Gly Ser Thr Glu Gln Arg Glu Val Gln Leu Glu Arg Lys Asp Asp Lys 360 Asp Lys Ala Gly Val Arg Thr Glu Lys Cys Trp Trp Asn Ser Arg Asn Val Asn Ala Ile Thr Glu Gln Leu Gly His Leu Thr Ser Ser Glu Arg Ser <210> <211> 964 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (63)..(869) <223> G28 <400> 9 60 gaaatctcaa caagaaccaa accaaacaac aaaaaaacat tottaataat tatotttotg tt atg tcg atg acg gcg gat tct caa tct gat tat gct ttt ctt gag Met Ser Met Thr Ala Asp Ser Gln Ser Asp Tyr Ala Phe Leu Glu 1 5 10 107 tcc ata cga cga cac tta cta gga gaa tcg gag ccg ata ctc agt gag Ser Ile Arg Arg His Leu Leu Gly Glu Ser Glu Pro Ile Leu Ser Glu 20 25 30155 tcg aca gcg agt tcg gtt act caa tct tgt gta acc ggt cag agc att Ser Thr Ala Ser Ser Val Thr Gln Ser Cys Val Thr Gly Gln Ser Ile 203 251 aaa ceg gtg tac gga ega aac eet age ttt age aaa etg tat eet tge Lys Pro Val Tyr Gly Arg Asn Pro Ser Phe Ser Lys Leu Tyr Pro Cys ttc acc gag agc tgg gga gat ttg ccg ttg aaa gaa aac gat tct gag Phe Thr Glu Ser Trp Gly Asp Leu Pro Leu Lys Glu Asn Asp Ser Glu 65 70 75 299 gat atg tta gtt tac ggt atc ctc aac gac gcc ttt cac ggc ggt tgg Asp Met Leu Val Tyr Gly Ile Leu Asn Asp Ala Phe His Gly Gly Trp 80 85 90 95347 gag cog tot tot tog tot toe gao gaa gat ogt ago tot tto cog agt 395 Glu Pro Ser Ser Ser Ser Ser Asp Glu Asp Arg Ser Ser Phe Pro Ser 443 gtt aag atc gag act ccg gag agt ttc gcg gcg gtg gat tct gtt ccg

							мот	15 6	00010	222	Liet	ina	CT25			
Val	Lys	Ile	Glu 115	Thr	Pro	Glu		15 S Phe 120							Pro	
gtc Val	aag Lys	aag Lys 130	gag Glu	aag Lys	acg Thr	agt Ser	cct Pro 135	gtt Val	tcg Ser	gcg Ala	gcg Ala	gtg Val 140	acg Thr	gcg Ala	gcg Ala	491
	gga Gly 145															539
	gcg Ala															587
	acg Thr															635
	ttc Phe															683
	aat Asn															731
	ttt Phe 225															779
	gcc Ala															827
	gtt Val												taa			869
ttt	tgati	tt (ctt	gtt	g at	gatt	atat	gat	tctt	caa	aaaa	igaa	gaa d	gtta	aataaa	929
aaaa	attc	gtt (atta	attaa	aa aa	aaaa	aaaa	a aaa	aa							964
<210 <210 <210 <210	1 > 2 2 > 1	10 268 PRT	idops	sis t	:hali	iana										
<40		10	•													
Met 1	Ser	Met	Thr	Ala 5	Авр	Ser	Gln	Ser	Asp 10	Tyr	Ala	Phe	Leu	Glu 15	Ser	
Ile	Arg	Arg	His 20	Leu	Leu	Gly	Glu	Ser 25	Glu	Pro	Ile	Leu	Ser 30	Glu	Ser	
Thr	Ala	Ser 35	Ser	Val	Thr	Gln	Ser 40	Cys	Val	Thr	Gly	Gln 45	Ser	Ile	Lys	
Pro	Val 50	Tyr	Gly	Arg	Asn	Pro 55	Ser	Phe	Ser	Lys	Leu 60	Tyr	Pro	Сув	Phe	
Thr 65	Glu	Ser	Trp	Gly	Asp 70	Leu	Pro	Leu	Lys	Glu 75	Asn	Asp	Ser	Glu	Asp 80	

Page 13

Met Leu Val Tyr Gly Ile Leu Asn Asp Ala Phe His Gly Gly Trp Glu 95

MBI15 Sequence Listing.ST25

Pro Ser Ser Ser Ser Asp Glu Asp Arg Ser Ser Phe Pro Ser Val

105 Lys Ile Glu Thr Pro Glu Ser Phe Ala Ala Val Asp Ser Val Pro Val Lys Lys Glu Lys Thr Ser Pro Val Ser Ala Ala Val Thr Ala Ala Lys 130 135 140 Gly Lys His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Phe Ala 145 150 160 Ala Glu Ile Arg Asp Pro Ala Lys Asn Gly Ala Arg Val Trp Leu Gly Thr Phe Glu Thr Ala Glu Asp Ala Ala Leu Ala Tyr Asp Arg Ala Ala Phe Arg Met Arg Gly Ser Arg Ala Leu Leu Asn Phe Pro Leu Arg Val Asn Ser Gly Glu Pro Asp Pro Val Arg Ile Lys Ser Lys Arg Ser Ser 210 215 220 Phe Ser Ser Ser Asn Glu Asn Gly Ala Pro Lys Lys Arg Arg Thr Val 225 230 235 240 Ala Ala Gly Gly Gly Met Asp Lys Gly Leu Thr Val Lys Cys Glu Val 245 250 255 Val Glu Val Ala Arg Gly Asp Arg Leu Leu Val Leu <210> · 11 <211> 1571 <212> DNA Arabidopsis thaliana <220> CDS <221> <222> (428) .. (1402) <223> G869 <400> 11 aggaacagtg aaaggttcgg ttttttgggt ttcgatctga taatcaacaa gaaaaaaggg 60 tttgatttat gtcggctggg tttgaatcga ctgtgatttt gtctttgatt catatctctt 120 ctccgatttc atcatcatct tccccatcat cgtcgtcttt gaaatcttgt cttctcaacg 180 ctcttcactt ctgctgtaat aagcagaggc ttgttctgga gactccttct ctttccatgc gcttaagacc caaaaggact tgttctagtg ttgaagtctt tggggggtttt cacataaagc 300 agcaaaagtt ttctttttc atagttcgct gagagttttg agttttgata ccaaaaaagt 360 tttgaccttt tagagtgatt ttttgttctt tctgttttct gggtattttt gaggagtggg 420 tttaaca atg gtt gcg att aga aag gaa cag tct ttg agt ggt gtt agt Met Val Ala Ile Arg Lys Glu Gln Ser Leu Ser Gly Val Ser 469 517 ago gag att aag aag aga got aag aga aac act ota tog too ott oot Page 14

Ser 15	Glu	Ile	Lys	Lys	Arg 20	Ala					List Leu			Leu	Pro 30	,
														gat Asp 45		565
														gtt Val		613
														cct Pro		661
														agt Ser		709
														cct Pro		757
														gct Ala 125		805
														act Thr		853
														ctt Leu		901
														gat Asp		949
														gtt Val		997
														aac Asn 205		1045
cct Pro	gat Asp	gac Asp	gtc Val 210	tcg Ser	acc Thr	gtt Val	gct Ala	cca Pro 215	act Thr	gct Ala	cca Pro	act Thr	cca Pro 220	aat Asn	gtt Val	1093
														act Thr		1141
cta Leu	cag Gln 240	atc Ile	cct Pro	gat Asp	ttt Phe	ggt Gly 245	ttc Phe	ttg Leu	gca Ala	gag Glu	gag Glu 250	caa Gln	caa Gln	gac Asp	cta Leu	1189
														ggc Gly		1237
														tta Leu 285		1285
gat Asp	ttc Phe	gac Asp	ttt Phe 290	gcg Ala	gat Asp	gtt Val	gaa Glu	gat Asp 295	ctt Leu	cag Gln	cta Leu	gct Ala	gac Asp 300	tct Ser	agt Ser	1333
														cca Pro		1381

Page 15

		ttt Phe				tag			-		List itgtt	_		gaaga	1
gtgt	ttt	gtt t	ttt	gtt	a to	gcttt	agta	att	taag	gaca	taca	aaaa	gtg (gtgt	tccgg
atte	gtagt	aa g	gatet	taag	ga ca	ataaa	gcc	g ggt	tttg	gcaa	ttag	ggaat	cg a	agttt	taatg
aagt	ttta	agt t	tate	gtttg											
<210 <211 <211 <211	L> :	12 324 PRT Arabi	idops	sis t	:hali	iana									
<400)> :	12													
Met 1	Val	Ala	Ile	Arg 5	Lys	Glu	Gln	Ser	Leu 10	Ser	Gly	Val	Ser	Ser 15	Glu
Ile	Lys	Lys	Arg 20	Ala	Lys	Arg	Asn	Thr 25	Leu	Ser	Ser	Leu	Pro 30	Gln	Glu
Thr	Gln	Pro 35	Leu	Arg	Lys	Val	Arg 40	Ile	Ile	Val	Asn	Asp 45	Pro	Tyr	Ala
Thr	Авр 50	Asp	Ser	Ser	Ser	Asp 55	Glu	Glu	Glu	Leu	Lys 60	Val	Pro	Lys	Pro
Arg 65	Lys	Met	Lys	Arg	Ile 70	Val	Arg	Glu	Ile	Asn 75	Phe	Pro	Ser	Met	Glu 80
Val	Ser	Glu	Gln	Pro 85	Ser	Glu	Ser	Ser	Ser 90	Gln	Asp	Ser	Thr	Lys 95	Thr ·
Asp	Gly	Lys	Ile 100	Ala	Val	Ser	Ala	Ser 105	Pro	Ala	Val	Pro	Arg 110	Lys	Lys
Pro	Val	Gly 115	Val	Arg	Gln	Arg	Lys 120	Trp	Gly	Lys	Trp	Ala 125	Ala	Glu	Ile
Arg	Asp 130	Pro	Ile	Lys	Lys	Thr 135	Arg	Thr	Trp	Leu	Gly 140	Thr	Phe	Asp	Thr
Leu 145	Glu	Glu	Ala	Ala	Lys 150	Ala	Tyr	Asp	Ala	Lys 155	Lys	Leu	Glu	Phe	Asp 160
Ala	Ile	Val	Ala	Gly 165	Asn	Val	Ser	Thr	Thr 170	Lys	Arg	Asp	Val	Ser 175	Ser
Ser	Glu	Thr	Ser 180	Gln	Cys	Ser	Arg	Ser 185	Ser	Pro	Val	Val	Pro 190	Val	Glu
Gln	Asp	Авр 195	Thr	Ser	Ala	Ser	Ala 200	Leu	Thr	Сув	Val	Asn 205	Asn	Pro	Asp
Asp	Val 210	Ser	Thr	Val	Ala	Pro 215	Thr	Ala	Pro	Thr	Pro 220	Asn	Val	Pro	Ala
Gly	Gly	Asn	Lys	Glu	Thr	Leu	Phe	Asp	Phe	Азр	Phe	Thr	Asn	Leu	Gln

225	MB1:	15 Sequence 1 235	Listing.ST25	240
Ile Pro Asp Phe Gly 245	Phe Leu Ala	Glu Glu Gln 250	Gln Asp Leu As 25	
Asp Cys Phe Leu Ala 260	Asp Asp Gln	Phe Asp Asp 265	Phe Gly Leu Le 270	u Asp
Asp Ile Gln Gly Phe 275	Glu Asp Asn 280	Gly Pro Ser	Ala Leu Pro As 285	p Phe
Asp Phe Ala Asp Val 290	Glu Asp Leu 295	Gln Leu Ala	Asp Ser Ser Ph	e Gly
Phe Leu Asp Gln Leu 305	Ala Pro Ile 310	Asn Ile Ser 315	Cys Pro Leu Ly	s Ser 320
Phe Ala Ala Ser				
<210> 13 <211> 920 <212> DNA <213> Arabidopsis t	haliana			
<220> <221> CDS <222> (1)(852) <223> G237				
<400> 13 atg gcg aag acg aaa	tat gga gag	aga cat agg		40
Met Ala Lys Thr Lys 1 5				
	Tyr Gly Glu	Arg His Arg 10 agc ttc atc	Lys Gly Leu Tr 15 ctc tct tat gg	Ser c cat 96
cct gaa gaa gac gag Pro Glu Glu Asp Glu	Tyr Gly Glu aag cta agg Lys Leu Arg	agc ttc atc Ser Phe Ile 25	Lys Gly Leu Tr 15 ctc tct tat gg Leu Ser Tyr Gl 30 tta caa agg aa	p Ser c cat 96 y His t ggg 144
cct gaa gaa gac gag Pro Glu Glu Asp Glu 20 tct tgc tgg acc act Ser Cys Trp Thr Thr	Tyr Gly Glu aag cta agg Lys Leu Arg gtt ccc atc Val Pro Ile 40 aga tgg att	Arg His Arg 10 agc ttc atc Ser Phe Ile 25 aaaa gct ggg Lys Ala Gly aat tac cta	Lys Gly Leu Tr 15 ctc tct tat gg Leu Ser Tyr Gl 30 tta caa agg aa Leu Gln Arg Ass 45	c cat 96 y His 96 t ggg 144 n Gly
cct gaa gaa gac gag Pro Glu Glu Asp Glu 20 tct tgc tgg acc act Ser Cys Tro Thr Thr 35 aag agc tgc aga tta Lys Ser Cys Arg Leu 50 agg gat atg att agt Arg Asp Met Ile Ser	Tyr Gly Glu aag cta agg Lys Leu Arg gtt ccc atc Val Pro Ile 40 aga tgg att Arg Trp Ile 55 gca gaa gaa	Arg His Arg 10 agc ttc atc Ser Phe Ile 25 aaaa gct ggg Lys Ala Gly aat tac cta Asn Tyr Leu gaa gag act	Lys Gly Leu Tr 15 ctc tct tat gg Leu Ser Tyr Gl 30 tta caa agg aa Leu Gln Arg As 45 aga cca ggg tt Arg Pro Gly Let 60 atc ttg acg tt	c cat 96 y His 96 t ggg 144 n Gly 192 u Lys 192
cct gaa gaa gac gag Pro Glu Glu Asp Glu 20 tct tgc tgg acc act Ser Cys Tro Thr Thr 35 aag agc tgc aga tta Lys Ser Cys Arg Leu 50 agg gat atg att agt Arg Asp Met Ile Ser	Tyr Gly Glu aag cta agg Lys Leu Arg gtt ccc atc Val Pro Ile 40 aga tgg att Arg Trp Ile 55 gca gaa gaa Ala Glu Glu 70 aag tgg tcg	Arg His Arg 10 agc ttc atc Ser Phe Ile 25 aaa gct ggg Lys Ala Gly aat tac cta Asn Tyr Leu gaa gag act Glu Glu Thr 75 caa ata gct	Lys Gly Leu Try 15 ctc tct tat gg Leu Ser Tyr Gly 30 tta caa agg aa Leu Gln Arg As: 45 aga cca ggg tt: Arg Pro Gly Lei 60 atc ttg acg tt: Ile Leu Thr Pho	c cat 96 y His 96 t ggg 144 n Gly 192 u Lys 240 e His 80 g gga 288
cct gaa gaa gac gag Pro Glu Glu Asp Glu 20 tct tgc tgg acc act Ser Cys Trp Thr Thr 35 aag agc tgc aga tta Lys Ser Cys Arg Leu 50 agg gat atg att agt Arg Asp Met Ile Ser 65 tct ccc ttg ggt aac Ser Pro Leu Gly Asn	Tyr Gly Glu aag cta agg Lys Leu Arg gtt ccc atc Val Pro Ile 40 aga tgg att Trp Ile 55 gca gaa gaa Ala Glu Glu 70 aag tgg tcg Lys Trp Ser ata aag aac	Arg His Arg 10 agc ttc atc Ser Phe Ile 25 aaa gct ggg Lys Ala Gly aat tac cta Asn Tyr Leu gaa gag act Glu Glu Thr 75 caa ata gct Gln Ile Ala 90 tat tgg cac	Lys Gly Leu Try 15 ctc tct tat gg Leu Ser Tyr Gl 30 tta caa agg aa Leu Gln Arg As: 45 aga cca ggg tta Arg Pro Gly Lei 60 atc ttg acg tt Ile Leu Thr Phi aaa ttc tta cc Lys Phe Leu Pr 95 tct cat ttg aca	c cat 96 y His 96 t ggg 144 n Gly 192 t cat 240 e His 80 g gga 288 c Gly 288
cct gaa gaa gac gag Pro Glu Glu Asp Glu 20 tct tgc tgg acc act Ser Cys Trp Thr Thr 35 aag agc tgc aga tta Lys Ser Cys Arg Leu 50 agg gat atg att agt Arg Asp Met Ile Ser 65 tct ccc ttg ggt aac Ser Pro Leu Gly Asn 85 aga aca gac aat gag Arg Thr Asp Asn Glu	Tyr Gly Glu aag cta agg Lys Leu Arg gtt ccc atc Val Pro Ile 40 aga tgg att Arg Trp Ile 55 gca gaa gaa Ala Glu Glu 70 aag tgg tcg Lys Trp Ser ata aag aac Ile Lys Asn cag agc tta	Arg His Arg 10 agc ttc atc Ser Phe Ile 25 aaa gct ggg Lys Ala Gly aat tac cta Asn Tyr Leu gaa gag act Glu Glu Thr 75 caa ata gct Gln Ile Ala 90 tat tgg cac Tyr Trp His 105 caa gat gca	Lys Gly Leu Try 15 ctc tct tat gg Leu Ser Tyr Gl 30 tta caa agg aa Leu Gln Arg As: 45 aga cca ggg tta Arg Pro Gly Lei 60 atc ttg acg tti Ile Leu Thr Pho aaa ttc tta ccc Lys Phe Leu Pro 95 tct cat ttg acg Ser His Leu Lyi 110 aaa tct att tca	c cat 96 y His 96 t ggg 144 n Gly 142 a aag 192 Lys 240 e His 80 g gga 288 c Gly 336 e Lys 336 c cct 384

gaa	acc	ttg	atc	tcg	aat	cac						ing.: aga		cta	gag	480
Glu 145	Thr	Leu	Ile	Ser	Asn 150	His	Val	Phe	Ser	Leu 155	Gln	Arg	Leu	Leu	Glu 160	
		tat.	t 02	tat		tos	~~=	733	200	220	002	22 t	220	anc	cat	528
Asn	Lys	Ser	Ser	Ser	ccc Pro	Ser	Gln	Glu	Ser	Asn	Gly	Asn	Asn	Ser	His	520
				165					170					175		
caa	tgt	tct	tct	gct	cct Pro	gag	att	cca	agg Arg	Ctt	ttc	ttc	tct	gaa Glu	tgg.	576
GIII	Cys	261	180	AIG	710	GIU	110	185	9	~~~		• • • • • • • • • • • • • • • • • • • •	190			
ctt	tct	tct	tca	tat	ccc	cac	acc	gat	tat	tcc	tct	gag	ttt	acc	gac	624
Leu	Ser	Ser 195	Ser	Tyr	Pro	His	Thr 200	Asp	Tyr	Ser	Ser	Glu 205	Phe	Thr	Asp	
								~+~		~~~	205		+ 0 =	act	+=+	672
Ser	Lys	His	Ser	Gln	gct Ala	Pro	Asn	Val	Glu	Glu	Thr	Leu	Ser	Ala	Tyr	0,72
	210					215					220					
gaa	gaa	atg	ggt	gat	gtt Val	gat	cag	ttc Phe	cat	tac Tvr	aac Asn	gaa Glu	atg Met	atg Met	atc Ile	720
225	Giu	Mec	Ory	rop	230	nap	U1			235					240	
aac	aac	agc	aac	tgg	act	ctt	aac	gac	att	gtg	ttt	ggt	tcc	aaa	tgt	768
Asn	Asn	Ser	Asn	Trp 245	Thr	Leu	Asn	Asp	Ile 250	Val	Phe	Gly	Ser	Lys 255	Cys	
										~~+	• • •	~~+	tat	22+	fat	816
aag Lys	aag Lys	Gln	gag Glu	Cat His	cat His	Ile	Tyr	Arg	Glu	Ala	Ser	Asp	Сув	Asn	Ser	010
_			260					265					270			
					tct Ser						taa	att	gcgti	tta		862
Ser	AIA	275	FILE	FIIC	Ser	FIO	280	1111	1114	****						
ttgi	taat	gta a	aatca	aaati	t c	taagg	gcaa	a ac	eggaa	aaaa	aaa	aaaa	aaa a	aaaa	aaaa	920
ttgi	taat	gta a	aatca	aati	t c	taagg	gcaa	a ac	cggaa	aaaa	aaa	aaaa	aaa a	aaaa	aaaa	920
<21	0> :	14	aatca	aati	ct c	taag	gcaa	a ac	cggaa	aaaa	aaa	aaaa	aaa a	aaaa	aaaa	920
<210 <210 <210	0> : 1> : 2> :	14 283 PRT					gcaa	a ac	eggaa	aaaa	aaa	aaaa	aaa a	aaaa	aaaa	920
<210 <21	0> : 1> : 2> :	14 283 PRT			t ct		gcaa	a ac	cggaa	aaaa	aaa	aaaa	aaa i	aaaa	aaaa	920
<210 <210 <210 <210 <400	0> : 1> : 2> : 3> :	14 283 PRT Arab:	idop:	sis (thal:	iana										920
<210 <211 <211 <211 <400 Met	0> : 1> : 2> : 3> :	14 283 PRT Arab:	idop:	sis (iana			His							920
<210 <210 <210 <210 <400	0> : 1> : 2> : 3> :	14 283 PRT Arab:	idop:	sis (thal:	iana								Trp		920
<210 <211 <211 <211 <400 Met	0> : 1> : 2> : 3> : 0> :	14 283 PRT Arab: 14	idop: Thr	sis (Lys 5	thal:	iana Gly	Glu	Arg	His 10	Arg	Ьув	Gly	Leu	Trp 15	Ser	920
<210 <211 <211 <211 <400 Met	0> : 1> : 2> : 3> : 0> :	14 283 PRT Arab: 14	idop: Thr	sis (Lys 5	thal: Tyr	iana Gly	Glu	Arg	His 10	Arg	Ьув	Gly	Leu	Trp 15	Ser	920
<210 <211 <211 <211 <400 Met 1	0> : 1> : 2> : 3> : 3> : Ala Glu	14 283 PRT Arab: 14 Lys Glu	idops Thr Asp 20	Lys 5 Glu	thal: Tyr Lys	iana Gly Leu	Glu	Arg Ser 25	His 10 Phe	Arg	Lys	Gly	Leu Tyr 30	Trp 15	Ser His	920
<210 <211 <211 <211 <400 Met 1	0> : 1> : 2> : 3> : 3> : Ala Glu	14 283 PRT Arab: 14 Lys Glu	idops Thr Asp 20	Lys 5 Glu	thal: Tyr	iana Gly Leu	Glu	Arg Ser 25	His 10 Phe	Arg	Lys	Gly	Leu Tyr 30	Trp 15	Ser His	920
<210 <211 <211 <211 <400 Met 1	0> : 1> : 2> : 3> : 0> : Cys	14 283 PRT Arab 14 Lys Glu Trp 35	idops Thr Asp 20	Lys 5 Glu Thr	Tyr Lys Val	iana Gly Leu Pro	Glu Arg	Arg Ser 25	His 10 Phe	Arg Ile Gly	Lys Leu Leu	Gly Ser Gln 45	Leu Tyr 30	Trp 15 Gly Asn	Ser His	920
<210 <211 <211 <211 <400 Met 1	0)> :: 1> :: 2> : 1> :: Cys	14 283 PRT Arab 14 Lys Glu Trp 35	idops Thr Asp 20	Lys 5 Glu Thr	Tyr Lys Val	Gly Leu Pro	Glu Arg	Arg Ser 25	His 10 Phe	Arg Ile Gly	Lys Leu Leu	Gly Ser Gln 45	Leu Tyr 30	Trp 15 Gly Asn	Ser His	920
<210 <211 <211 <211 <400 Met 1	0> : 1> : 2> : 3> : 0> : Cys	14 283 PRT Arab 14 Lys Glu Trp 35	idops Thr Asp 20	Lys 5 Glu Thr	Tyr Lys Val	iana Gly Leu Pro	Glu Arg	Arg Ser 25	His 10 Phe	Arg Ile Gly	Lys Leu Leu	Gly Ser Gln 45	Leu Tyr 30	Trp 15 Gly Asn	Ser His	920
<211 <211 <211 <400 Met 1 Pro	0)> :: 11> :: 22> :: 33> :: Ala Glu Cys Ser 50	14 283 PRT Arab: 14 Lys Glu Trp 35	Thr Asp 20 Thr	Lys 5 Glu Thr	Tyr Lys Val	Gly Leu Pro	Glu Arg Ile 40	Arg Ser 25 Lys	Hiss 10 Phe Ala	Arg Ile Gly Leu	Lys Leu Leu Arg	Gly Ser Gln 45	Leu Tyr 30 Arg	Trp 15 Gly Asn	Ser His Gly Lys	920
<211 <211 <211 <400 Met 1 Pro	0)> :: 11> :: 22> :: 33> :: Ala Glu Cys Ser 50	14 283 PRT Arab: 14 Lys Glu Trp 35	Thr Asp 20 Thr	Lys 5 Glu Thr	Tyr Lys Val	Gly Leu Pro	Glu Arg Ile 40	Arg Ser 25 Lys	Hiss 10 Phe Ala	Arg Ile Gly Leu	Lys Leu Leu Arg	Gly Ser Gln 45	Leu Tyr 30 Arg	Trp 15 Gly Asn	Ser His Gly Lys	920
<211 <211 <211 <211 <211 <211 <211 <211	0)> :: 1> :: 2> :: 2> :: 33> :: Ala Glu Cys Ser 50 Asp	14 283 PRT Arab: 14 Lys Glu Trp 35 Cys	Thr Asp 20 Thr Arg	Lys 5 Glu Thr	Tyr Lys Val Arg	Gly Leu Pro Trp 55	Glu Arg Ile 40 Glu	Arg Ser 25 Lys Asn	His 10 Phe Ala Tyr	Arg Ile Gly Leu Thr 75	Lys Leu Arg 60	Gly Ser Gln 45 Pro	Tyr 30 Arg Gly	Trp 15 Gly Asn Leu	Ser His Gly Lys His	920
<211 <211 <211 <211 <211 <211 <211 <211	0)> :: 1> :: 2> :: 2> :: 33> :: Ala Glu Cys Ser 50 Asp	14 283 PRT Arab: 14 Lys Glu Trp 35 Cys	Thr Asp 20 Thr Arg	Lys 5 Glu Thr	Tyr Lys Val Arg	Gly Leu Pro Trp 55	Glu Arg Ile 40 Glu	Arg Ser 25 Lys Asn	His 10 Phe Ala Tyr	Arg Ile Gly Leu Thr 75	Lys Leu Arg 60	Gly Ser Gln 45 Pro	Tyr 30 Arg Gly	Trp 15 Gly Asn Leu	Ser His Gly Lys	920
<211 <211 <211 <211 <211 <211 <211 <400 Met 1 Pro Ser Ly8 Arg 65	0)> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> ::	14 283 PRT Arab Arab 14 Lys Glu Trp 35 Cys Met	Thr Asp 20 Thr Arg	Lys 5 Glu Thr Leu Ser Asn 85	Tyr Lys Val Arg Ala 70 Lys	Gly Leu Pro Trp 55 Glu Trp	Glu Arg Ile 40 Glu Ser	Arg Ser 25 Lys Asn Glu	His 10 Phe Ala Tyr Glu	Arg Ile Gly Leu Thr 75	Leu Leu Arg 60	Gly Ser Gln 45 Pro	Tyr 30 Arg Gly	Trp 15 Gly Asn Leu Phe	Ser His Gly Lys His 80	920
<211 <211 <211 <211 <211 <211 <211 <400 Met 1 Pro Ser Ly8 Arg 65	0)> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> :: 1> :: 2> ::	14 283 PRT Arab Arab 14 Lys Glu Trp 35 Cys Met	Thr Asp 20 Thr Arg	Lys 5 Glu Thr Leu Ser Asn 85	Tyr Lys Val Arg Ala 70 Lys	Gly Leu Pro Trp 55 Glu Trp	Glu Arg Ile 40 Glu Ser	Arg Ser 25 Lys Asn Glu	His 10 Phe Ala Tyr Glu Ile 90	Arg Ile Gly Leu Thr 75	Leu Leu Arg 60	Gly Ser Gln 45 Pro	Tyr 30 Arg Gly	Trp 15 Gly Asn Leu Phe Pro 95	Ser His Gly Lys His	920

Lys Trp Leu Lys Ser Gln Ser Leu Gln Asp Ala Lys Ser Ile Ser Pro

MBI15 Sequence Listing.ST25 120 125

Pro Ser Ser Ser Ser Ser Leu Val Ala Cys Gly Glu Arg Asn Pro 130 135 140

Glu Thr Leu Ile Ser Asn His Val Phe Ser Leu Gln Arg Leu Leu Glu 145 150 155 160

Asn Lys Ser Ser Ser Pro Ser Gln Glu Ser Asn Gly Asn Asn Ser His 165 170 175

Gln Cys Ser Ser Ala Pro Glu Ile Pro Arg Leu Phe Phe Ser Glu Trp 180 185 190

Leu Ser Ser Ser Tyr Pro His Thr Asp Tyr Ser Ser Glu Phe Thr Asp 195 200 205

Ser Lys His Ser Gln Ala Pro Asn Val Glu Glu Thr Leu Ser Ala Tyr 210 215 220

Glu Glu Met Gly Asp Val Asp Gln Phe His Tyr Asn Glu Met Met Ile 225 230 235 240

Asn Asn Ser Asn Trp Thr Leu Asn Asp Ile Val Phe Gly Ser Lys Cys 245 250 255

Lys Lys Gln Glu His His Ile Tyr Arg Glu Ala Ser Asp Cys Asn Ser 260 265 270

Ser Ala Glu Phe Phe Ser Pro Pro Thr Thr Thr 275 280

<210> 15

<211> 1302

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (331)..(1149)

115

<223> G409

<400> 15
tgtctctctc tctggctctc tttctcttaa cgtgatcata acgtgattcg aaaattggat 60
atagataggt ttcttgttgg atcttgatcc ctctggaaaa ggaggggaga atagcagttc 120
atgatgggat tttgtatctg cccgttggag tcacctgcga gattactatg gagtacaagc 180
ttcttccgcc ataagatcat gatcttctaa tccttcctac ttcttcccat ctttttaatc 240
atcttctcgc tatctctgct tcctctttct ctctgtttcc tctttctcag aactcagaag 300
tagttgttgt tttattctg ttgatcaaaa atg gaa tcc aat tcg ttt ttc ttc Met Glu Ser Asn Ser Phe Phe Phe 1

gat cca tct gct tca cac ggc aac agc atg ttc ttc ctt ggg aat ctc Asp Pro Ser Ala Ser His Gly Asn Ser Met Phe Phe Leu Gly Asn Leu 10 15 20

aat ccc gtc gtc caa gga gga gga gca aga tcg atg atg aac atg gag
Asn Pro Val Val Gln Gly Gly Ala Arg Ser Met Met Asn Met Glu

Page 19

402

25	MBI1	5 Sequence Lis	sting.ST25	40
gaa act tcg aag cga Glu Thr Ser Lys Arg 45	agg ccc ttc Arg Pro Phe	ttt agc tcc co Phe Ser Ser Pr 50	ct gag gat ctc ro Glu Asp Leu 55	tac 498 Tyr
gac gat gac ttt tac Asp Asp Asp Phe Tyr 60	Asp Asp Gln	ttg cct gaa aa Leu Pro Glu Ly 65	ag aag cgt cgc ys Lys Arg Arg 70	ctc 546 Leu
act acc gaa caa gtg Thr Thr Glu Gln Val 75	cat ctg ctg His Leu Leu 80	gag aaa agc tt Glu Lys Ser Pi	tc gag aca gag ne Glu Thr Glu 85	aac 594 Asn
aag cta gag cct gaa Lys Leu Glu Pro Glu 90	cgc aag act Arg Lys Thr 95	Gln Leu Ala Ly	ag aag ctt ggt ys Lys Leu Gly 00	cta 642 Leu
cag cca agg caa gtg Gln Pro Arg Gln Val 105	gct gtc tgg Ala Val Trp 110	ttt cag aat co Phe Gln Asn Ai 115	gc cga gct cgt rg Arg Ala Arg	tgg 690 Trp 120
aaa aca aaa cag ctt Lys Thr Lys Gln Leu 125	gag aga gac Glu Arg Asp	tac gat ctt ct Tyr Asp Leu Le 130	tc aag tcc act eu Lys Ser Thr 135	tac 738 Tyr
gac caa ctt ctt tct Asp Gln Leu Leu Ser 140	aac tac gac Asn Tyr Asp	tcc atc gtc at Ser Ile Val Me 145	tg gac aac gat et Asp Asn Asp 150	aag 786 Lys
ctc aga tcc gag gtt Leu Arg Ser Glu Val 155	act tcc ctg Thr Ser Leu 160	acc gaa aag ct Thr Glu Lys Le	tt cag ggc aaa eu Gln Gly Lys 165	caa 834 Gln
gag aca gct aat gaa Glu Thr Ala Asn Glu 170	cca cct ggt Pro Pro Gly 175	Gln Val Pro G	aa cca aac caa lu Pro Asn Gln 80	ctt 882 Leu
gat ccg gtt tac att Asp Pro Val Tyr Ile 185	aat gcg gca Asn Ala Ala 190	gca atc aaa ad Ala Ile Lys Th 195	cc gag gac cgg hr Glu Asp Arg	tta 930 Leu 200
agt tca ggg agc gtt Ser Ser Gly Ser Val 205	ggg agc gcg Gly Ser Ala	gta cta gac ga Val Leu Asp As 210	ac gac gca cct sp Asp Ala Pro 215	caa 978 Gln
cta cta gac agc tgt Leu Leu Asp Ser Cys 220	gac tct tac Asp Ser Tyr	ttc cca agc at Phe Pro Ser II 225	tc gta ccc atc le Val Pro Ile 230	caa 1026 Gln
gac aac agc aac gcc Asp Asn Ser Asn Ala 235	agt gat cat Ser Asp His 240	gac aat gac co Asp Asn Asp A	gg agc tgt ttc rg Ser Cys Phe 245	gcc 1074 Ala
gac gtc ttt gtg ccc Asp Val Phe Val Pro 250	acc act tca Thr Thr Ser 255	Pro Ser His A	at cat cac ggt sp His His Gly 60	gaa 1122 Glu
tca ttg gct ttc tgg Ser Leu Ala Phe Trp 265	gga tgg cct Gly Trp Pro 270	tag aaaaccacto	c tgataataaa	1169
tgtgtgttta tttaagtt				
tgactcgtgg aataattc	cg ctgttcaacg	gtattttat c	agttgcatt atat	
atgaaaaaaa aaa				1302
<pre><210> 16 <211> 272 <212> PRT <213> Arabidopsis</pre>	thaliana			

<400> 16

MBI15 Sequence Listing.ST25

Met Glu Ser Asn Ser Phe Phe Phe Asp Pro Ser Ala Ser His Gly Asn 1 10 15

Ser Met Phe Phe Leu Gly Asn Leu Asn Pro Val Val Gln Gly Gly Gly 20 25 30

Ala Arg Ser Met Met Asn Met Glu Glu Thr Ser Lys Arg Arg Pro Phe 35 40

Phe Ser Ser Pro Glu Asp Leu Tyr Asp Asp Asp Phe Tyr Asp Asp Gln 50 55

Leu Pro Glu Lys Lys Arg Arg Leu Thr Thr Glu Gln Val His Leu Leu 65 70 75 80

Glu Lys Ser Phe Glu Thr Glu Asn Lys Leu Glu Pro Glu Arg Lys Thr 85 90 95

Gln Leu Ala Lys Lys Leu Gly Leu Gln Pro Arg Gln Val Ala Val Trp 100 105 110

Phe Gln Asn Arg Arg Ala Arg Trp Lys Thr Lys Gln Leu Glu Arg Asp 115 120 125

Tyr Asp Leu Leu Lys Ser Thr Tyr Asp Gln Leu Leu Ser Asn Tyr Asp 130 135 140

Ser Ile Val Met Asp Asn Asp Lys Leu Arg Ser Glu Val Thr Ser Leu 145 150 155 160

Thr Glu Lys Leu Gln Gly Lys Gln Glu Thr Ala Asn Glu Pro Pro Gly 165 170 175

Gln Val Pro Glu Pro Asn Gln Leu Asp Pro Val Tyr Ile Asn Ala Ala 180 185 190

Ala Ile Lys Thr Glu Asp Arg Leu Ser Ser Gly Ser Val Gly Ser Ala 195 200 205

Val Leu Asp Asp Asp Ala Pro Gln Leu Leu Asp Ser Cys Asp Ser Tyr 210 215 220

Phe Pro Ser Ile Val Pro Ile Gln Asp Asn Ser Asn Ala Ser Asp His 225 230 235 240

Asp Asn Asp Arg Ser Cys Phe Ala Asp Val Phe Val Pro Thr Thr Ser 245 250 255

Pro Ser His Asp His His Gly Glu Ser Leu Ala Phe Trp Gly Trp Pro

<210> 17

<211> 2545 <212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS <222> (103)..(2322) <223> G418

<400> 17 acatatgttt taaattettt gtetgaatet tacaggatee gagagagaga getetggaae 60														60		
gata	_										at a	tg g let G	ga a	ta a	ct	114
aaa Lys 5	act Thr	tct Ser	cct Pro	aat Asn	act Thr 10	aca Thr	att Ile	ctc Leu	ttg Leu	aag Lys 15	act Thr	ttt Phe	cac His	aat Asn	aat Asn 20	162
tct Ser	atg Met	tcc Ser	caa Gln	gat Asp 25	tat Tyr	cat His	cat His	cat His	cat His 30	cat His	cat His	aat Asn	caa Gln	cac His 35	caa Gln	210
gga Gly	ggt Gly	atc Ile	ttc Phe 40	aac Asn	ttc Phe	tct Ser	aat Asn	gga Gly 45	ttc Phe	gac Asp	cga Arg	tca Ser	gat Asp 50	tct Ser	CCC Pro	258
aat Asn	tta Leu	aca Thr 55	act Thr	cag Gln	cag Gln	aag Lys	caa Gln 60	gag Glu	cat His	caa Gln	agg Arg	gta Val 65	gag Glu	atg Met	gac Asp	306
gag Glu	gaa Glu 70	tct Ser	tca Ser	gtc Val	gcc Ala	gga Gly 75	ggt Gly	agg Arg	att Ile	ccg Pro	gtc Val 80	tac Tyr	gaa Glu	tca Ser	gcc Ala	354
ggt Gly 85	atg Met	tta Leu	tcc Ser	gaa Glu	atg Met 90	ttt Phe	aat Asn	ttc Phe	ccc Pro	gga Gly 95	agc Ser	agc Ser	ggt Gly	gga Gly	gga Gly 100	402
aga Arg	gat Asp	ctc Leu	gac Asp	ctc Leu 105	ggc Gly	caa Gln	tct Ser	ttc Phe	cgg Arg 110	tca Ser	aat Asn	agg Arg	cag Gln	ttg Leu 115	ctt Leu	450
gag Glu	gag Glu	caa Gln	cat His 120	cag Gln	aat Asn	att Ile	ccg Pro	gct Ala 125	atg Met	aat Asn	gct Ala	acg Thr	gat Asp 130.	tca Ser	gcc Ala	498
acc Thr	gcc Ala	acc Thr 135	gca Ala	gcc Ala	gcc Ala	atg Met	cag Gln 140	tta Leu	ttc Phe	ttg Leu	atg Met	aat Asn 145	cca Pro	ccg Pro	cca Pro	546
ccg Pro	caa Gln 150	caa Gln	cca Pro	ccg Pro	tct Ser	ccg Pro 155	tca Ser	tcc Ser	aca Thr	act Thr	tcc Ser 160	cca Pro	agg Arg	agc Ser	cac His	594
His 165	Asn	Ser	tca Ser	Thr	Leu 170	His	Met	Leu	Leu	Pro 175	Ser	Pro	ser	Tnr	180	642
Thr	Thr	His	cat His	Gln 185	Asn	Tyr	Thr	Asn	His 190	Met	Ser	Met	His	195	Leu	690
Pro	His	Gln	cat His 200	His	Gln	Gln	Ile	Ser 205	Thr	Trp	Gln	Ser	Ser 210	Pro	Авр	738
His	His	His 215		His	His	Asn	Ser 220	Gln	Thr	Glu	Ile	Gly 225	Thr	Val	нів	786
Val	Glu 230	Asn	agc Ser	Gly	Gly	His 235	Gly	Gly	Gln	Gly	Leu 240	Ser	Leu	ser	Leu	834
tca Ser 245	Ser	tct Ser	tta Leu	gag Glu	gct Ala 250	Ala	gca Ala	aaa Lys	Ala	gaa Glu 255	Glu	tat Tyr	aga Arg	aac Asn	att Ile 260	882

MBI15 Sequence Listing.ST25

							WBI	15 5	eque	псе	List	ing.	ST25			
						tct Ser										930
caa Gln	ttc Phe	aag Lys	act Thr 280	ctt Leu	ctt Leu	gct Ala	aat Asn	tct Ser 285	tct Ser	caa Gln	cat His	cac His	cat His 290	caa Gln	gta Val	978
tta Leu	aac Asn	caa Gln 295	ttc Phe	cga Arg	tca Ser	tct Ser	ccg Pro 300	gct Ala	gct Ala	tct Ser	tcc Ser	tct Ser 305	tcc Ser	atg Met	gca Ala	1026
						aac Asn 315	Ser									1074
ttg Leu 325	ttg Leu	gaa Glu	gag Glu	ttt Phe	tgt Cys 330	agt Ser	gtt Val	gga Gly	aga Arg	gga Gly 335	ttt Phe	ttg Leu	aag Lys	aag Lys	aac Asn 340	1122
						aac Asn										1170
						gcc Ala										1218
						gag Glu										1266
						gac Asp 395										1314
						tct Ser										1362
			Tyr			ttg Leu		Gln								1410
						gtt Val										1458
ctt Leu	ctt Leu	999 Gly 455	gac Asp	aaa Lys	gat Asp	gca Ala	gcg Ala 460	gga Gly	atc Ile	tct Ser	tct Ser	tcc Ser 465	G1y 999	tta Leu	aca Thr	1506
aaa Lys	ggt Gly 470	gaa Glu	act Thr	ccg Pro	cgt Arg	ttg Leu 475	cgt Arg	ttg Leu	cta Leu	gag Glu	caa Gln 480	agt Ser	ttg Leu	cgt Arg	cag Gln	1554
caa Gln 485	cgt Arg	gcg Ala	ttt Phe	cat His	caa Gln 490	atg Met	ggt Gly	atg Met	atg Met	gaa Glu 495	caa Gln	gaa Glu	gct Ala	tgg Trp	cgg Arg 500	1602
cca Pro	caa Gln	cgc Arg	ggt Gly	ttg Leu 505	cct Pro	gaa Glu	cgc Arg	tcc Ser	gtc Val 510	aat Asn	ata Ile	ctt Leu	aga Arg	gct Ala 515	tgg Trp	1650
						cac His										1698
						ggt Gly										1746
						cgt Arg			Lys		Met					1794

	550					555	MBI	15 S	eque	nce 1	List: 560	ing.	ST25			
			gaa Glu													1842
aac Asn	gaa Glu	gaa Glu	gat Asp	caa Gln 585	gaa Glu	aca Thr	aaa Lys	aac Asn	agc Ser 590	aac Asn	gac Asp	gac Asp	aag Lys	agc Ser 595	aca Thr	1890
aaa Lys	tcc Ser	aac Asn	aac Asn 600	aat Asn	gaa Glu	agc Ser	aac Asn	ttc Phe 605	act Thr	gcc Ala	gtt Val	cgg Arg	acc Thr 610	act Thr	tca Ser	1938
caa Gln	act Thr	cca Pro 615	acg Thr	aca Thr	acc Thr	gca Ala	cca Pro 620	gac Asp	gca Ala	tca Ser	gac Asp	gca Ala 625	gac Asp	gca Ala	gca Ala	1986
gta Val	gcg Ala 630	aca Thr	ggc Gly	cac His	cgt Arg	cta Leu 635	aga Arg	tcc Ser	aac Asn	att Ile	aat Asn 640	gct Ala	tac Tyr	gaa Glu	aac Asn	2034
gac Asp 645	gct Ala	tca Ser	tca Ser	ctt Leu	cta Leu 650	ctc Leu	cct Pro	tcc Ser	tct Ser	tat Tyr 655	tcc Ser	aac Asn	gcc Ala	gcc Ala	gct Ala 660	2082
cct Pro	gcc Ala	gct Ala	gtt Val	tct Ser 665	gac Asp	gac Asp	ttg Leu	aat Asn	tct Ser 670	cgt Arg	tac Tyr	ggt Gly	ggc Gly	tca Ser 675	gac Asp	2130
gcg Ala	ttt Phe	tcc Ser	gcc Ala 680	gtt Val	gcc Ala	acg Thr	tgt Cys	caa Gln 685	caa Gln	agt Ser	gta Val	ggt Gly	999 Gly 690	ttc Phe	gat Asp	2178
gat Asp	gct Ala	gac Asp 695	atg Met	gat Asp	ggt Gly	gtt Val	aac Asn 700	gtt Val	ata Ile	agg Arg	ttt Phe	999 Gly 705	aca Thr	aac Asn	cct Pro	2226
act Thr	ggt Gly 710	gac Asp	gtg Val	tct Ser	ctc Leu	acg Thr 715	ctt Leu	ggt Gly	tta Leu	cgc Arg	cac His 720	gct Ala	gga Gly	aac Asn	atg Met	2274
cct Pro 725	gac Asp	aaa Lys	gac Asp	gct Ala	tct Ser 730	ttc Phe	tgc Cys	gtt Val	aga Arg	gag Glu 735	ttt Phe	999 999	ggt Gly	ttt Phe	tag	2322
ttt	gctt	ttg	tcac	tcca	tt ta	aatt	aatt	a at	tata	gttt	tcc	attc	tta	ctta	ttttaa	2382
ttg	aaaa	tct :	attt	ttgt	ct c	ttaa	aagt	c ca	aaca	atac	att	agtc	tag	ccct	cctctg	2442
Ctt	tttt	ttt	tcta	tctc	gt ga	aaga	gaag	a aa	acga	tacg	taa	atcc	ctt	cgaa	aactaa	2502
tgt	acgt	tgt .	acga	ctta	tt g	tttt	cata	a aa	aaaa	aaaa	aaa					2545
<21 <21		18 739														

<211> 739 <212> PRT <213> Arabidopsis thaliana

Met Gly Ile Thr Lys Thr Ser Pro Asn Thr Thr Ile Leu Leu Lys Thr 1 5 10 15

Phe His Asn Asn Ser Met Ser Gln Asp Tyr His His His His His 20 25 30

Asn Gln His Gln Gly Gly Ile Phe Asn Phe Ser Asn Gly Phe Asp Arg 35 40 45

Ser Asp Ser Pro Asn Leu Thr Thr Gln Gln Lys Gln Glu His Gln Arg Page 24

Val Glu Met Asp Glu Glu Ser Ser Val Ala Gly Gly Arg Ile Pro Val 65 70 75 80

Tyr Glu Ser Ala Gly Met Leu Ser Glu Met Phe Asn Phe Pro Gly Ser 85 90 95

Ser Gly Gly Arg Asp Leu Asp Leu Gly Gln Ser Phe Arg Ser Asn 100 105 110

Arg Gln Leu Leu Glu Glu Gln His Gln Asn Ile Pro Ala Met Asn Ala 115 120 125

Thr Asp Ser Ala Thr Ala Thr Ala Ala Ala Met Gln Leu Phe Leu Met 130 135 140

Asn Pro Pro Pro Pro Gln Gln Pro Pro Ser Pro Ser Ser Thr Thr Ser 145 150 155 160

Pro Arg Ser His His Asn Ser Ser Thr Leu His Met Leu Leu Pro Ser 165 170 175

Pro Ser Thr Asn Thr Thr His His Gln Asn Tyr Thr Asn His Met Ser 180 185 190

Met His Gln Leu Pro His Gln His His Gln Gln Ile Ser Thr Trp Gln 195 200 205

Ser Ser Pro Asp His His His His His His Asn Ser Gln Thr Glu Ile 210 215 220

Gly Thr Val His Val Glu Asn Ser Gly Gly His Gly Gly Gln Gly Leu 225 230 230 235 240

Ser Leu Ser Leu Ser Ser Leu Glu Ala Ala Ala Lys Ala Glu Glu 245 250 255

Tyr Arg Asn Ile Tyr Tyr Gly Ala Asn Ser Ser Asn Ala Ser Pro His 260 265 270

His Gln Tyr Asn Gln Phe Lys Thr Leu Leu Ala Asn Ser Ser Gln His 275 280 285

His His Gln Val Leu Asn Gln Phe Arg Ser Ser Pro Ala Ala Ser Ser 290 295 300

Ser Ser Met Ala Ala Val Asn Ile Leu Arg Asn Ser Arg Tyr Thr Thr 305 310 315 320

Ala Ala Gln Glu Leu Leu Glu Glu Phe Cys Ser Val Gly Arg Gly Phe

Leu Lys Lys Asn Lys Leu Gly Asn Ser Ser Asn Pro Asn Thr Cys Gly 340 345 350

- MBI15 Sequence Listing.ST25
 Gly Asp Gly Gly Gly Ser Ser Pro Ser Ser Ala Gly Ala Asn Lys Glu
 355 360 365
- His Pro Pro Leu Ser Ala Ser Asp Arg Ile Glu His Gln Arg Arg Lys 370 375 380
- Val Lys Leu Leu Thr Met Leu Glu Glu Val Asp Arg Arg Tyr Asn His 385 395 400
- Tyr Cys Glu Gln Met Gln Met Val Val Asn Ser Phe Asp Ile Val Met 405 410 415
- Gly His Gly Ala Ala Leu Pro Tyr Thr Ala Leu Ala Gln Lys Ala Met 420 430
- Ser Arg His Phe Arg Cys Leu Lys Asp Ala Val Ala Ala Gln Leu Lys 435 440 445
- Gln Ser Cys Glu Leu Leu Gly Asp Lys Asp Ala Ala Gly Ile Ser Ser 450 455 460
- Ser Gly Leu Thr Lys Gly Glu Thr Pro Arg Leu Arg Leu Leu Glu Gln 465 470 480
- Ser Leu Arg Gln Gln Arg Ala Phe His Gln Met Gly Met Met Glu Gln 485 490 495
- Glu Ala Trp Arg Pro Gln Arg Gly Leu Pro Glu Arg Ser Val Asn Ile 500 505 510
- Leu Arg Ala Trp Leu Phe Glu His Phe Leu His Pro Tyr Pro Ser Asp 515 520 525
- Ala Asp Lys His Leu Leu Ala Arg Gln Thr Gly Leu Ser Arg Asn Gln 530 535 540
- Val Ser Asn Trp Phe Ile Asn Ala Arg Val Arg Leu Trp Lys Pro Met 545 550 555 560
- Val Glu Glu Met Tyr Gln Gln Glu Ser Lys Glu Arg Glu Arg Glu Glu 565 570 575
- Glu Leu Glu Glu Asn Glu Glu Asp Gln Glu Thr Lys Asn Ser Asn Asp 580 585 590
- Asp Lys Ser Thr Lys Ser Asn Asn Asn Glu Ser Asn Phe Thr Ala Val
- Arg Thr Thr Ser Gln Thr Pro Thr Thr Thr Ala Pro Asp Ala Ser Asp 610 615 620
- Ala Asp Ala Ala Val Ala Thr Gly His Arg Leu Arg Ser Asn Ile Asn 625 630 635
- Ala Tyr Glu Asn Asp Ala Ser Ser Leu Leu Leu Pro Ser Ser Tyr Ser 645 650 655

MBI15 Sequence Listing.ST25

Asn Ala Ala Pro Ala Ala Val Ser Asp Asp Leu Asn Ser Arg Tyr
660 665 670

Gly Gly Ser Asp Ala Phe Ser Ala Val Ala Thr Cys Gln Gln Ser Val 675 680 685

Gly Gly Phe Asp Asp Ala Asp Met Asp Gly Val Asn Val Ile Arg Phe 690 695 700

Gly Thr Asn Pro Thr Gly Asp Val Ser Leu Thr Leu Gly Leu Arg His 705 710 715 720

Ala Gly Asn Met Pro Asp Lys Asp Ala Ser Phe Cys Val Arg Glu Phe 725 730 735

Gly Gly Phe

<210> 19 <211> 1240 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS <222> (88)..(1020) <223> G591

<400> 19 gtaaatctct ctttgaaggt tcctaactcg ttaatcgtaa ctcacagtga ctcgttcgag 60 tcaaagtctc tgtctttagc tcaaacc atg gct agt aac aac cct cac gac aac Met Ala Ser Asn Asn Pro His Asp Asn 114 ctt tct gac caa act cct tct gat gat ttc ttc gag caa atc ctc ggc Leu Ser Asp Gln Thr Pro Ser Asp Asp Phe Phe Glu Gln Ile Leu Gly 162 ctt cct aac ttc tca gcc tct tct gcc gcc ggt tta tct gga gtt gac Leu Pro Asn Phe Ser Ala Ser Ser Ala Ala Gly Leu Ser Gly Val Asp 210 gga gga tta ggt ggt gga gca ccg cct atg atg ctg cag ttg ggt tcc Gly Gly Leu Gly Gly Gly Ala Pro Pro Met Met Leu Gln Leu Gly Ser 45 50 55 258 gga gaa gaa gga agt cac atg ggt ggc tta gga gga agt gga cca act Gly Glu Glu Gly Ser His Met Gly Gly Leu Gly Gly Ser Gly Pro Thr 60 65 70306 ggg ttt cac aat cag atg ttt cct ttg ggg tta agt ctt gat caa ggg Gly Phe His Asn Gln Met Phe Pro Leu Gly Leu Ser Leu Asp Gln Gly 354 80 aaa gga cct ggg ttt ctt aga cct gaa gga gga cat gga agt ggg aaa Lys Gly Pro Gly Phe Leu Arg Pro Glu Gly Gly His Gly Ser Gly Lys 90 95 100 105 402 aga ttc tca gat gat gtt gtt gat aat cga tgt tct tct atg aaa cct Arg Phe Ser Asp Asp Val Val Asp Asn Arg Cys Ser Ser Met Lys Pro 450

gtt ttc cac ggg cag cct atg caa cag cca cct cca tcg gcc cca cat Val Phe His Gly Gln Pro Met Gln Gln Pro Pro Pro Ser Ala Pro His 125 130 135

498

MBI15 Sequence Listing.ST25 cag cct act tca atc cgt ccc agg gtt cga gct agg cgt ggt cag gct Gln Pro Thr Ser Ile Arg Pro Arg Val Arg Ala Arg Arg Gly Gln Ala 140 145 150	546
act gat cca cat agc atc gct gag cgg cta cgt aga gaa aga ata gca Thr Asp Pro His Ser Ile Ala Glu Arg Leu Arg Arg Glu Arg Ile Ala 155 160 165	594
gaa cgg atc agg gcg ctg cag gaa ctt gta cct act gtg aac aag acc Glu Arg Ile Arg Ala Leu Gln Glu Leu Val Pro Thr Val Asn Lys Thr 170 175 180 185	642
gat aga gct gct atg atc gat gag att gtc gat tat gta aag ttt ctc Asp Arg Ala Ala Met Ile Asp Glu Ile Val Asp Tyr Val Lys Phe Leu 190 195 200	690
agg ctc caa gtc aag gtt ttg agc atg aac cga ctt ggt gga gcc ggt Arg Leu Gln Val Lys Val Leu Ser Met Asn Arg Leu Gly Gly Ala Gly 205 210 215	738
gcg gtt gct cca ctt gtt act gat atg cct ctt tca tca tca gtt gag Ala Val Ala Pro Leu Val Thr Asp Met Pro Leu Ser Ser Val Glu 220 225 230	786
gat gaa acg ggt gag ggt gga agg act ccg caa cca gcg tgg gag aaa Asp Glu Thr Gly Glu Gly Gly Arg Thr Pro Gln Pro Ala Trp Glu Lys 235 240 245	834
tgg tct aac gat ggg act gaa cgt caa gtg gct aaa ctg atg gaa gag Trp Ser Asn Asp Gly Thr Glu Arg Gln Val Ala Lys Leu Met Glu Glu 250 255 260 265	882
aac gtt gga gcc gcg atg cag ctt ctt caa tca aag gct ctt tgt atg Asn Val Gly Ala Ala Met Gln Leu Leu Gln Ser Lys Ala Leu Cys Met 270 275 280	930
atg cca atc tca ttg gca atg gca att tac cat tct caa cct ccg gat Met Pro Ile Ser Leu Ala Met Ala Ile Tyr His Ser Gln Pro Pro Asp 285 290 295	978
aca tct tca gtg gtc aag cct gag aac aat cct cca cag tag Thr Ser Ser Val Val Lys Pro Glu Asn Asn Pro Pro Gln 300 305 310	1020
gatttctgca ataaagagtt tgtacagcta atccaactgt ccaacatggg tttttcttct	1080
gctctaatga ctctggtttc ttctctcctc tctcaccgac ttgaaaggta aaaaagtgaa	1140
aaaggctttg tagatggaat caatgtagga tttgcagtag agggcaaaaa aatgtcatat	1200
agctcaattg atcaagtctt aaaaaaaaaaa aaaaaaaaaa	1240
<210> 20 <211> 310 <212> PRT <213> Arabidopsis thaliana	
<400> 20	
Met Ala Ser Asn Asn Pro His Asp Asn Leu Ser Asp Gln Thr Pro Ser 1 10 15	
Asp Asp Phe Phe Glu Gln Ile Leu Gly Leu Pro Asn Phe Ser Ala Ser 20 25 30	

Pro Pro Met Met Leu Gln Leu Gly Ser Gly Glu Glu Gly Ser His Met 50 . 60 Page 28

Ser Ala Ala Gly Leu Ser Gly Val Asp Gly Gly Leu Gly Gly Gly Ala 35 4040

MBI15 Sequence Listing.ST25

Gly Gly Leu Gly Gly Ser Gly Pro Thr Gly Phe His Asn Gln Met Phe

Pro Leu Gly Leu Ser Leu Asp Gln Gly Lys Gly Pro Gly Phe Leu Arg

Pro Glu Gly Gly His Gly Ser Gly Lys Arg Phe Ser Asp Asp Val Val 100 105 110

Asp Asn Arg Cys Ser Ser Met Lys Pro Val Phe His Gly Gln Pro Met 115 120 125

Gln Gln Pro Pro Pro Ser Ala Pro His Gln Pro Thr Ser Ile Arg Pro 130 135 140

Arg Val Arg Ala Arg Gly Gln Ala Thr Asp Pro His Ser Ile Ala 145 150 160

Glu Arg Leu Arg Glu Arg Ile Ala Glu Arg Ile Arg Ala Leu Gln 165 170 175

Glu Leu Val Pro Thr Val Asn Lys Thr Asp Arg Ala Ala Met Ile Asp 180 185 190

Glu Ile Val Asp Tyr Val Lys Phe Leu Arg Leu Gln Val Lys Val Leu

Ser Met Asn Arg Leu Gly Gly Ala Gly Ala Val Ala Pro Leu Val Thr 210 215 220

Asp Met Pro Leu Ser Ser Ser Val Glu Asp Glu Thr Gly Glu Gly Gly

Arg Thr Pro Gln Pro Ala Trp Glu Lys Trp Ser Asn Asp Gly Thr Glu 245 250 255

Arg Gln Val Ala Lys Leu Met Glu Glu Asn Val Gly Ala Ala Met Gln 265 270

Leu Leu Gln Ser Lys Ala Leu Cys Met Met Pro Ile Ser Leu Ala Met 275 280 285

Ala Ile Tyr His Ser Gln Pro Pro Asp Thr Ser Ser Val Val Lys Pro 295

Glu Asn Asn Pro Pro Gln

<210>

<211> 1179 <212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (109)..(966)

G525

<400> 21 cttctctctt ctcaaaaacc cttccctctt cgtctccaaa caacaacaaa cacaacaaca 60														60		
acaaaaatct tacaagaaga tcatttttag aaaccctatt aggataaa atg gat tac $$\tt 11'$ Met Asp Tyr $\tt 1$														117		
gag Glu	gca Ala 5	tca Ser	aga Arg	atc Ile	Val	gaa Glu 10	atg Met	gta Val	gaa Glu	gat Asp	gaa Glu 15	gaa Glu	cat His	ata Ile	gat Asp	165
cta Leu 20	cca Pro	cca Pro	gga Gly	ttc Phe	aga Arg 25	ttt Phe	cac His	cct Pro	act Thr	gat Asp 30	gaa Glu	gaa Glu	ctc Leu	ata Ile	act Thr 35	213
cac His	tac Tyr	ctc Leu	Lys	cca Pro 40	aag Lys	gtt Val	ttc Phe	aac Asn	act Thr 45	ttc Phe	ttc Phe	tct Ser	gct Ala	act Thr 50	gcc Ala	261
att Ile	ggt Gly	gaa Glu	gtt Val 55	gat Asp	ctc Leu	aac Asn	aag Lys	att Ile 60	gag Glu	cct Pro	tgg Trp	gac Asp	tta Leu 65	cca Pro	tgg Trp	309
aag Lys	gct Ala	aag Lys 70	atg Met	gga Gly	gaa Glu	aaa Lys	gaa Glu 75	tgg Trp	tat Tyr	ttc Phe	ttc Phe	tgt Cys 80	gtg Val	aga Arg	gac Asp	357
cgg Arg	aaa Lys 85	tac Tyr	ccg Pro	acc Thr	ggt Gly	tta Leu 90	agg Arg	aca Thr	aac Asn	cgg Arg	gcg Ala 95	aca Thr	gaa Glu	gcc Ala	ggt Gly	405
tat Tyr 100	tgg Trp	aaa Lys	gcc Ala	aca Thr	gga Gly 105	aaa Lys	gac Asp	aaa Lys	gag Glu	ata Ile 110	ttc Phe	aag Lys	gga Gly	aaa Lys	tca Ser 115	453
ctt Leu	gtg Val	ggt Gly	atg Met	aag Lys 120	aaa Lys	act Thr	ttg Leu	gtt Val	ttc Phe 125	tat Tyr	aaa Lys	gga Gly	aga Arg	gct Ala 130	cct Pro	501
aaa Lys	gga Gly	gtt Val	aaa Lys 135	acc Thr	aat Asn	tgg Trp	gtt Val	atg Met 140	cat His	gaa Glu	tat Tyr	cgt Arg	tta Leu 145	gaa Glu	ggc Gly	549
aaa Lys	tat Tyr	tgt Сув 150	att Ile	gaa Glu	aat Asn	ctt Leu	ccc Pro 155	caa Gln	aca Thr	gct Ala	aag Lys	aac Asn 160	gaa Glu	tgg Trp	gtt Val	597
ata Ile	tgt Cys 165	cgt Arg	gtt Val	ttc Phe	caa Gln	aaa Lys 170	cgt Arg	gcc Ala	gat Asp	ggt Gly	aca Thr 175	aag Lys	gtt Val	cca Pro	atg Met	645
tca Ser 180	atg Met	ctt Leu	gat Asp	cca Pro	cac His 185	att Ile	aac Asn	cga Arg	atg Met	gaa Glu 190	cca Pro	gcc Ala	ggt Gly	tta Leu	cct Pro 195	693
tcg Ser	tta Leu	atg Met	gat Asp	tgt Cys 200	tct Ser	caa Gln	cga Arg	gac Asp	tcc Ser 205	ttc Phe	acc Thr	ggt Gly	tcg Ser	tcg Ser 210	tct Ser	741
cac His	gtg Val	acc Thr	tgc Cys 215	ttc Phe	tcc Ser	gac Asp	caa Gln	gaa Glu 220	acc Thr	gaa Glu	gac Asp	aaa Lys	aga Arg 225	ctt Leu	gtc Val	789
cac His	gag Glu	tcc Ser 230	aaa Lys	gac Asp	ggt Gly	ttt Phe	ggt Gly 235	Ser	ctg Leu	ttt Phe	tac Tyr	ser 240	Asp	cct Pro	ctg Leu	837
ttt Phe	tta Leu 245	caa Gln	gac Asp	aat Asn	tat Tyr	tcg Ser 250	Leu	atg Met	aag Lys	ctg Leu	ttg Leu 255	Leu	gac Asp	ggt Gly	caa Gln	885
gaa Glu 260	Thr	caa Gln	ttc Phe	tcc Ser	ggc Gly 265	Lys	cct Pro	ttc Phe	Asp	ggt Gly 270	Arg	gat Asp	tcg Ser	tcc	ggt Gly 275	933

MBI15 Sequence Listing.ST25

	a ttg gat to u Leu Asp Cy 280			gttgtataag	ttatgttgta	986
gacttgtagt	agtcatgtgt	tcgtgtgtgt	gaatgaatat	tcttgttaca	tttttttgta	1046
aaaaaggaga	aaaaaatatg	ctagaaagtc	aattgctttt	gttatgtagc	attagtgttt	1106
tttatgtact	caatagactt	cctaattaaa	taaaaatctt	aatttatttg	ccaaaaaaaa	1166
aaaaaaaaa	aaa					1179

<210> 22

<211> 285

<212> PRT

<213> Arabidopsis thaliana

<400> 22

Met Asp Tyr Glu Ala Ser Arg Ile Val Glu Met Val Glu Asp Glu Glu 1 5 15

His Ile Asp Leu Pro Pro Gly Phe Arg Phe His Pro Thr Asp Glu Glu 20 25 30

Leu Ile Thr His Tyr Leu Lys Pro Lys Val Phe Asn Thr Phe Phe Ser 35 40 45

Ala Thr Ala Ile Gly Glu Val Asp Leu Asn Lys Ile Glu Pro Trp Asp 50 60

Leu Pro Trp Lys Ala Lys Met Gly Glu Lys Glu Trp Tyr Phe Phe Cys 65 70 75 80

Val Arg Asp Arg Lys Tyr Pro Thr Gly Leu Arg Thr Asn Arg Ala Thr 85 90 95

Glu Ala Gly Tyr Trp Lys Ala Thr Gly Lys Asp Lys Glu Ile Phe Lys 100 105 110

Gly Lys Ser Leu Val Gly Met Lys Lys Thr Leu Val Phe Tyr Lys Gly

Arg Ala Pro Lys Gly Val Lys Thr Asn Trp Val Met His Glu Tyr Arg 130 135 140

Leu Glu Gly Lys Tyr Cys Ile Glu Asn Leu Pro Gln Thr Ala Lys Asn 145 150 155 160

Glu Trp Val Ile Cys Arg Val Phe Gln Lys Arg Ala Asp Gly Thr Lys 165 170 175

Val Pro Met Ser Met Leu Asp Pro His Ile Asn Arg Met Glu Pro Ala 180 185 190

Gly Leu Pro Ser Leu Met Asp Cys Ser Gln Arg Asp Ser Phe Thr Gly 195 200 205

Ser Ser Ser His Val Thr Cys Phe Ser Asp Gln Glu Thr Glu Asp Lys 210 215 220

Arg Leu Val 225	. His Glu	Ser Lys 230	Asp Gly	Phe Gly 235	Ser Leu	Phe Tyr	Ser 240
Asp Pro Le	Phe Leu 245		Asn Tyr	Ser Leu 250	Met Lys	Leu Leu 255	Leu
Asp Gly Gl	Glu Thr 260	Gln Phe	Ser Gly 265	Lys Pro	Phe Asp	Gly Arg 270	Asp
Ser Ser Gl 27		Glu Leu	Asp Cys 280	Val Trp	Asn Phe 285		
<210> 23 <211> 890 <212> DNA <213> Ara	oidopsis	thaliana					
<220> <221> CDS <222> (55 <223> G54)(738) 5					·	
<400> 23 gcaaccttca	aactaaaa	act cgaga	gacaa gaa	aatcctca	gaatcttt	aa ctta	atg 57 Met 1
gcg ctc ga Ala Leu Gl	g gct cti u Ala Lei 5	aca tca Thr Ser	cca aga Pro Arg 10	tta gct Leu Ala	tct ccg Ser Pro	att cct Ile Pro 15	cct 105 Pro
ttg ttc ga Leu Phe Gl 20	u Asp Se:	t tca gtc r Ser Val	ttc cat Phe His 25	gga gtc Gly Val	gag cac Glu His 30	tgg aca Trp Thr	aag 153 Lys
ggt aag cg Gly Lys Ar 35	a tct aag g Ser Ly	g aga tca s Arg Ser 40	aga tcc Arg Ser	gat ttc Asp Phe	cac cac His His 45	caa aac Gln Asn	ctc 201 Leu
act gag ga Thr Glu Gl 50	u Glu Ty	r Leu Ala 55	Phe Cys	Leu Met	ren ren	Ala Alg	65 65
aac cgt ca Asn Arg Gl	n Pro Pro 70	o Pro Pro	Pro Ala	Val Glu 75	rys ren	80	пАв
tgt agc gt Cys Ser Va	1 Cys As 85	p Lys Thr	Phe Ser	Ser Tyr	GIN AIA	95	GIY
cac aag go His Lys Al	a Ser Hi	в Arg Lys	Asn Leu 105	Ser Gin	Thr Leu 110	ser Gry	GIY
gga gat ga Gly Asp An 115	p His Se	r Thr Ser 120	Ser Ala	Thr Thr	125	Ala Val	·
act gga ag Thr Gly So 130	er Gly Ly	s Ser His	val Cys	140	сув Авп	гув зег	145
cct tcc g Pro Ser G	ly Gln Al 15	a Leu Gly O	Gly His	Lys Arg	сув нів	160	GIY
aac aac a	ac atc aa	c act agt	agc gtg	tcc aac Page		ggt gcg	ggg 585

Asn	Asn	Asn	Ile 165	Asn	Thr	Ser			eque Ser						Gly	
					agc Ser											633
					ttc Phe											681
					aag Lys 215											729
	ctt Leu	taa	ggaa	att	ac t	taga	acgat	ta ag	gatt	cgtl	t tgl	tata	tgt			778
tgaç	gagtt	gt	gtagg	gaatt	t gt	tgad	etgta	a cat	acca	aat	tgga	actt	ga (etga	ttccaa	838
ttct	tctt	gt t	cttt	catt	t ta	aaaa	attat	taa	aacc	gatt	ctt	acca	aca a	aa		890
<210 <211 <212 <213	l> 2 !> I !> <i>I</i>	24 227 PRT Arabi	idops	sis t	hali	iana										
			Glu	Ala	Leu	Thr	Ser	Pro	Ara	Leu	Ala	Ser	Pro	Ile	Pro	
1		200	-	5	204		501		10	204		501		15		
Pro	Leu	Phe	Glu 20	Asp	Ser	Ser	Val	Phe 25	His	Gly	Val	Glu	His 30	Trp	Thr	
Lys	Gly	Lys 35	Arg	Ser	Lys	Arg	Ser 40	Arg	Ser	Asp	Phe	His 45	His	Gln	Asn	
Leu	Thr 50	Glu	Glu	Glu	Tyr	Leu 55	Ala	Phe	Суз	Leu	Met 60	Leu	Leu	Ala	Arg	•
Авр 55	Asn	Arg	Gln	Pro	Pro 70	Pro	Pro	Pro	Ala	Val 75	Glu	Lys	Leu	Ser	Tyr 80	
Lys	Сув	Ser	Val	Cys 85	Asp	Lys	Thr	Phe	Ser 90	Ser	Tyr	Gln	Ala	Leu 95	Gly	
3ly	His	Lys	Ala 100	Ser	His	Arg	Lys	Asn 105	Leu	Ser	Gln	Thr	Leu 110	Ser	Gly .	
3ly	Gly	Asp 115	Asp	His	Ser	Thr	Ser 120	Ser	Ala	Thr	Thr	Thr 125	Ser	Ala	Val	
Thr	Thr 130	Gly	Ser	Gly	Lys	Ser 135	His	Val	Cys	Thr	Ile 140	Сув	Asn	Lys	Ser	
Phe 145	Pro	Ser	Gly	Gln	Ala 150	Leu	Gly	Gly	His	Lys 155	Arg	Сув	His	Tyr	Glu 160	
3ly	Asn	Asn	Asn	Ile	Asn	Thr	Ser	Ser	Val	Ser	Asn	Ser	Glu	Gly	Ala	

MBI15 Sequence Listing.ST25
Gly Ser Thr Ser His Val Ser Ser Ser His Arg Gly Phe Asp Leu Asn
180 185 190

Ile Pro Pro Ile Pro Glu Phe Ser Met Val Asn Gly Asp Asp Glu Val

Met Ser Pro Met Pro Ala Lys Lys Pro Arg Phe Asp Phe Pro Val Lys 210 215 220

Leu Gln Leu

<210> 25 <211> 1126

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (282)..(920)

<223> G865

<400> 25 atccccactt gttgttcatc accaagccaa gctccatgtc ctagtcactc cacagattcc 60 120 ctatcatcat caattogttt caaacttagt tootttcaaa gtottgtaca tatatacaca cacacctatt attotottgg tgtgtttgtg tgttacatat acgtgtgagt acatactttg 180 ttgtaaaagt ggatcggagg tatggaaagg gaccggttcc accggaaaca tcggcggcgg 240 cggatgataa ttcgtcttgg aacgagactg atgtcaccgc c atg gtc tcc gct ctc Met Val Ser Ala Leu 296 age cgt gtc ata gag aat ccg aca gac ccg ccg gtc aaa caa gag ctt Ser Arg Val Ile Glu Asn Pro Thr Asp Pro Pro Val Lys Gln Glu Leu 344 gat aaa tcg gat caa cat caa cca gac caa gat caa cca aga aga aga Asp Lys Ser Asp Gln His Gln Pro Asp Gln Asp Gln Pro Arg Arg 392 cac tat aga ggc gta agg cag aga cca tgg ggt aaa tgg gcg gca gaa His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly Lys Trp Ala Ala Glu 440 atc cgc gat cca aag aaa gca gcc cgt gtc tgg ctc ggg act ttc gag Ile Arg Asp Pro Lys Lys Ala Ala Arg Val Trp Leu Gly Thr Phe Glu 488 acg gca gag gaa gct gct tta gcc tat gac cga gct gcc ctc aaa ttc Thr Ala Glu Glu Ala Ala Leu Ala Tyr Asp Arg Ala Ala Leu Lys Phe 536 aaa ggc acc aag gct aaa ctg aac ttc cct gaa cgg gtc caa ggc cct Lys Gly Thr Lys Ala Lys Leu Asn Phe Pro Glu Arg Val Gln Gly Pro 584 95 act acc acc aca acc att tct cat gca cca aga gga gtt agt gaa tcc Thr Thr Thr Thr Ile Ser His Ala Pro Arg Gly Val Ser Glu Ser 632 atg aac tca cct cct cct cga cct ggt cca cct tca act act act act Met Asn Ser Pro Pro Pro Arg Pro Gly Pro Pro Ser Thr Thr Thr 680 tcg tgg cca atg act tat aac cag gac ata ctt caa tac gct cag ttg Ser Trp Pro Met Thr Tyr Asn Gln Asp Ile Leu Gln Tyr Ala Gln Leu 135 140 145 728

MRTIC	Sequence	Lighing	СТЭБ
MBITZ	Sequence	LIBCING.	. 5125

ctt acg agt aac aat gag gtt gat tta tca tac tac acg tcg act ctc Leu Thr Ser Asn Asn Glu Val Asp Leu Ser Tyr Tyr Thr Ser Thr Leu 150 155 160 165	776
ttc agt caa cct ttt tca acg cct tct tca tct tct tct tcc tcc caa Phe Ser Gln Pro Phe Ser Thr Pro Ser Ser Ser Ser Ser Ser Gln 170 175 180	824
cag acg cag caa cag cag cta caa caa caa caa cag cag cgt gaa gaa Gln Thr Gln	872
gaa gag aag aat tat ggt tac aat tat tat aac tac cca aga gaa taa Glu Glu Lys Asn Tyr Gly Tyr Asn Tyr Tyr Asn Tyr Pro Arg Glu 200 205 210	920
tctaattatt attgttggtc gaatcagttt tataaatagc tatcatagtt tcatttttgg	980
tttccgtaac ctttgttgca tggaaaatat gaatgaacga gggacatgtg taacaatttg 10	040
tttgtgtttc gtaaatgtta gttgtatttg gatttgctga agtttgattt tctgagcata 11	100
aatcatttga cggtcaaaaa aaaaaa 11	126
<210> 26 <211> 212	

PRT

Arabidopsis thaliana

<400> 26

Met Val Ser Ala Leu Ser Arg Val Ile Glu Asn Pro Thr Asp Pro Pro

Val Lys Gln Glu Leu Asp Lys Ser Asp Gln His Gln Pro Asp Gln Asp 20 25 30

Gln Pro Arg Arg Arg His Tyr Arg Gly Val Arg Gln Arg Pro Trp Gly 35 40 .

Lys Trp Ala Ala Glu Ile Arg Asp Pro Lys Lys Ala Ala Arg Val Trp 50 55 60

Leu Gly Thr Phe Glu Thr Ala Glu Glu Ala Ala Leu Ala Tyr Asp Arg 65 70 80

Ala Ala Leu Lys Phe Lys Gly Thr Lys Ala Lys Leu Asn Phe Pro Glu 85 90 95

Arg Val Gln Gly Pro Thr Thr Thr Thr Thr Ile Ser His Ala Pro Arg

Gly Val Ser Glu Ser Met Asn Ser Pro Pro Pro Arg Pro Gly Pro Pro 115 120 125

Ser Thr Thr Thr Thr Ser Trp Pro Met Thr Tyr Asn Gln Asp Ile Leu 130 135 140

Gln Tyr Ala Gln Leu Leu Thr Ser Asn Asn Glu Val Asp Leu Ser Tyr

Tyr Thr Ser Thr Leu Phe Ser Gln Pro Phe Ser Thr Pro Ser Ser

Ser Ser Ser Ser Gln Gln Thr Gln Gln Gln Gln Leu Gln Gln Gln

Gln Gln Arg Glu Glu Glu Lys Asn Tyr Gly Tyr Asn Tyr Tyr Asn 200

Tyr Pro Arg Glu

<210> 27 <211> 1152 <212> DNA <213> Arabidopsis thaliana <220> <221> CDS (76) .. (1008) <222> <223> G881 <400> 27 gtgaccgaag aaagcaaatt gagactacgc accaactagt cctttggttt gtatcttaag 60 ataaaggttt ctttt atg gac ggt tct tcg ttt ctc gac atc tct ctc gat
Met Asp Gly Ser Ser Phe Leu Asp Ile Ser Leu Asp 111 ctc aac acc aat cct ttc tcc gca aaa ctt ccg aag aag gag gtc tca Leu Asn Thr Asn Pro Phe Ser Ala Lys Leu Pro Lys Lys Glu Val Ser 159 gtt ttg gct tct act cac tta aag agg aaa tgg ttg gag caa gac gag Val Leu Ala Ser Thr His Leu Lys Arg Lys Trp Leu Glu Gln Asp Glu 207 35 agc gca agt gag tta cga gag gag cta aac aga gtt aat tca gag aac Ser Ala Ser Glu Leu Arg Glu Glu Leu Asn Arg Val Asn Ser Glu Asn 255 aag aag cta aca gag atg tta gct aga gtc tgt gag agc tac aac gaa Lys Lys Leu Thr Glu Met Leu Ala Arg Val Cys Glu Ser Tyr Asn Glu 65 70 75303 cta cat aat cat ttg gag aag ctt cag agt cgc cag agc cct gaa atc Leu His Asn His Leu Glu Lys Leu Gln Ser Arg Gln Ser Pro Glu Ile 351 85 gag cag acc gat ata ccg ata aag aaa aga aaa caa gac ccg gat gag Glu Gln Thr Asp Ile Pro Ile Lys Lys Arg Lys Gln Asp Pro Asp Glu 399 ttc tta ggc ttt cct att gga ctc agt agt gga aaa act gag aac agc Phe Leu Gly Phe Pro Ile Gly Leu Ser Ser Gly Lys Thr Glu Asn Ser 110 115 120 447 tcc agc aac gaa gat cat cat cat cat cat cag caa cat gag cag aaa Ser Ser Asn Glu Asp His His His His Gln Gln His Glu Gln Lys 495 135 aat cag ctt ctt tca tgt aaa aga cca gtc act gat agc ttc aac aaa Asn Gln Leu Ser Cys Lys Arg Pro Val Thr Asp Ser Phe Asn Lys 543 gca aaa gtt tcg act gtc tac gtg cct act gaa aca tcg gac aca agc Ala Lys Val Ser Thr Val Tyr Val Pro Thr Glu Thr Ser Asp Thr Ser 591

Page 36

185

639

ttg aca gtt aaa gat gga ttt caa tgg agg aaa tac gga caa aag gtt Leu Thr Val Lys Asp Gly Phe Gln Trp Arg Lys Tyr Gly Gln Lys Val

180

														ttt Phe		687
ccg Pro 205	tct Ser	tgt Cys	cca Pro	gta Val	aaa Lys 210	aag Lys	aag Lys	gta Val	caa Gln	cgc Arg 215	agc Ser	gca Ala	gag Glu	gat Asp	cca Pro 220	735
tct Ser	tta Leu	ctt Leu	gta Val	gcg Ala 225	aca Thr	tac Tyr	gaa Glu	999 Gly	acg Thr 230	cat His	aac Asn	cac His	ttg Leu	ggt Gly 235	cca Pro	783
aat Asn	gct Ala	tct Ser	gaa Glu 240	G1y 999	gat Asp	gct Ala	Thr	agc Ser 245	cag Gln	ggt Gly	G1 y 999	tca Ser	agc Ser 250	aca Thr	gtg Val	831
														aaa Lys		879
gaa Glu	agg Arg 270	gat Asp	aat Asn	acg Thr	atg Met	caa Gln 275	gag Glu	gtt Val	ctg Leu	att Ile	caa Gln 280	caa Gln	atg Met	gcg Ala	tca Ser	927
tcg Ser 285	tta Leu	aca Thr	aaa Lys	gat Asp	tcg Ser 290	aaa Lys	ttt Phe	aca Thr	gct Ala	gct Ala 295	ctt Leu	gct Ala	gct Ala	gct Ala	ata Ile 300	975
tct Ser	Gly ggg	agg Arg	tta Leu	atg Met 305	gag Glu	caa Gln	tct Ser	aga Arg	aca Thr 310	tga	acg	ttt	tag	tgaa	tgtatt	1028
gtt	ttg	ttt	gttt	agaa	tg a	tct	tcgt	t tt	cgaai	ttgt	gtc	tttc	gat	tagg	agataa	1088
aaga	atgt	ata 1	taaa	tatt	at aa	agtag	gatg	a ag	aaat	egta	taa	gtaa	aaa a	aaaa	aaaaa	1148
aaa	a															1152

<210> 28

<211> 310 <212> PRT

<213> Arabidopsis thaliana

<400> 28

Met Asp Gly Ser Ser Phe Leu Asp Ile Ser Leu Asp Leu Asn Thr Asn 1 10 15

Pro Phe Ser Ala Lys Leu Pro Lys Lys Glu Val Ser Val Leu Ala Ser 25 30

Thr His Leu Lys Arg Lys Trp Leu Glu Gln Asp Glu Ser Ala Ser Glu 35 40 45

Leu Arg Glu Glu Leu Asn Arg Val Asn Ser Glu Asn Lys Lys Leu Thr 50 55 60

Glu Met Leu Ala Arg Val Cys Glu Ser Tyr Asn Glu Leu His Asn His 65 70 75 80

Leu Glu Lys Leu Gln Ser Arg Gln Ser Pro Glu Ile Glu Gln Thr Asp 85 90 95

Ile Pro Ile Lys Lys Arg Lys Gln Asp Pro Asp Glu Phe Leu Gly Phe 100 105 110

?ro	Ile	Gly 115	Leu	Ser	Ser	Gly	MBI: Lys 120	15 Se Thr	equer Glu	Asn	Ser	Ser 125	Ser	Asn	Glu	
Asp	His 130	His	His	His	His	Gln 135	Gln	His	Glu	Gln	Lys 140	Asn	Gln	Leu	Leu	
Ser 145	Cys	Lys	Arg	Pro	Val 150	Thr	Asp	Ser	Phe	Asn 155	Lys	Ala	Lys	Val	Ser 160	
Thr	Val	Tyr	Val	Pro 165	Thr	Glu	Thr	Ser	Asp 170	Thr	Ser	Leu	Thr	Val 175	Lys	
Asp	Gly	Phe	Gln 180	Trp	Arg	Lys	Tyr	Gly 185	Gln	Lys	Val	Thr	Arg 190	Asp	Asn	
Pro	Ser	Pro 195		Ala	Tyr	Phe	Arg 200	Сув	Seŗ	Phe	Ala	Pro 205	Ser	Сув	Pro	
Val	Lys 210		Lys	Val	Gln	Arg 215	Ser	Ala	Glu	Asp	Pro 220	Ser	Leu	Leu	Val	
Ala 225		Tyr	Glu	Gly	Thr 230	His	Asn	His	Leu	Gly 235	Pro	Asn	Ala	Ser	Glu 240	
Gly	Asp	Ala	Thr	Ser 245		Gly	Gly	Ser	Ser 250	Thr	Val	Thr	Leu	Asp 255	Leu	
Val	Asn	Gly	Cys 260	His	Arg	Leu	Ala	Leu 265	Glu	Lys	Asn	Glu	Arg 270	Asp	Asn	
Thr	Met	Gln 275		ı Val	Leu	Ile	Gln 280	Gln	Met	Ala	Ser	Ser 285	Leu	Thr	. FAs	
Asp	Ser 290		Phe	Thr	. Ala	Ala 295	Leu		Ala	Ala	.·Il∈ 300	e Ser	Gly	Arg	Leu	
Met 305		ı Glr	sei	Arg	310	;										
<21 <21 <21 <21	1> 12>	29 1276 DNA Aral		psis	thal	liana	ı									
<22	20> 21> 22> 23>	CDS (47) G89		1150)		,									
<40 ta	00> atcc	29 gatt	cgt	cttc	atc 1	tgatt	ccc	tc c	cttco	egaga	a at	aata	atg Met 1	tac Tyr	ccg Pro	55
Pr	a cc o Pr 5	t cc o Pr	c tc o Se	a ag r Se	c ater	c tac e Ty: 10	c gc	t cc a Pr	t cc	g ate	g ct t Le 15	u va	g aa l As	t tg n Cy	c tcc s Ser	103
99 G1 20	у Су	c cg s Ar	g ac g Th	g cc r Pr	t ct o Le 25	u Gl	g ct n Le	c cc u Pr	o se	30	у АТ	c cg a Ar	a tc g Se	t at r Il	t cgc e Arg 35	151
										Page	38					

								13 3	cque	nce	DIBC	ıng.	3123			
						gtt Val										. 199
						tcc Ser										247
						cct Pro										295
						cgt Arg 90										343
						atg Met										391
						ctc Leu										439
						caa Gln	Asn									487
gta Val	cag Gln	gga Gly 150	tgc Cys	aca Thr	gca Ala	ggc Gly	gac Asp 155	tca Ser	ctt Leu	gtc Val	ttc Phe	cac His 160	tac Tyr	tct Ser	ggt Gly	535
cat His	ggt Gly 165	tcg Ser	cgt Arg	caa Gln	aga Arg	aac Asn 170	tac Tyr	aac Asn	ggt Gly	gat Asp	gaa Glu 175	gtt Val	gat Asp	ggc Gly	tat Tyr	583
						ctg Leu										631
						acc Thr										679
						gat Asp										727
						atg Met										775
						ggt Gly 250										823
						tgt Cys										871
						acg Thr										919
att Ile	caa Gln	gca Ala	att Ile 295	gaa Glu	cgc Arg	agc Ser	gca Ala	caa Gln 300	ggc Gly	aca Thr	acc Thr	tat Tyr	gga Gly 305	agc Ser	ctt Leu	967
						aca Thr										1015
						acg Thr									g1y.	1063

MBI15 Sequence Listing.ST25 325 330 335	
gga agt gcg att ggg gga tta aga cag gag cct caa ctg act gct tgc Gly Ser Ala Ile Gly Gly Leu Arg Gln Glu Pro Gln Leu Thr Ala Cys 340 345 350 355	1111
caa aca ttc gat gtc tat gca aag cct ttc act ctc tag taaaggacaa Gln Thr Phe Asp Val Tyr Ala Lys Pro Phe Thr Leu 360 365	1160
gtcacttttt atgtatagcg agtgtgattt gagaatccgt ccatataacc accttttgtt	1220
tottattttt atttttottt caaaagaata aaggaaaaca ttgatttggt gattcg	1276
<210> 30 <211> 367 <212> PRT <213> Arabidopsis thaliana	
<400> 30	
Met Tyr Pro Pro Pro Pro Ser Ser Ile Tyr Ala Pro Pro Met Leu Val 1 5 10	
Asn Cys Ser Gly Cys Arg Thr Pro Leu Gln Leu Pro Ser Gly Ala Arg 20 25 30	
Ser Ile Arg Cys Ala Leu Cys Gln Ala Val Thr His Ile Ala Asp Pro 35 40 45	
Arg Thr Ala Pro Pro Pro Gln Pro Ser Ser Ala Pro Ser Pro Pro Pro 50 55 60	
Gln Ile His Ala Pro Pro Gly Gln Leu Pro His Pro His Gly Arg Lys 65 70 75 80	
Arg Ala Val Ile Cys Gly Ile Ser Tyr Arg Phe Ser Arg His Glu Leu 85 90 95	
Lys Gly Cys Ile Asn Asp Ala Lys Cys Met Arg His Leu Leu Ile Asn 100 105 110	
Lys Phe Lys Phe Ser Pro Asp Ser Ile Leu Met Leu Thr Glu Glu Glu 115 120 125	
Thr Asp Pro Tyr Arg Ile Pro Thr Lys Gln Asn Met Arg Met Ala Leu 130 135 140	
Tyr Trp Leu Val Gln Gly Cys Thr Ala Gly Asp Ser Leu Val Phe His 145 150 155 160	
Tyr Ser Gly His Gly Ser Arg Gln Arg Asn Tyr Asn Gly Asp Glu Val 165 170 175	
Asp Gly Tyr Asp Glu Thr Leu Cys Pro Leu Asp Phe Glu Thr Gln Gly 180 185 190	
Met Ile Val Asp Asp Glu Ile Asn Ala Thr Ile Val Arg Pro Leu Pro 195 200 205	
His Gly Val Lys Leu His Ser Ile Ile Asp Ala Cys His Ser Gly Thr Page 40	

192

MBI15 Sequence Listing.ST25 210

Val Leu Asp Leu Pro Phe Leu Cys Arg Met Asn Arg Ala Gly Gln Tyr 230

Val Trp Glu Asp His Arg Pro Arg Ser Gly Leu Trp Lys Gly Thr Ala

Gly Gly Glu Ala Ile Ser Ile Ser Gly Cys Asp Asp Asp Gln Thr Ser

Ala Asp Thr Ser Ala Leu Ser Lys Ile Thr Ser Thr Gly Ala Met Thr

Phe Cys Phe Ile Gln Ala Ile Glu Arg Ser Ala Gln Gly Thr Thr Tyr

Gly Ser Leu Leu Asn Ser Met Arg Thr Thr Ile Arg Asn Thr Gly Asn

Asp Gly Gly Gly Ser Gly Gly Val Val Thr Thr Val Leu Ser Met Leu 325 330 335

Leu Thr Gly Gly Ser Ala Ile Gly Gly Leu Arg Gln Glu Pro Gln Leu

Thr Ala Cys Gln Thr Phe Asp Val Tyr Ala Lys Pro Phe Thr Leu

<210>

<211> 726

<212> DNA

Arabidopsis thaliana

<220>

<221>

(1)..(726) <222>

G378

<400> 31 atg gcc tcg tca tca tca tct tat aga ttc caa tct ggg tct tac Met Ala Ser Ser Ser Ser Ser Tyr Arg Phe Gln Ser Gly Ser Tyr 48

cct ctt tcg tca agt cct tct ctt ggg aat ttc gtc gaa cgc att aaa Pro Leu Ser Ser Pro Ser Leu Gly Asn Phe Val Glu Arg Ile Lys 96

gac gct tgt cat ttc ctt gtc tct gct gtt ttg ggt acc att atc tcc Asp Ala Cys His Phe Leu Val Ser Ala Val Leu Gly Thr Ile Ile Ser 144

gcg atc ttg acc ttc ttc ttc gca cta gtg ggc aca ttg cta ggg gca Ala Ile Leu Thr Phe Phe Phe Ala Leu Val Gly Thr Leu Leu Gly Ala

ctt aca gga gct ttg ata ggt caa gaa act gag agt ggt ttc att aga Leu Thr Gly Ala Leu Ile Gly Gln Glu Thr Glu Ser Gly Phe Ile Arg 240

gga gca gca att gga gcc att tcg gga gct gtt ttc tct atc gag gtc Gly Ala Ala Ile Gly Ala Ile Ser Gly Ala Val Phe Ser Ile Glu Val 85 90 95 288

Phe Glu Ser Ser Leu Asp Leu Trp Lys Ser Asp Glu Ser Gly Phe Gly

MBI15 Sequence Listing.ST25

Cys Phe Leu Tyr Leu Ile Asp Val Ile Val Ser Leu Leu Ser Gly Arg

Leu Val Arg Glu Arg Ile Gly Pro Ala Met Leu Ser Ala Val Gln Ser 130 135 140

Gln Met Gly Ala Val Asp Thr Ala Phe Asp Asp His Thr Ser Leu Phe 145 150 155 160

Asp Thr Gly Gly Ser Lys Gly Leu Thr Gly Asp Leu Val Glu Lys Ile 165 170 175

Pro Lys Met Thr Ile Thr Gly Asn Asn Asn Thr Asp Ala Ser Glu Asn 180 185 190

Thr Asp Ser Cys Ser Val Cys Leu Gln Asp Phe Gln Leu Gly Glu Thr 195 200 205

Val Arg Ser Leu Pro His Cys His His Met Phe His Leu Pro Cys Ile 210 215 220

Asp Asn Trp Leu Leu Arg His Gly Ser Cys Pro Met Cys Arg Arg Asp 225 230 235 240

Ile

<210> 33

<211> 1370

<212> DNA

<213> Arabidopsis thaliana

<220>

<221> CDS

<222> (184)..(969)

<223> G569

<400> 33 gtcgacccac gcgtccgggt ttttctttta tcctcttatc gctaatctgg agctctatat 60 agactataaa gggtttttga ttgattcggg agctcgagat ttgacttctt ttagctgatt 120 cggcaagttt gtatctagaa aggatcgatt ggtgaggtca atagtggttg gtgggtttta 180 gta atg gaa gac ggt gag ctt gat ttc tcc aat cag gaa gtg ttt tcg
Met Glu Asp Gly Glu Leu Asp Phe Ser Asn Gln Glu Val Phe Ser 228 agt tcg gag atg ggt gaa tta cca cct agc aat tgt tcg atg gat agt Ser Ser Glu Met Gly Glu Leu Pro Pro Ser Asn Cys Ser Met Asp Ser 276 ttc ttt gat ggg ctt tta atg gat act aat gct gct tgt acc cac act Phe Phe Asp Gly Leu Leu Met Asp Thr Asn Ala Ala Cys Thr His Thr 324 cac acc tgt aac ccc act gga cca gag aac act cat act cac acg tgc His Thr Cys Asn Pro Thr Gly Pro Glu Asn Thr His Thr His Thr Cys 372 ttc cat gtc cac acc aag att ctc ccg gat gag agc gat gaa aaa gtt Phe His Val His Thr Lys Ile Leu Pro Asp Glu Ser Asp Glu Lys Val 420

70

tet	act	aat	oat	aca	act	gag	MBI:	l5 S(tat	eque:	nce 1 aaq	List: aaq	ing.:	gaa	aag	aga	468
Ser	Thr	Asp	Asp	Thr	Ala 85	Ğlü	Ser	Cys	ĞĬÿ	Lys 90	Lys	ĞĨy	Ğlu	ГÀв	Arg 95	
80													-	226	220	516
cct Pro	ttg Leu	gga Glv	aac Asn	cgg Arg	gaa Glu	gcg Ala	gtt Val	aga Arg	aag Lys	Tyr	aga Arg	gag Glu	aag Lys	nys	Lys	310
		•		100					105					110		
gct	aaa	gct	gct	tct	ttg	gag	gat	gag	gtt	gca	agg	ctt	agg	gcg	gtg Val	564
Ala	Lys	Ala	Ala 115	Ser	Leu	GIU	Asp	120	vai	Ala	Aig	neu	Arg 125	A.u	742	
22 +	can	caq	cto	ata	aao	agg	tta	caa	aat	caq	gct	acc	ttg	gaa	gct	612
Asn	Gln	Gln	Leu	Val	Lys	Arg	Leu 135	Gln	Asn	Gln	Āla	Thr 140	Leu	Glu	Ala	
		130										_	~~~	303	ata	660
gag Glu	gtt Val	tcg Ser	agg	ctt Leu	aag Lys	Cys	Leu	CCC Lėu	Val	Asp	Leu	Arg	gga Gly	Arg	Ile	000
	145					150					155					
gat	gga	gag	att	gga	tct	ttt	cct	tat	cag	aaa	cct	atg	gct Ala	gca	aat Asn	708
Asp 160	Gly	GIu	TTE	GIY	165	Pne	PLO	ıyı	GIII	170	110	1100	Āla		175	
att	cct	tet	ttc	tcq	cac	atq	atg	aat	cct	tgt	aat	gta	caa	tgt	gat	756
Ile	Pro	Ser	Phe	Ser 180	His	Met	Met	Asn	Pro 185	Сув	Asn	Val	Gln	Сув 190	Asp	
										~~~	ata	221	200	caa	gaa	B04
gat Asp	gaa Glu	gtt Val	tat Tyr	Cys	Pro	Gln	Asn	Val	Phe	Gly	Val	Asn	agc	GIII	Glu	
			195					200					205			
ggt	gcc	tcg	atc	aat	gac	caa	999	tta	agt	ggt	tgt	gat	ttt Phe	gat Asp	cag Gln	852
GIY	Ala	210		ASII	Азр	GIII	215	ДС	001	027	-7-	220	)	•		
cta	caa	tac	atq	gct	aat	cag	aac	tta	aat	gga	aat	gga	aac	gga	tca	900
Leu	Gln 225	Cys	Met	Āla	Asn	Gln 230	. Asn	Leu	Asn	Gly	Asn 235	GIY	Asn Asn	GIY	Ser	
	_							tro	aat	aac	aga	aaa	aat	ggg	cat	948
Phe	Ser	Asn	Val	Asn	Thr	Ser	Val	Ser	Asn	грув	Arg	Lys	ĞĨy	Gĺy	His 255	
240					245	•				250					-	999
cgt	gca	tca	aga	gca	gtt Val	tga	ago	atca	tca	agct	tgta	ict a	tcta	tttc	C	999
77.5	, ,,,,,			260	)											
acc	agca	itag	atat	tgta	itt c	caaa	taag	ıt t <u>e</u>	ıtaga	gtto	ago	tgca	agga	tcag	cttcgc	1059
tca	ıgctt	:tga	gggg	ittgg	ıtg g	tgtg	gtct	t to	tttg	tggc	acg	agt	gaga	tcta	tggaca	1119
															gttgct	1179
															ıggggct	1239
																1299
-															agttgc	1359
tti	gtt	gact	taaa	atgt	gat a	atat	agct	t a	ictti	tacı	t tga	aaaa	aaaa	aaaa	aaaaaa	
aaa	aaaa	aaaa	a													1370
	10> 11>	34 261														
<2	12> 13>	PRT	hi đơ	osie	tha:	lian	a									
	00>	34	٠		<b>.</b> .		_ ==		~ n~	n (21)	יוים ח	u Va	) Dh	e Se	r Ser	
Me 1	t Gl	u As	p GI	y G1	u Le	u AS	h hu	e 5e	10			_ •a		15	r Ser	

Ser Glu Met Gly Glu Leu Pro Pro Ser Asn Cys Ser Met Asp Ser Phe Page 44

MBI15 Sequence Listing.ST25

Phe Asp Gly Leu Leu Met Asp Thr Asn Ala Ala Cys Thr His Thr His 35 40 45

Thr Cys Asn Pro Thr Gly Pro Glu Asn Thr His Thr His Thr Cys Phe 50 55 60

His Val His Thr Lys Ile Leu Pro Asp Glu Ser Asp Glu Lys Val Ser 65 70 75 80

Thr Asp Asp Thr Ala Glu Ser Cys Gly Lys Lys Gly Glu Lys Arg Pro 85 90 95

Leu Gly Asn Arg Glu Ala Val Arg Lys Tyr Arg Glu Lys Lys Lys Ala 100 105 110

Lys Ala Ala Ser Leu Glu Asp Glu Val Ala Arg Leu Arg Ala Val Asn 115 120 125

Gln Gln Leu Val Lys Arg Leu Gln Asn Gln Ala Thr Leu Glu Ala Glu 130 135 140

Val Ser Arg Leu Lys Cys Leu Leu Val Asp Leu Arg Gly Arg Ile Asp 145 150 150 160

Gly Glu Ile Gly Ser Phe Pro Tyr Gln Lys Pro Met Ala Ala Asn Ile 165 170 175

Pro Ser Phe Ser His Met Met Asn Pro Cys Asn Val Gln Cys Asp Asp 180 190

Glu Val Tyr Cys Pro Gln Asn Val Phe Gly Val Asn Ser Gln Glu Gly

Ala Ser Ile Asn Asp Gln Gly Leu Ser Gly Cys Asp Phe Asp Gln Leu 210 215 220

Gln Cys Met Ala Asn Gln Asn Leu Asn Gly Asn Gly Asn Gly Ser Phe 225 230 235 240

Ser Asn Val Asn Thr Ser Val Ser Asn Lys Arg Lys Gly Gly His Arg 245 250 255

Ala Ser Arg Ala Val 260

<210> 35

<211> 1638

<212> DNA

<213> Arabidopsis thaliana

<220>

221> CDS

<222> (267)..(1259)

<223> G558

<400> 35

										<b>.</b>		0	ma r			
ggaa	tttc	gg a	tcgt	gtct	c tc	tctg	MBI1 tttc	5 Se	quen gttt	caa	tccg	ng.s	cg a	atca	agccc	60
ttta	cttg	tg c	acct	tcaa	g at	ttcg	tttt	ttc	cagc	gcc	caga	atgc	tc c	gggt	gacca	120
acat	ttgt	tc c	tgat	tcat	t tc	ctat	tggt	tcg	tatt	gtc	tgtg	caca	ca a	gaga	aattt	180
caag	aagt	tg t	tact	aaaa	g ag	aggc	caca	agt	ggat	att	gtct	ttgt	ta t	caag	tgtta	240
gtac	agaa	aa g	tggt	gaga	a ag	taat	atg Met 1	gct Ala	gat Asp	acc Thr	agt Ser 5	ccg Pro	aga Arg	act Thr	gat Asp	293
gtc Val 10	tca Ser	aca Thr	gat Asp	gac Asp	gac Asp 15	aca Thr	gat Asp	cat His	Pro	gat Asp 20	ctt Leu	999 Gly	tcg Ser	gag Glu	gga Gly 25	341
gca Ala	cta Leu	gtg Val	aat Asn	act Thr 30	gct Ala	gct Ala	tct Ser	gat Asp	tcg Ser 35	agt Ser	gac Asp	cga Arg	tcg Ser	aag Lys 40	gga Gly	389
aag Lys	atg Met	gat Asp	caa Gln 45	aag Lys	act Thr	ctt Leu	cgt Arg	agg Arg 50	ctt Leu	gct Ala	caa Gln	aac Asn	cgt Arg 55	gag Glu	gca Ala	437
gca Ala	agg Arg	aaa Lys 60	agc Ser	aga Arg	ttg Leu	agg Arg	aag Lys 65	aag Lys	gct Ala	tat Tyr	gtt Val	cag Gln 70	cag Gln	cta Leu	gag Glu	485
aac Asn	agc Ser 75	cgc Arg	ttg Leu	aaa Lys	cta Leu	acc Thr 80	cag Gln	ctt Leu	gag Glu	cag Gln	gag Glu 85	ctg Leu	caa Gln	aga Arg	gca Ala	533
aga Arg 90	cag Gln	cag Gln	ggc Gly	gtc Val	ttc Phe 95	att Ile	tca Ser	ggc Gly	aca Thr	gga Gly 100	gac Asp	cag Gln	gcc Ala	cat His	tct Ser 105	581
act Thr	ggt Gly	gga Gly	aat Asn	ggt Gly 110	gct Ala	ttg Leu	gcg Ala	ttt Phe	gat Asp 115	gct Ala	gaa Glu	cat His	tca Ser	cgg Arg 120	tgg Trp	629
ttg Leu	gaa Glu	gaa Glu	aag Lys 125	aac Asn	aag Lys	caa Gln	atg Met	aac Asn 130	gag Glu	ctg Leu	agg Arg	tct Ser	gct Ala 135	ctg Leu	aat Asn	677
gcg Ala	cat His	gca Ala 140	ggt Gly	gat Asp	tct Ser	gag Glu	ctt Leu 145	cga Arg	ata Ile	ata Ile	gtc Val	gat Asp 150	ggt Gly	gtg Val	atg Met	725
gct Ala	cac His 155	tat Tyr	gag Glu	gag Glu	ctt Leu	ttc Phe 160	agg Arg	ata Ile	aag Lys	agc Ser	aat Asn 165	gca Ala	gct Ala	aag Lys	aat Asn	773
gat Asp 170	Val	ttt Phe	cac His	ttg Leu	cta Leu 175	Ser	ggc Gly	atg Met	tgg Trp	aaa Lys 180	aca Thr	cca Pro	gct Ala	gag Glu	aga Arg 185	821
tgt Cys	ttc Phe	ttg Leu	tgg Trp	ctc Leu 190	Gly	gga Gly	ttt Phe	cgt Arg	tca Ser 195	ser	gaa Glu	ctt Leu	cta Leu	aag Lys 200	ctt Leu	869
ctg Leu	gcg Ala	aat Asn	cag Gln 205	Leu	gag Glu	cca Pro	atg Met	aca Thr 210	GIU	aga Arg	cag Gln	ttg Leu	atg Met 215	Gry	ata Ile	917
Asn	Asn	Leu 220	Gln	Gln	Thr	Ser	225	Gin	Ala	GIU	Авр	230	Leu	Ser	caa Gln	965
Gly	Met 235	Glu	Ser	Lev	Gln	240	Ser	Leu	Ala	Asp	245	reu	261	261	Gly	1013
act Thi 250	Leu	ggt	tca Sex	agt Ser	Ser 255	Ser	999 Gly	aat Asn	vai	gca Ala 260	ser	tac Tyr	atg Met	ggt Gly	cag Gln 265	1061

#### MBI15 Sequence Listing.ST25

atg gcc atg gca atg gga aag tta ggt aca ctc gaa gga ttt atc cgc Met Ala Met Ala Met Gly Lys Leu Gly Thr Leu Glu Gly Phe Ile Arg 270 275 280	1109
cag gct gat aat ttg aga cta caa aca ttg caa cag atg ata aga gta Gln Ala Asp Asn Leu Arg Leu Gln Thr Leu Gln Gln Met Ile Arg Val 285 290 295	1157
tta aca acg aga cag tca gca cgt gct cta ctt gca ata cac gat tac Leu Thr Thr Arg Gln Ser Ala Arg Ala Leu Leu Ala Ile His Asp Tyr 300 305 310	1205
ttc tca cgg cta cga gct cta agc tcc tta tgg ctt gct cga ccc aga Phe Ser Arg Leu Arg Ala Leu Ser Ser Leu Trp Leu Ala Arg Pro Arg 315 320 325	1253
gag tga aactgtattt tggtcacatg tcagctgtac aaaatccata tggacacaaa Glu 330	1309
accaggagag actattaatc aacacttgtc agattettet taccaaatcc atcaacaaat	1369
aagcaaattt ctgggaaaca aaagactett tgtatgtagg tttettetae atggttgtgg	1429
taattcatgt tgttttagtt gtagtcatca gtttttaatt tagcatttga aaagttcaat	1489
gttgtttata tagcatcttc gattatctta gaaaggttat tgaattttgt ttttttttgt	1549
tacttttgtg tgtggtaaag gtgttttaac cttgcaactt ctgtactgta	1609
aatattaaga tgttctattt gagttttgt	1638

<210> 36

<400> 36

Met Ala Asp Thr Ser Pro Arg Thr Asp Val Ser Thr Asp Asp Asp Thr 1 5 10 15

Asp His Pro Asp Leu Gly Ser Glu Gly Ala Leu Val Asn Thr Ala Ala 20 25 30

Ser Asp Ser Ser Asp Arg Ser Lys Gly Lys Met Asp Gln Lys Thr Leu 35 40 45

Arg Arg Leu Ala Gln Asn Arg Glu Ala Ala Arg Lys Ser Arg Leu Arg 50 60

Lys Lys Ala Tyr Val Gln Gln Leu Glu Asn Ser Arg Leu Lys Leu Thr 65 70 75 80

Gln Leu Glu Gln Glu Leu Gln Arg Ala Arg Gln Gln Gly Val Phe Ile 85 90 95

Ser Gly Thr Gly Asp Gln Ala His Ser Thr Gly Gly Asn Gly Ala Leu 100 105 110

Ala Phe Asp Ala Glu His Ser Arg Trp Leu Glu Glu Lys Asn Lys Gln 115 120 125

Met Asn Glu Leu Arg Ser Ala Leu Asn Ala His Ala Gly Asp Ser Glu 130 135 140

<211> 330

<212> PRT

<213> Arabidopsis thaliana

Leu 145	Arg	Ile	Ile	Val	Asp 150	Gly	Val	Met	Ala	His 155	Tyr	Glu	Glu	Leu	Phe 160	
Arg	Ile	Lys	Ser	Asn 165	Ala	Ala	Lys	Asn	Asp 170	Val	Phe	His	Leu	Leu 175	Ser	
Gly	Met	Trp	Lys 180	Thr	Pro	Ala	Glu	Arg 185	Сув	Phe	Leu	Trp	Leu 190	Gly	Gly	
Phe	Arg	Ser 195	Ser	Glu	Leu	Leu	Lys 200	Leu	Leu	Ala	Asn	Gln 205	Leu	Glu	Pro	
Met	Thr 210	Glu	Arg	Gln	Leu	Met 215	Gly	Ile	Asn	Asn	Leu 220	Gln	Gln	Thr	Ser	
Gln 225	Gln	Ala	Glu	Asp	Ala 230	Leu	Ser	Gln	Gly	Met 235	Glu	Ser	Leu	Gln	Gln 240	
Ser	Leu	Ala	Asp	Thr 245	Leu	Ser	Ser	Gly	Thr 250	Leu	Gly	Ser	Ser	Ser 255	Ser	
Gly	Asn	Val	Ala 260	Ser	Tyr	Met	Gly	Gln 265	Met	Ala	Met	Ala	Met 270	Gly	Lys	
Leu	Gly	Thr 275		Glu	Gly	Phe	Ile 280		Gln	Ala	Asp	Asn 285	Leu	Arg	Leu	
Gln	Thr 290		Gln	Gln	Met	11e 295		Val	Lęu	Thr	Thr 300	Arg	Gln	Ser	Ala	
Arg 305		Leu	Leu	Ala	Ile 310		Asp	Tyr	Phe	Ser 315	Arg	Leu	Arg	Ala	Leu 320	
Ser	Ser	Leu	Trp	Leu 325		Arg	Pro	Arg	Glu 330							
<21 <21 <21 <21	1> 2>	37 436 DNA Arab	idop	sis	thal	iana										
<22 <22 <22 <22	1> 2>	CDS (83) G139	(3 6	13)												
<40 tcg	0> acct	37 cgt	ttcc	tttc	ct c	ctct	ctto	c ta	ıccat	tagt	acg	gttac	tgg	agct	gatctc	60
acg	įtata	ttt	tgga	tcgt	aa t	c at Me 1	g ga t As	ic gg sp Gl	gc ga ly Gl	a ga u As 5	it tt sp Pl	t go ne Al	c gg a Gl	a aa y Ly	ng gcg ys Ala 10	112
gct Ala	gct Ala	gaa Glu	gco Ala	aag Lys	g gga s Gly	ttg Lev	aac Asr	ccg Pro	g gga 5 Gly 20	tta Lev	ato 1116	gtg Val	cto Lev	ctt Leu 25	gtt Val	160
gtt Val	gga Gly	ggt Gly	ccg Pro	ctt Le	ctt Leu	gtg Val	tto Phe	c cta Leu 35	a ato ı Ile	gco Ala	a aa	tac n Tyn	gtg Val	Let	tac Tyr	208

							MBI	15 S	eque	nce :	List	ing.	ST25			
		gct Ala 45														256
aaa Lys	aag Lys 60	aag Lys	ctc Leu	aag Lys	cgg Arg	gag Glu 65	aag Lys	cta Leu	aag Lys	caa Gln	gga Gly 70	gtc Val	cct Pro	gtc Val	cct Pro	304
gga Gly 75		taa	aago	cago	ett a	agct	tect	t ca	actto	gtgc	tco	ette	aag			353
cggt	ttt	tgt t	:cggt	taco	ca aa	attto	acco	ttg	geggg	gttt	tttt	ctto	ct t	tact	tctgt	413
cato	agga	att a	tctt	tgag	gg co	et						٠.				436
<210 <211 <212 <213	> '  >	38 76 PRT Arabi	idops	sis t	:hali	iana										
<400	)> :	38														
Met 1	Asp	Gly	Glu	Asp 5	Phe	Ala	Gly	Lys	Ala 10	Ala	Ala	Glu	Ala	Lув 15	Gly	
Leu	Asn	Pro	Gly 20	Leu	Ile	Val	Leu	Leu 25	Val	Val	Gly	Gly	Pro 30	Leu	Leu	
Val	Phe	Leu 35	Ile	Ala	Asn	Tyr	Val 40	Leu	Tyr	Val	Туг	Ala 45	Gln	Lys	Asn	
Leu	Pro 50	Pro	Arg	Lув	Lys	Lys 55	Pro	Val	Ser	Lys	Lys 60	Lys	Leu	Lys	Arg	
Glu 65	Lys	Leu	Lys	Gln	Gly 70	Val	Pro	Val	Pro	Gly 75	Glu					
<210 <211 <212 <213	.> : !> I	39 L470 DNA Arabi	dops	sis t	hali	lana										
<220 <221 <222 <223	.> ( !> '	CDS (280) 3265	(1	317)	١											
<400		39 ett g	gaag	jccaa	aa to	aaac	cttt	cct	tcaa	tcc	tcaa	atti	tc g	gaaaa	ittttc	60
tctt	ttg	ett t	acgt	tcto	et ca	atto	ttat	ttg	taag	gaaa	gttt	gtto	ct t	taat	caatc	120
aaat	caaa	aga g	jactt	ttga	aa ga	ttgt	ttco	caa	tttg	gegt	caat	cggg	gat o	cgagt	caaat	180
ctga	aato	ett c	tcca	ctca	at Ca	tctg	acta	a taa	gact	taa	tcaa	aggga	act t	tttg	ttcgg	240
gttt	ggtt	tt a	aacg	tctt	g ga	ttcg	aagt	: ggt	taag		let A			aat a Asn <i>A</i>	sn	294
gga Gly	ggt Gly	tca Ser	agc Ser	tca Ser 10	ctt Leu	cca Pro	cct Pro	ttc Phe	ctt Leu 15	act Thr	aaa Lys	aca Thr	tat Tyr	gaa Glu 20	atg Met	342
gtt Val	gat Asp	gat Asp	tct Ser	tct Ser	tct Ser	gac Asp	tcg Ser	gtc Val	gtt Val	gct Ala	tgg Trp	agc Ser	gaa Glu	aac Asn	aac Asn	390

1302

MBI15 Sequence Listing.ST25 aaa agc ttc atc gtc aag aat cca gca gag ttt tca aga gac ctt ctt Lys Ser Phe Ile Val Lys Asn Pro Ala Glu Phe Ser Arg Asp Leu Leu 40 45 50438 ccg aga ttc ttc aag cat aag aat ttc tca agt ttc atc cgt cag ctt Pro Arg Phe Phe Lys His Lys Asn Phe Ser Ser Phe Ile Arg Gln Leu 486 aat aca tat ggt ttt cga aaa gta gat cct gag aaa tgg gaa ttc ttg Asn Thr Tyr Gly Phe Arg Lys Val Asp Pro Glu Lys Trp Glu Phe Leu 70 75 80 85 534 aat gat gat ttt gtt aga ggt cga cct tac ctt atg aag aac att cat Asn Asp Asp Phe Val Arg Gly Arg Pro Tyr Leu Met Lys Asn Ile His 582 aga cga aaa ccg gtt cat agc cac tcg tta gtg aat cta caa gcg caa Arg Arg Lys Pro Val His Ser His Ser Leu Val Asn Leu Gln Ala Gln 630 aat cct ttg acg gaa tca gaa aga cgg agc atg gag gat cag ata gaa Asn Pro Leu Thr Glu Ser Glu Arg Arg Ser Met Glu Asp Gln Ile Glu 678 aga ctg aaa aat gag aaa gaa ggc ctt ctt gcg gag tta cag aac caa Arg Leu Lys Asn Glu Lys Glu Gly Leu Leu Ala Glu Leu Gln Asn Gln 135 726 gag caa gaa cgg aaa gag ttt gag ctg caa gta acg aca ttg aaa gat Glu Gln Glu Arg Lys Glu Phe Glu Leu Gln Val Thr Thr Leu Lys Asp 774 cgg tta caa cat atg gaa caa cat cag aaa tca ata gtg gca tat gtt Arg Leu Gln His Met Glu Gln His Gln Lys Ser Ile Val Ala Tyr Val 170 175 180 822 tca cag gtt ttg gga aaa cca gga ctt tca cta aac ctc gaa aac cat Ser Gln Val Leu Gly Lys Pro Gly Leu Ser Leu Asn Leu Glu Asn His 185 190 195 870 gag aga aga aaa aga aga ttt caa gag aac tct ctt cct cca agc agt Glu Arg Arg Lys Arg Arg Phe Gln Glu Asn Ser Leu Pro Pro Ser Ser 200 205 210 918 tca cac ata gaa cag gtc gaa aag tta gaa tct tcg cta acg ttt tgg Ser His Ile Glu Gln Val Glu Lys Leu Glu Ser Ser Leu Thr Phe Trp 966 gag aat ctt gta tcg gaa tca tgc gag aag agc ggt ttg cag tca tca Glu Asn Leu Val Ser Glu Ser Cys Glu Lys Ser Gly Leu Gln Ser Ser 1014 agc atg gat cat gat gca gct gag tca agt cta agt att ggc gat aca Ser Met Asp His Asp Ala Ala Glu Ser Ser Leu Ser Ile Gly Asp Thr 1062 cga ccc aaa tca tcg aag att gat atg aac tca gag ccg ccc gtt acc Arg Pro Lys Ser Ser Lys Ile Asp Met Asn Ser Glu Pro Pro Val Thr 265 270 275 1110 gtt act gcg cct gct cca aaa aca ggc gtt aac gat gac ttt tgg gaa Val Thr Ala Pro Ala Pro Lys Thr Gly Val Asn Asp Asp Phe Trp Glu 1158 285 caa tgt ttg aca gag aac cct gga tca acc gag caa caa gaa gtt cag Gln Cys Leu Thr Glu Asn Pro Gly Ser Thr Glu Gln Gln Glu Val Gln 295 300 305 1206 tca gag aga aga gat gtc ggt aat gat aat aat ggt aat aag att gga Ser Glu Arg Arg Asp Val Gly Asn Asp Asn Asn Gly Asn Lys Ile Gly 1254

aat caa agg acg tat tgg tgg aat tca ggg aat gta aat aac att aca

MBI15 Sequence Listing.ST25

Asn Gln Arg Thr Tyr Trp Trp Asn Ser Gly Asn Val Asn Asn Ile Thr 330 335 340

gag aaa gct tct tga catgaatgag gtttttgtaa aatagttttc ttttggttcc 1357 Glu Lys Ala Ser

actgagatta ttgtatgtgt tcattattta ttactctgtt tctgtaaaaa caaatctctc 1417 tattgtttga ggcaggagtg acataaatgc atatgcagaa ttggtttcaa aaa 1470

<210> 40

<211> 345

<212> PRT

<213>. Arabidopsis thaliana

<400> 40

Met Asp Glu Asn Asn Gly Gly Ser Ser Ser Leu Pro Pro Phe Leu Thr

Lys Thr Tyr Glu Met Val Asp Asp Ser Ser Ser Asp Ser Val Val Ala 20 25 30

Trp Ser Glu Asn Asn Lys Ser Phe Ile Val Lys Asn Pro Ala Glu Phe 35 40 45

Ser Arg Asp Leu Leu Pro Arg Phe Phe Lys His Lys Asn Phe Ser Ser 50 60

Phe Ile Arg Gln Leu Asn Thr Tyr Gly Phe Arg Lys Val Asp Pro Glu 65 70 75 80

Lys Trp Glu Phe Leu Asn Asp Asp Phe Val Arg Gly Arg Pro Tyr Leu 85 90 95

Met Lys Asn Ile His Arg Arg Lys Pro Val His Ser His Ser Leu Val

Asn Leu Gln Ala Gln Asn Pro Leu Thr Glu Ser Glu Arg Arg Ser Met

Glu Asp Gln Ile Glu Arg Leu Lys Asn Glu Lys Glu Gly Leu Leu Ala 130 135 140

Glu Leu Gln Asn Gln Glu Gln Glu Arg Lys Glu Phe Glu Leu Gln Val 145 150 155 160

Thr Thr Leu Lys Asp Arg Leu Gln His Met Glu Gln His Gln Lys Ser 165 170 175

Ile Val Ala Tyr Val Ser Gln Val Leu Gly Lys Pro Gly Leu Ser Leu 180 185 190

Asn Leu Glu Asn His Glu Arg Arg Lys Arg Arg Phe Gln Glu Asn Ser 195 200 205

Leu Pro Pro Ser Ser Ser His Ile Glu Gln Val Glu Lys Leu Glu Ser 210 215 220

							MBI1	L5 S6	eguer	ice L	ıstı	ng.:	5125					
Ser 225	Leu	Thr	Phe	Trp	Glu 230	Asn	Leu	Val	Ser	Glu   235	Ser	Суз	Glu	Lys	Se 24	r O		
Gly	Leu	Gln	Ser	Ser 245	Ser	Met	Asp	His	Asp 250	Ala .	Ala	Glu	Ser	Ser 255	Le	eu		
Ser	Ile	Gly	Asp 260	Thr	Arg	Pro	Lys	Ser 265	Ser	Lys	Ile	Asp	Met 270	Asn	Se	er		
Glu	Pro	Pro 275	Val	Thr	Val	Thr	Ala 280	Pro	Ala	Pro	Lys	Thr 285	Gly	Val	As	sn		
Asp	Asp 290	Phe	Trp	Glu	Gln	Cys 295	Leu	Thr	Glu	Asn	Pro 300	Gly	Ser	Thr	· G	lu		
Gln 305	Gln	Glu	Val	Gln	Ser 310	Glu	Arg	Arg	Asp	Val 315	Gly	Asn	Asp	Asn	3:	sn 20		
Gly	Asn	Lys	Ile	Gly 325	Asn	Gln	Arg	Thr	Tyr 330	Trp	Trp	Asn	Ser	Gly 335	, A	sn (		
Val	Asn	Asn	11e 340	Thr	Glu	Lys	Ala	Ser 345	•									
<21 <21 <21 <21	1> 2>	41 913 DNA Arab	oidop	sis	thal	iana											•	
<22 <22 <22 <22	1> 2>	CDS (52) G100	) (5 )6	783)														
<40 gat	10> :aaat	41 caa	tca	acaaa	ac a	aaaa	aaac	et ci	tatag	gttag	ttt	.ctc	tgaa	a a M 1		tac Tyr	5'	7
gga Gl	a cag	g tg n Cy 5	c aat	t ata	a gaa a Glu	tco Ser	gad Ası	tao Ty:	c gct r Ala	t ttg a Leu	ttg Lev	gae Gli 15	g tc u Se	g at r Il	a a e :	aca Thr	10	5
cgi	cae Hi:	s Le	g ct u Le	a gga u Gl	a gga y Gly	998 7 Gly 25	a gga y Gl	a ga y Gl	g aa u Asi	c gag n Glu	cto Lev 30	g cg 1 Ar	a ct g Le	caa uAs	t e	gag Glu	15	3
tc: Se: 35	a ac	a cc r Pr	g ag o Se	t to r Se	g tgi r Cy: 40	t tto	c ace	a ga r Gl	g ag u Se	t tgg r Trp 45	g gga 5 Gl _y	a gg y Gl	t tt y Le	g cc u Pr		ttg Leu 50	20	1
aa Ly	a ga s Gl	g aa u As	t ga n As	t tc p Se 55	a gaq r Gl	giga u As	c at p Me	g tt t Le	g gt u Va 60	g tad 1 Ty	c Gly	a ct y Le	c ct u Le	c aa u Ly 65		gat Asp	24	9
gc Al	c tt a Ph	c ca e Hi	t tt .s Ph 70	e As	c ac p Th	g tc r Se	a tc r Se	a to r Se 75	I AS	c tto p Le	g ag u Se	c tg r Cy	t ct s Le 80		t ne	gat Asp	29	7
t t Ph	t cc e Pr	g gg G Al	la Va	t aa 1 Ly	a gt s Va	c ga 1 Gl	g cc u Pr 90	OTI	t ga ir Gl	g aa u As	c tt n Ph	t ac e Th 95		eg af La Mo	tg et	gag Glu	34	15
ga G1	g aa u Ly 10	78 PI	ca aa co Ly	ıg aa /s Ly	a go rs Al	g at a Il	e PI	g gt o Va	t ac	g ga ir Gl	g ac u Th 11		ca gi la Va	tg aa al L	ag ys	gcg Ala	39	93
										Dage	52							

									_			_				
					gtg Val 120											441
					gcg Ala											489
					gat Asp											537
					cgc Arg											585
					ccg Pro											633
tcg Ser 195	tcg Ser	tcg Ser	tcg Ser	tcg Ser	tcc Ser 200	tct Ser	tct Ser	acg Thr	tcg Ser	tcg Ser 205	tct Ser	gaa Glu	aac Asn	G1y 999	aag Lys 210	681
					aaa Lys											729
					ggt Gly											777
tca Ser	taa	gttt	gato	tt g	gtgtg	ttt	g ta	gttg	aata :	gtt	ttgo	tat	aaat	gttg	jag	833
gcad	caaç	jta a	aagt	gtto	e eg	tgat	gtaa	att	agtt	act	aaac	agag	jcc a	itata	tcttc	893
aato	aaaa	aa a	laaaa	aaaa	ıa											913

<210> 42

<211> 243 <212> PRT

<213> Arabidopsis thaliana

<400> 42

Met Tyr Gly Gln Cys Asn Ile Glu Ser Asp Tyr Ala Leu Leu Glu Ser 1 5 15

Ile Thr Arg His Leu Leu Gly Gly Gly Gly Glu Asn Glu Leu Arg Leu 25 30

As nGlu Ser Thr Pro Ser Ser Cys Phe Thr Glu Ser Trp Gly Gly Leu 35 40 45

Pro Leu Lys Glu Asn Asp Ser Glu Asp Met Leu Val Tyr Gly Leu Leu 50 60

Lys Asp Ala Phe His Phe Asp Thr Ser Ser Ser Asp Leu Ser Cys Leu 65 70 70 80

Phe Asp Phe Pro Ala Val Lys Val Glu Pro Thr Glu Asn Phe Thr Ala 85 90 95

Met Glu Glu Lys Pro Lys Lys Ala Ile Pro Val Thr Glu Thr Ala Val

Lys	Ala	Lys 115	His	Tyr	Arg	Gly	Val 120	Arg	Gln	Arg	Pro	Trp 125	Gly	Lys	Phe	
Ala	Ala 130	Glu	Ile	Arg	Asp	Pro 135	Ala	Lys	Asn	Gly	Ala 140	Arg	Val	Trp	Leu	
Gly 145	Thr	Phe	Glu	Thr	Ala 150	Glu	Asp	Ala	Ala	Leu 155	Ala	Tyr	Asp	Ile	Ala 160	
Ala	Phe	Arg	Met	Arg 165	Gly	Ser	Arg	Ala	Leu 170	Leu	Asn	Phe	Pro	Leu 175	Arg	
Val	Asn	Ser	Gly 180	Glu	Pro	Asp	Pro	Val 185	Arg	Ile	Thr	Ser	Lys 190	Arg	Ser	
Ser	Ser	Ser 195	Ser	Ser	Ser	Ser	Ser 200	Ser	Ser	Thr	Ser	Ser 205	Ser	Glu	Asn	
Gly	Lys 210	Leu	Lys	Arg	Arg	Arg 215	Lys	Ala	Glu	Asn	Leu 220	Thr	Ser	Glu	Val	
Val 225	Gln	Val	Lys	Сув	Glu 230	Val	Gly	Asp	Glu	Thr 235	Arg	Val	Asp	Glu	Leu 240	
Leu	Val	Ser														
	1> 2>	43 912 DNA Arab	idop	sis	thal	iana										
	1>	CDS (53) G130	(8 9	59)										٠		
<40 cgt	0> :cgac	43 ctc	ttaa	ttaa	ga c	gact	tgag	ja ga	.gaaa	gaaa	gat	acgt	.gga	ag a	itg acc let Thr	58
aaa Lys	tct Ser	gga Gly 5	gag Glu	aga Arg	cca Pro	aaa Lys	cag Glr	j aga n Arg	cag Gln	agg Arg	aaa Lys	999 Gly	tta Lei	tgg Tr	tca Ser	106
cct Pro	gaa Glu 20	gaa Glu	gac Asp	cag Gln	, aag Lys	ctc Leu 25	aag Lys	g agt s Ser	tto Phe	ato Ile	cto Lev	tct Sei	cgt Arg	g Gly	cat His	154
gct Ala 35	tgo a Cys	tgg Trp	acc Thr	act Thr	gtt Val	ccc Pro	ato Ile	c cta e Leu	gct Ala	gga Gly 45	tto Lev	caa Glr	a agg	g aat g Asi	ggg Gly 50	202
		ta:	aga	a tta	agg	tgg	att	c aat	tac	cta	aga	CC	gga Gl	a cta y Lei	a aag 1 Lys	250
aaa Lys	a ago s Sei	Cys	Arg	55	ı Arç	111	, 110		60	. Let		,		65		
Ly	s Sei	c Cys	a Arg	Lev 55	- gaa	oaa	a gaa	а даа	60 a qaq	aco	ato	tte	act	65 t tta	a cat u His	298

Ser	Ser	Leu 85	Gly	Asn	Lys	Trp		15 S Arg							Gly	
aga Arg	aca Thr 100	gac Asp	aac Asn	gag Glu	att Ile	aag Lys 105	aac Asn	tat Tyr	tgg Trp	cat His	tcc Ser 110	tat Tyr	ctg Leu	aag Lys	aag Lys	394
					caa Gln 120											442
					tca Ser											490
					cac His											538
					tcc Ser											586
					cct Pro											634
					gat Asp 200	Tyr										682
					gat Asp					Glu						730
					tct Ser											778
					cat His											826
					gac Asp					taa	gaag	jagto	gaa t	atga	atcgta	879
agag	gaad	cat a	agct	agtt	a ct	tgtg	ttac	ago	2							912
<210 <211 <212 <213	l> 2 ?> I	14 268 PRT Arabi	idops	sis t	:hali	.ana										
<400	)> 4	4														,
Met 1	Thr	Lys	Ser	Gly 5	Glu	Arg	Pro	Lys	Gln 10	Arg	Gln	Arg	Lys	Gly 15	Leu	
,Trp	Ser	Pro	Glu 20	Glu	Asp	Gln	Lув	Leu 25	Lys	Ser	Phe	Ile	Leu 30	Ser	Arg	
Gly	His	Ala 35	Сув	Trp	Thr	Thr	Val 40	Pro	Ile	Leu	Ala	Gly 45	Leu	Gln	Arg	
Asn	Gly 50	Lys	Ser	Сув	Arg	Leu 55	Arg	Trp	Ile	Asn	Tyr 60	Leu	Arg	Pro	Gly	
Leu 65	Lys	Arg	Gly	Ser	Phe 70	Ser	Glu	Glu	Glu	Glu 75	Glu	Thr	Ile	Leu	Thr 80	

Leu	His	Ser	Ser	Leu 85	Gly	Asn	Lys	Trp	Ser 90	Arg	Ile	Ala	Lys	Tyr 95	Leu	
Pro	Gly	Arg	Thr 100	Asp	Asn	Glu	Ile	Lys 105	Asn	Tyr	Trp	His	Ser 110	Tyr	Leu	
Lys	Lys	Arg 115	Trp	Leu	Lys	Ser	Gln 120	Pro	Gln	Leu	Lys	Ser 125	Gln	Ile	Ser	
Asp	Leu 130	Thr	Glu	Ser	Pro	Ser 135	Ser	Leu	Leu	Ser	Сув 140	Gly	Lys	Arg	Asn	
Leu 145	Glu	Thr	Glu	Thr	Leu 150	Asp	His	Val	Ile	Ser 155	Phe	Gln	ГЛа	Phe	Ser 160	
Glu	Asn	Pro	Thr	Ser 165	Ser	Pro	Ser	Lys	Glu 170	Ser	Asn	Asn	Asn	Met 175	Ile	
Met	Asn	Asn	Ser 180	Asn	Asn	Leu	Pro	Lys 185	Leu	Phe	Phe	Ser	Glu 190	Trp	Ile	
Ser	Ser	Ser 195		Pro	His	Ile	Asp 200	Tyr	Ser	Ser	Ala	Phe 205	Thr	Asp	Ser	
Lys	His 210		Asn	Glu	Thr	Gln 215	Asp	Gln	Ile	Asn	Glu 220	Glu	Glu	Val	Met	
Met 225		Asn	Asn	Asn	Asn 230	Tyr	Ser	Ser	Leu	Glu 235	Asp	Val	Met	Leu	Arg 240	
Thr	Asp	Phe	Leu	Gln 245		Asp	His	Glu	Tyr 250	Ala	, Asn	Tyr	Tyr	255		
Gly	Asp	Phe	Phe 260		Asn	Ser	Asp	Gln 265	Asn	Tyr	Val			•	•	
	.1> .2>	45 1575 DNA Arab		sis	thal	iana										
<22	1> 2>	CDS (1). G255	. (15 io	75)												
	00> g gct ; Ala		tat Tyr	tac Tyr 5	cct Pro	aat Asr	agt Ser	gto Val	ggc Gly 10	atg Met	caa Glr	tct Sei	ctt Leu	tac Tyr 15	caa Gln	48
gaa Glu	tco Sei	att Ile	tac Tyr 20	cto Lev	aac Asn	gaa Glu	caa Glr	caa Glr 25	caa Glr	caa Glr	caa Glr	a caa a Gli	caa Glr 30	gct	tct Ser	96
tct Sei	tco Sei	tct Ser 35	gct Ala	gca Ala	tct Ser	tto Phe	tco Ser 40	gag Glu	g att 1 Ile	gtt Val	tco Sei	gg( Gl) 45	gat Asp	gtt Val	cga L Arg	144
aad	c aad	gaç	ato	gta	a ttt	: ato	cca	a cca		a ago Page		c gta	a gco	gto	e aac	192

Asn	Asn 50	Glu	Met	Val	Phe	Ile 55		15 S Pro							Asn	
					tca Ser 70											240
					ggt Gly											288
ccg Pro	ttt Phe	cag Gln	tat Tyr 100	cat His	tac Tyr	cag Gln	aat Asn	ctt Leu 105	tcg Ser	aac Asn	caa Gln	ttg Leu	agt Ser 110	tac Tyr	aat Asn	336
aat Asn	ctt Leu	aat Asn 115	cct Pro	tct Ser	act Thr	atg Met	tct Ser 120	gat Asp	gag Glu	aat Asn	999 999	aag Lys 125	agc Ser	ttg Leu	agt Ser	384
					tct Ser											432
					ggt Gly 150											480
					agt Ser											528
					gat Asp											576
ttg Leu	999 Gly	aat Asn 195	aag Lys	aag Lys	atg Met	aag Lys	aat Asn 200	gat Asp	aaa Lys	ggt Gly	caa Gln	gac Asp 205	ttt Phe	cac His	aat Asn	624
					att Ile											. 672
					cgt Arg 230											720
					gag Glu											768
					gca Ala											816
					aca Thr											864
					gac Asp											912
				Glu	aga Arg 310											960
					tta Leu											1008
					atg Met											1056

Page 57

							MBI	15 Se	equer	ice I	Listi	ing.S	T25	~~~	cat	1104
tta Leu	Pro	gaa Glu 355	aac Asn	Ser	gtc Val	Ser	Ile 360	Leu	Arg	Ala	Trp	Leu 365	Phe	Glu	His	
ttc	ctt	cat	cca	tat	cct	aaa	gaa	tca	gag	aaa	atc	atg	ctt	tca	aag	1152
Phe	Leu 370	His	Pro	Tyr	Pro	Lys 375	Ğlu	Ser	Ğlü	Lys	11e 380	Met	Leu	Ser	Lys	
cag	aca	gga	cta	tcg	aaa	aac	cag	gtt	gca	aat	tgg	ttt	att	aac	gcg	1200
385		Gly			390					395					400	
aga	gtt	cga Arg	cta	tgg	aaa	cca	atg	att	gaa	gag	atg	tat	aaa Lvs	gaa Glu	gag Glu	1248
Arg	Val	Arg	Dea	405	nys	110	nec		410			-1-	_,-	415		
ttt	gga	gaa	tca	gca	gag	tta	ctc	tct	aac	tct	aat	caa	gac	acc	aaa	1296
Phe	ĞĨy	Ğlu	Ser 420	Āla	Glu	Leu	Leu	Ser 425	Asn	Ser	Asn	Gln	Asp 430	Thr	Lys	
		cag		202	+a+	<b>636</b>	ctc	222	cac	naa	gac	tct	tca	tct	tca	1344
aaa Lys	Met	Gln	Glu	Thr	Ser	Gln	Leu	Lys	His	Glu	Asp	ser	Ser	Ser	Ser	
		435					440					445				
caa	caa	cag Gln	aat	cag	gga	aac Asn	aac Asn	aac Asn	aac Asn	aac Asn	atc Ile	cca Pro	tat Tyr	aca Thr	tct Ser	1392
GIII	450	GIII	non	0111	OLY	455					460					
qat	gca	gaa	caa	aac	cta	gtc	ttt	gca	gat	cct	aaa	cca	gac	cgt	gct	1440
Asp 465	Ala	Ğlu	Gln	Asn	Leu 470	Val	Phe	Ala	Asp	Pro 475	Lys	Pro	Asp	Arg	480	
								a <b>t-a</b>	226	tat	cet	000	+++	aat	att	1488
act Thr	act Thr	gga Gly	gat	Tyr	gac Asp	Ser	Leu	Met	Asn	Tyr	His	Gly	Phe	GIA	Ile	
				485				_	490					495		
gat	gat	tac Tyr	aat	cgt	tac	gtt	ggc	ctt	gga	aac	caa	caa	gat	ggc	aga Arg	1536
Asp	Asp	туг	500		ıyı	Val	GIY	505	Gry	ASII	0	<b></b>	510	0-7	*5	
tat	tct	aat	ccc	cat	caa	tta	cac	gac	ttt	gtt	gtc	taa				1575
Tyr	Ser	Asn 515	Pro	His	Gln	Leu	His 520	Asp	Phe	Val	Val					
		313					320									
<21	0>	46														•
<21 <21		524 PRT														
<21		Arab	idop	sis	thal	iana										
<40	0>	46														
Met	Ala	Val	Туг	Tyr	Pro	Asn	Ser	Val	Gly	Met	Gln	Ser	Leu	Tyr	Gln	
1				5					10					15		
<i>α</i> 1	Cor	• т1а	. The same	· T.611	λen	G1 u	G1 n	Glr	Gln	Gln	Gln	Gln	Gln	Ala	Ser	
GIU	Ser	. 116	20	пес	, Mail	GIU		25					30			
														_		
Ser	Ser	Ser	Ala	Ala	Ser	Phe	Ser 40	Glu	Ile	· Val	Ser	Gly	Asp	val	Arg	
		در														
Asn	Asr	ı Glu	ı Met	: Val	Phe	Ile	Pro	Pro	Thr	Ser	Asp	Val	Ala	val	Asn	
	50					55					60					
al.	, n	, 11-7	ጥሎ-	r ttn.T	Ça~	. ga-	· Der	) Dar	) I.e.	ı Ser	Phe	His	Glv	/ Glv	Gly	
65	, ASI	. val	. 4111	. 701	70					75					80	

Leu Ser Leu Ser Leu Gly Asn Gln Ile Gln Ser Ala Val Ser Val Ser 85 90 95

MBI15 Sequence Listing.ST25

Pro Phe Gln Tyr His Tyr Gln Asn Leu Ser Asn Gln Leu Ser Tyr Asn 100 105 110

Asn Leu Asn Pro Ser Thr Met Ser Asp Glu Asn Gly Lys Ser Leu Ser 115 120 125

Val His Gln His His Ser Asp Gln Ile Leu Pro Ser Ser Val Tyr Asn 130 135 140

Asn Asn Gly Asn Asn Gly Val Gly Phe Tyr Asn Asn Tyr Arg Tyr Glu 145 150 155 160

Thr Ser Gly Phe Val Ser Ser Val Leu Arg Ser Arg Tyr Leu Lys Pro 165 170 175

Thr Gln Gln Leu Leu Asp Glu Val Val Ser Val Arg Lys Asp Leu Lys 180 185 190

Leu Gly Asn Lys Lys Met Lys Asn Asp Lys Gly Gln Asp Phe His Asn 195 200 205

Gly Ser Ser Asp Asn Ile Thr Glu Asp Asp Lys Ser Gln Ser Gln Glu 210 215 220

Leu Ser Pro Ser Glu Arg Gln Glu Leu Gln Ser Lys Lys Ser Lys Leu 225 230 235 240

Leu Thr Met Val Asp Glu Val Asp Lys Arg Tyr Asn Gln Tyr His His 245 250 255

Gln Met Glu Ala Leu Ala Ser Ser Phe Glu Met Val Thr Gly Leu Gly 260 265 270

Ala Ala Lys Pro Tyr Thr Ser Val Ala Leu Asn Arg Ile Ser Arg His 275 280 285

Phe Arg Cys Leu Arg Asp Ala Ile Lys Glu Gln Ile Gln Val Ile Arg 290 295 300

Gly Lys Leu Gly Glu Arg Glu Thr Ser Asp Glu Gln Gly Glu Arg Ile 305 310 315 320

Pro Arg Leu Arg Tyr Leu Asp Gln Arg Leu Arg Gln Gln Arg Ala Leu 325 330 335

His Gln Gln Leu Gly Met Val Arg Pro Ala Trp Arg Pro Gln Arg Gly 340 345 350

Leu Pro Glu Asn Ser Val Ser Ile Leu Arg Ala Trp Leu Phe Glu His

Phe Leu His Pro Tyr Pro Lys Glu Ser Glu Lys Ile Met Leu Ser Lys 370 375 380

Gln Thr Gly Leu Ser Lys Asn Gln Val Ala Asn Trp Phe Ile Asn Ala 385 390 395 400

Arg Val Arg Leu Trp Lys Pro Met Ile Glu Glu Met Tyr Lys Glu Glu 405 410 415	
Phe Gly Glu Ser Ala Glu Leu Leu Ser Asn Ser Asn Gln Asp Thr Lys 420 425 430	
Lys Met Gln Glu Thr Ser Gln Leu Lys His Glu Asp Ser Ser Ser Ser 445	
Gln Gln Asn Gln Gly Asn Asn Asn Asn Ile Pro Tyr Thr Ser 450 455 460	
Asp Ala Glu Gln Asn Leu Val Phe Ala Asp Pro Lys Pro Asp Arg Ala 465 470 475 480	
Thr Thr Gly Asp Tyr Asp Ser Leu Met Asn Tyr His Gly Phe Gly Ile 485 490 495	
Asp Asp Tyr Asn Arg Tyr Val Gly Leu Gly Asn Gln Gln Asp Gly Arg 500 505 510	
Tyr Ser Asn Pro His Gln Leu His Asp Phe Val Val 515 520	
<210> 47 <211> 1983 <212> DNA <213> Arabidopsis thaliana	
<220> <221> CDS <222> (73)(1956) <223> G965	
<400> 47 gattctctgt gtatgtctga atccttacag gatccaagag ctttggaaaa aagatataat	60
gaataacaag at atg ggt tta gct act aca act tct tct atg tca caa gat Met Gly Leu Ala Thr Thr Thr Ser Ser Met Ser Gln Asp 1 5 10	111
tat cat cat cac caa gga atc ttt tcc ttc tct aat gga ttc cac cga Tyr His His His Gln Gly Ile Phe Ser Phe Ser Asn Gly Phe His Arg 15 20 25	159
tca tca tca acc act cat cag gag gaa gta gat gaa tcc gcc gtc gtc Ser Ser Ser Thr Thr His Gln Glu Glu Val Asp Glu Ser Ala Val Val 30 35 40 45	207
tcc ggt gct caa att ccg gtt tat gaa acc gcc gga atg ttg tct gaa Ser Gly Ala Gln Ile Pro Val Tyr Glu Thr Ala Gly Met Leu Ser Glu 50 55 60	25 <b>5</b>
atg ttt gct tac cct ggc gga ggt ggc ggc ggt tcc ggt gga gag att Met Phe Ala Tyr Pro Gly Gly Gly Gly Gly Ser Gly Gly Glu Ile 65 70 75	303
ctt gat cag tct act aaa cag ttg cta gag caa caa aac cgt cac aac	
Leu Asp Gln Ser Thr Lys Gln Leu Leu Glu Gln Gln Asn Arg His Asn 80 85 90  aac aac aat aac tca act ctt cat atg tta tta cca aat cat cat caa	351 399

## MBI15 Sequence Listing.ST25

									•	_						
					gac Asp 115											447
					tct Ser											495
					cac His											543
					gcc Ala											591
					gat Asp											639
					aat Asn 195											687
					cac His											735
					gcg Ala											783
					tat Tyr											831
					aga Arg											879
					aat Asn 275											927
					aca Thr											975
gat Asp	cgg Arg	att Ile	gaa Glu 305	cat His	caa Gln	aga Arg	aga Arg	aaa Lys 310	gtc Val	aag Lys	cta Leu	cta Leu	tct Ser 315	atg Met	ctt Leu	1023
gaa Glu	g <u>ag</u> Glu	gtg Val 320	gac Asp	cga Arg	cgg Arg	Tyr	aac Asn 325	His	tac Tyr	tgc Cys	Glu	caa Gln 330	atg Met	caa Gln	atg Met	1071
					gac Asp											1119
					caa Gln 355											1167
					gtt Val											1215
gat Asp	aaa Lys	gag Glu	gcg Ala 385	gca Ala	GJ Y 999	gct Ala	gca Ala	tcc Ser 390	tcg Ser	G1y 999	tta Leu	acc Thr	aaa Lys 395	G1y 999	gaa Glu	1263
					ttg Leu											1311

		400					MBI1 405	15 Se	equen	ice I	List	ing.9	ST25			
ttt Phe	cat His 415	cat His	atg Met	ggt Gly	atg Met	atg Met 420	gag Glu	caa Gln	gag Glu	gca Ala	tgg Trp 425	aga Arg	ccg Pro	caa Gln	cgt Arg	1359
ggt Gly 430	ttg Leu	cct Pro	gaa Glu	cgc Arg	tcc Ser 435	gtt Val	aat Asn	atc Ile	ctt Leu	aga Arg 440	gct Ala	tgg Trp	cta Leu	ttc Phe	gag Glu 445	1407
cat His	ttt Phe	ctt Leu	aat Asn	ccg Pro 450	tac Tyr	cca Pro	agc Ser	gat Asp	gct Ala 455 _.	gat Asp	aag Lys	cac His	ctc Leu	tta Leu 460	gca Ala	1455
cga Arg	cag Gln	act Thr	ggt Gly 465	tta Leu	tcc Ser	aga Arg	aat Asn	cag Gln 470	gtg Val	tca Ser	aat Asn	tgg Trp	ttc Phe 475	ata Ile	aat Asn	1503
gct Ala	agg Arg	gtt Val 480	cgc Arg	cta Leu	tgg Trp	aaa Lys	cca Pro 485	atg Met	gtg Val	gaa Glu	gag Glu	atg Met 490	tat Tyr	caa Gln	caa Gln	1551
gaa Glu	gca Ala 495	aaa Lys	gaa Glu	aga Arg	gaa Glu	gaa Glu 500	gca Ala	gaa Glu	gaa Glu	gaa Glu	aat Asn 505	GIU	aat Asn	caa Gln	caa Gln	1599
caa Gln 510	caa Gln	aga Arg	aga Arg	cag Gln	caa Gln 515	caa Gln	aca Thr	aac Asn	aac Asn	aac Asn 520	gac Asp	acg Thr	aaa Lys	ccc Pro	aac Asn 525	1647
aac Asn	aat Asn	gaa Glu	aac Asn	aac Asn 530	ttc Phe	act Thr	gtc Val	ata Ile	acc Thr 535	gca Ala	caa Gln	act Thr	cca Pro	acg Thr 540	1111	1695
atg Met	aca Thr	tcg Ser	aca Thr 545	His	cac His	gaa Glu	aac Asn	gac Asp 550	Ser	tca Ser	t t c Phe	ctc Leu	ser Ser 555	502	gtc	1743
gcc Ala	gcc	gct Ala 560	Ser	cac His	ggc	ggt	tca Ser 565	Ast	gcg Ala	ttc Phe	acc Thr	gtc Val 570	, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	acg Thr	tgt Cys	1791
cag Gln	caa Gln 575	Asp	gto Val	agt Ser	gac Asp	tto Phe 580	His	gto Val	gac Asp	gga Gly	gat Asp 585	נוט נ	gtg Val	aac Asr	gtc Val	1839
ata Ile 590	Arg	tto Phe	ggg Gly	acc Thr	aaa Lys 595	Gli	act Thr	ggt Gl	gac Asp	gtg Val 600	. 561	ctt Lei	acg Thr	ctt Lev	ggt Gly 605	1887
cta Lei	cgo Arg	cac His	tct Sei	gg Gly 610	Asr	att 1116	cct Pro	gat Asp	aag Lys 615	ASI	act Thi	t tot	tto Phe	tco Ser 620	gtt Val	1935
aga	gad J Asj	ttt Phe	gga Gly 625	y As <u>r</u>	ttt Phe	tag	g tct	tct	tgt	ttct	caai	ttt a	attca	atc		1983
<2: <2:	10> 11> 12> 13>	48 627 PRT Aral	oido	psis	tha:	lian	a					•				
	00>	48														
Me 1	t Gl	y Le	u Al	a Th	r Th	r Th	r Se	r Se	r Me	t Se	r Gl	n As	р Ту	r Hi 15	s His	
Hi	s Gl	n Gl	y Il 20		e Se	r Ph	e Se	r As 25	n Gl	y Ph	e Hi	s Ar	g Se 30	r Se	r Ser	

Thr Thr His Gln Glu Glu Val Asp Glu Ser Ala Val Val Ser Gly Ala

35

Gln Ile Pro Val Tyr Glu Thr Ala Gly Met Leu Ser Glu Met Phe Ala 50 55 60

Tyr Pro Gly Gly Gly Gly Gly Gly Ser Gly Gly Glu Ile Leu Asp Gln 65 70 75 80

Ser Thr Lys Gln Leu Leu Glu Gln Gln Asn Arg His Asn Asn Asn Asn Asn 90 95

Asn Ser Thr Leu His Met Leu Leu Pro Asn His His Gln Gly Phe Ala

Phe Thr Asp Glu Asn Thr Met Gln Pro Gln Gln Gln His Phe Thr 115 120 125

Trp Pro Ser Ser Ser Ser Asp His His Gln Asn Arg Asp Met Ile Gly 130 135 140

Thr Val His Val Glu Gly Gly Lys Gly Leu Ser Leu Ser Leu Ser Ser 145 150 160

Ser Leu Ala Ala Ala Lys Ala Glu Glu Tyr Arg Ser Ile Tyr Cys Ala 165 170 175

Ala Val Asp Gly Thr Ser Ser Ser Ser Asn Ala Ser Ala His His His 180 185 190

Gln Phe Asn Gln Phe Lys Asn Leu Leu Leu Glu Asn Ser Ser Gln 195 200 205

His His His Gln Val Val Gly His Phe Gly Ser Ser Ser Ser Ser 210 215 220

Pro Met Ala Ala Ser Ser Ser Ile Gly Gly Ile Tyr Thr Leu Arg Asn 225 230 235 240

Ser Lys Tyr Thr Lys Pro Ala Gln Glu Leu Leu Glu Glu Phe Cys Ser 245 250 255

Val Gly Arg Gly His Phe Lys Lys Asn Lys Leu Ser Arg Asn Asn Ser 260 265 270

Asn Pro Asn Thr Thr Gly Gly Gly Gly Gly Gly Gly Ser Ser Ser Ser 275 280 285

Ala Gly Thr Ala Asn Asp Ser Pro Pro Leu Ser Pro Ala Asp Arg Ile 290 295 300

Glu His Gln Arg Arg Lys Val Lys Leu Leu Ser Met Leu Glu Glu Val 305 310 315 320

Asp Arg Arg Tyr Asn His Tyr Cys Glu Gln Met Gln Met Val Val Asn 325 330 335

- MBI15 Sequence Listing.ST25 Ser Phe Asp Gln Val Met Gly Tyr Gly Ala Ala Val Pro Tyr Thr Thr 340 345 350
- Leu Ala Gln Lys Ala Met Ser Arg His Phe Arg Cys Leu Lys Asp Ala 355 360 365
- Val Ala Val Gln Leu Lys Arg Ser Cys Glu Leu Leu Gly Asp Lys Glu 370 375 380
- Ala Ala Gly Ala Ala Ser Ser Gly Leu Thr Lys Gly Glu Thr Pro Arg 385 390 395 400
- Leu Arg Leu Leu Glu Gln Ser Leu Arg Gln Gln Arg Ala Phe His His 405 410 415
- Met Gly Met Met Glu Glu Glu Ala Trp Arg Pro Gln Arg Gly Leu Pro 420 425 430
- Glu Arg Ser Val Asn Ile Leu Arg Ala Trp Leu Phe Glu His Phe Leu 435 440 445
- Asn Pro Tyr Pro Ser Asp Ala Asp Lys His Leu Leu Ala Arg Gln Thr 450 455 460
- Gly Leu Ser Arg Asn Gln Val Ser Asn Trp Phe Ile Asn Ala Arg Val 465 470 475 480
- Arg Leu Trp Lys Pro Met Val Glu Glu Met Tyr Gln Gln Glu Ala Lys 485 490 495
- Glu Arg Glu Glu Ala Glu Glu Glu Asn Glu Asn Gln Gln Gln Arg 500 505 510
- Arg Gln Gln Gln Thr Asn Asn Asn Asp Thr Lys Pro Asn Asn Asn Glu 515 520 525
- Asn Asn Phe Thr Val Ile Thr Ala Gln Thr Pro Thr Thr Met Thr Ser 530 535
- Thr His His Glu Asn Asp Ser Ser Phe Leu Ser Ser Val Ala Ala Ala 545 550 560
- Ser His Gly Gly Ser Asp Ala Phe Thr Val Ala Thr Cys Gln Gln Asp 565 570 575
- Val Ser Asp Phe His Val Asp Gly Asp Gly Val Asn Val Ile Arg Phe 580 585 590
- Gly Thr Lys Gln Thr Gly Asp Val Ser Leu Thr Leu Gly Leu Arg His 595 600 605
- Ser Gly Asn Ile Pro Asp Lys Asn Thr Ser Phe Ser Val Arg Asp Phe 610 615 620

Gly Asp Phe

<210> 49

<21		1420			:											
<21:		ONA Arab:	idops	sis (	thal	Lana										
<220 <220 <220 <220	l> ( 2>	CDS (138) 3793	) (:	1046)	)											
<400		49 tca (	cctaa	acaa	ct co	cacae	caaga	a tto	ette	gcat	ggaa	aagt	tgt 1	totto	cgactt	60
ctc	ttct	cta a	actc	gctal	tc ti	tta	actca	a cc	cagc	cca	ctg	agtc	gaa a	aatti	caaac	120
ctt	tact	cgt 1	ttcc	1	atg g Met <i>l</i> 1								His A			170
atc Ile	tcc Ser	gat Asp	cca Pro 15	tct Ser	cct Pro	acc Thr	gac Asp	gat Asp 20	ttc Phe	ttc Phe	gag Glu	cag Gln	atc Ile 25	ctc Leu	G1y 999	218
					ggt Gly											<b>266</b>
					ccg Pro											314
					atg Met 65											362
					ttt Phe											410
gga Gly	cat His	ggc Gly	ttt Phe 95	ctt Leu	aaa Lys	cct Pro	gat Asp	gaa Glu 100	act Thr	ggt Gly	aaa Lys	cgt Arg	ttc Phe 105	caa Gln	gac Asp	458
					cga Arg											506
					cca Pro											554
att Ile 140	cgg Arg	cct Pro	aga Arg	gtt Val	agg Arg 145	gct Ala	agg Arg	cga Arg	ggt Gly	caa Gln 150	gct Ala	acc Thr	gat Asp	cca Pro	cat His 155	602
agc Ser	atc Ile	gct Ala	gag Glu	agg Arg 160	ctc Leu	cga Arg	agg Arg	Glu	aga Arg 165	ata Ile	gca Ala	gaa Glu	cgg Arg	atc Ile 170	agg Arg	650
tcg Ser	ttg Leu	cag Gln	gaa Glu 175	ctt Leu	gta Val	cct Pro	acc Thr	gtt Val 180	aac Asn	aag Lys	aca Thr	gat Asp	agg Arg 185	gct Ala	gct Ala	698
					gtc Val											746
aag Lув	gtc Val 205	ctg Leu	agc Ser	atg Met	agc Ser	cgt Arg 210	ctt Leu	ggt Gly	gga Gly	gcc Ala	ggt Gly 215	gct Ala	gtc Val	gca Ala	cca Pro	794
					cca Pro				Ser		Glu					842

MBI15 Sequence Listing.ST25	
220 225 230 235	200
gcc gtg tgg gag aaa tgg tca aac gat ggg aca gag agg caa gtg gct Ala Val Trp Glu Lys Trp Ser Asn Asp Gly Thr Glu Arg Gln Val Ala 240 245 250	890
aag ctg atg gaa gaa aac gtt gga gca gcg atg caa ctt ttg caa tca Lys Leu Met Glu Glu Asn Val Gly Ala Ala Met Gln Leu Leu Gln Ser 255 260 265	938
aag gct ctt tgc ata atg ccg atc tca ttg gca atg gcg att tac cat Lys Ala Leu Cys Ile Met Pro Ile Ser Leu Ala Met Ala Ile Tyr His 270 275 280	986
tct cag cca cca gac aca tct tct tca atc gtc aaa cca gag atg aat Ser Gln Pro Pro Asp Thr Ser Ser Ser Ile Val Lys Pro Glu Met Asn 285 290 295	1034
cct cca ccg tag attitigtic atccaacggt ccccagctga tgattgacat Pro Pro Pro 300	1086
tttgctctgt ttcccactac tagacttttg tgactcatga aaggtaagta aaaaggcatt	1146
ggagatggaa totaagtagg atttgtgcag taaagaagta aaacgggato tgtcaaaaga	1206
aggaaaaagc tctcgcttgc ttggctagta tttatcattt tgatgaaagt aactcttttt	1266
tgttcaaaga ctttagtgtg attttcagga ccaagggctt tgagggtagt gctagctgta	1326
gtaatagtaa tgaaggtgtg ggatcgtgtt tttgaattat gtaaaaaagg aagaaaaaac	1386
aaatgttggt attatattat ggttttgcct gaaa	1420
<210> 50 <211> 302 <212> PRT <213> Arabidopsis thaliana	
<400> 50	
Met Ala Asn Asn Asn Ile Pro His Asp Ser Ile Ser Asp Pro Ser 1 10 15	
Pro Thr Asp Asp Phe Phe Glu Gln Ile Leu Gly Leu Ser Asn Phe Ser 20 25 30	
Gly Ser Ser Gly Ser Gly Leu Ser Gly Ile Gly Gly Val Gly Pro Pro 35 40 45	
Pro Met Met Leu Gln Leu Gly Ser Gly Asn Glu Gly Asn His Asn His 50 55 60	
Met Gly Ala Ile Gly Gly Gly Gly Pro Val Gly Phe His Asn Gln Met 65 70 75 80	
Phe Pro Leu Gly Leu Ser Leu Asp Gln Gly Lys Gly His Gly Phe Leu 85 90 95	
Lys Pro Asp Glu Thr Gly Lys Arg Phe Gln Asp Asp Val Leu Asp Asn 100 105 110	
Arg Cys Ser Ser Met Lys Pro Ile Phe His Gly Gln Pro Met Ser Gln 115 120 125	
Pro Ala Pro Pro Met Pro His Gln Gln Ser Thr Ile Arg Pro Arg Val	

MBI15 Sequence Listing.ST25

130 Arg Ala Arg Arg Gly Gln Ala Thr Asp Pro His Ser Ile Ala Glu Arg 150 155 Leu Arg Arg Glu Arg Ile Ala Glu Arg Ile Arg Ser Leu Gln Glu Leu Val Pro Thr Val Asn Lys Thr Asp Arg Ala Ala Met Ile Asp Glu Ile 180 185 190 Val Asp Tyr Val Lys Phe Leu Arg Leu Gln Val Lys Val Leu Ser Met 195 200 205 Ser Arg Leu Gly Gly Ala Gly Ala Val Ala Pro Leu Val Thr Glu Met 210 215 220 Pro Leu Ser Ser Ser Val Glu Asp Glu Thr Gln Ala Val Trp Glu Lys Trp Ser Asn Asp Gly Thr Glu Arg Gln Val Ala Lys Leu Met Glu Glu 245 250 255 Asn Val Gly Ala Ala Met Gln Leu Leu Gln Ser Lys Ala Leu Cys Ile Met Pro Ile Ser Leu Ala Met Ala Ile Tyr His Ser Gln Pro Pro Asp Thr Ser Ser Ser Ile Val Lys Pro Glu Met Asn Pro Pro Pro 295 <211> 1198 <212> DNA <213> Arabidopsis thaliana <220> CDS <221> (96)..(1052) <222> <223> G764 <400> 51 atcgaattcg cggccgctcg atatctttac aaccattaaa caaaaaattt ggccactaca agttgaaaaa gttttgatta tatctaatcg ctgaa atg gat tac aag gta tca Met Asp Tyr Lys Val Ser 113 161 gat tta cca cct ggt ttc aga ttt cac cca act gat gaa gaa ctt ata Asp Leu Pro Pro Gly Phe Arg Phe His Pro Thr Asp Glu Glu Leu Ile 25 30 35 209 aca cac tat cta aga cca aag gtt gta aac tct ttt ttc tct gct ata
Thr His Tyr Leu Arg Pro Lys Val Val Asn Ser Phe Phe Ser Ala Ile
40 45 50 257 gct att ggt gaa gtt gat ctc aac aaa gtc gag cct tgg gac ttg cct Ala Ile Gly Glu Val Asp Leu Asn Lys Val Glu Pro Trp Asp Leu Pro 305

55	MBI	I15 Sequence L 65	isting.ST25	70
tgg aag gct aag ctt Trp Lys Ala Lys Leu 75	ggg gaa aaa Gly Glu Lys	a gag tgg tac s Glu Trp Tyr 80 ·	ttc ttt tgc gta Phe Phe Cys Val 85	aga 353 Arg
gac cga aaa tac ccg Asp Arg Lys Tyr Pro 90	act ggt tta Thr Gly Leu	a aga acg aat u Arg Thr Asn 95	cgt gct act aaa Arg Ala Thr Lys 100	gcc 401 Ala
ggt tat tgg aaa gct Gly Tyr Trp Lys Ala 105	aca ggg aaa Thr Gly Lys 110	B Asp Lys Glu	atc ttc aaa ggg Ile Phe Lys Gly 115	aaa 449 Lys
tct ctt gtt ggt atg Ser Leu Val Gly Met 120	aag aaa aca Lys Lys Thr 125	r Leu Val Phe	tac aaa gga aga Tyr Lys Gly Arg 130	gct 497 Ala
cct aaa gga gta aaa Pro Lys Gly Val Lys 135	aca aat tgg Thr Asn Trp 140	g gtc atg cat p Val Met His 145	gag tat cga tta Glu Tyr Arg Leu	gaa 545 Glu 150
ggc aaa ttc gct atc Gly Lys Phe Ala Ile 155	Asp Asn Leu	c tct aaa acc u Ser Lys Thr 160	gct aag aac gaa Ala Lys Asn Glu 165	tgt 593 Cys
gtt att agt cgt gtt Val Ile Ser Arg Val 170	ttt cat aca Phe His Thr	a cgg act gat r Arg Thr Asp 175	ggt acg aag gag Gly Thr Lys Glu 180	cat 641 His
atg tcc gtt ggt tta Met Ser Val Gly Leu 185	cct ccg ctg Pro Pro Leu 190	u Met Asp Ser	tct cca tat cta Ser Pro Tyr Leu 195	aag 689 Lys
agt aga gga caa gac Ser Arg Gly Gln Asp 200	tct tta gcc Ser Leu Ala 205	c ggg acc acc a Gly Thr Thr	ctt ggt ggg ttg Leu Gly Gly Leu 210	ttg 737 Leu
tct cac gtt acc tac Ser His Val Thr Tyr 215	ttc tcc gac Phe Ser Asp 220	c caa aca acc p Gln Thr Thr 225	gat gac aag agt Asp Asp Lys Ser	ctt 785 Leu 230
gtg gcc gat ttt aaa Val Ala Asp Phe Lys 235	Thr Thr Met	g ttt ggt tcc et Phe Gly Ser 240	gga tcg act aad Gly Ser Thr Ass 245	Pne
tta cca aac ata ggt Leu Pro Asn Ile Gly 250	tct cta cta Ser Leu Leu	a gac ttc gat au Asp Phe Asp 255	cct ctg ttt cta Pro Leu Phe Leu 260	caa 881 Gln
aac aat tot toa gta Asn Asn Ser Ser Val 265	a ctg aag atg Leu Lys Met 270	et Leu Leu Asp	aat gaa gaa acc Asn Glu Glu Thi 275	caa 929
ttt aag aag aat ctt Phe Lys Lys Asn Lev 280	cac aat to His Asn Ser 285	ca ggt tca tca er Gly Ser Ser	gag agt gaa cta Glu Ser Glu Lev 290	aca 977 Thr
gcg agt tct tgg caa Ala Ser Ser Trp Gli 295	a ggt cac aat n Gly His Ass 300	at tot tat ggt on Ser Tyr Gly 305	ser thr Giv Pro	1 gtg 1025 5 Val 310
aat ctt gat tgc gt Asn Leu Asp Cys Val 31!	l Trp Lys Pho	tc tga atttgga he	aaa tcgaaaattt	1072
ggatgttaac taggggg	tat atagggtt	ttt taaaaacagt	gtatatatgc gtt	atgtgtt 1132
agctttagat tctagga	tat acaaagat	tga cactaataga	ttcttataac att	ttgtaaa 1192
aaaaaa				1198

<210> 52 <211> 318

MBI15 Sequence Listing.ST25

<212> PRT

<213> Arabidopsis thaliana

<400> 52

Met Asp Tyr Lys Val Ser Arg Ser Gly Glu Ile Val Glu Gly Glu Val 1 5 10 15

Glu Asp Ser Glu Lys Ile Asp Leu Pro Pro Gly Phe Arg Phe His Pro 25 30

Thr Asp Glu Glu Leu Ile Thr His Tyr Leu Arg Pro Lys Val Val Asn 35 40 45

Ser Phe Phe Ser Ala Ile Ala Ile Gly Glu Val Asp Leu Asn Lys Val 50 60

Glu Pro Trp Asp Leu Pro Trp Lys Ala Lys Leu Gly Glu Lys Glu Trp 65 70 75 80

Tyr Phe Phe Cys Val Arg Asp Arg Lys Tyr Pro Thr Gly Leu Arg Thr 85 90 95

Asn Arg Ala Thr Lys Ala Gly Tyr Trp Lys Ala Thr Gly Lys Asp Lys

Glu Ile Phe Lys Gly Lys Ser Leu Val Gly Met Lys Lys Thr Leu Val 115 120 125

Phe Tyr Lys Gly Arg Ala Pro Lys Gly Val Lys Thr Asn Trp Val Met 130 135 140

His Glu Tyr Arg Leu Glu Gly Lys Phe Ala Ile Asp Asn Leu Ser Lys 145 150 160

Thr Ala Lys Asn Glu Cys Val Ile Ser Arg Val Phe His Thr Arg Thr 165 170 175

Asp Gly Thr Lys Glu His Met Ser Val Gly Leu Pro Pro Leu Met Asp 180 185 190

Ser Ser Pro Tyr Leu Lys Ser Arg Gly Gln Asp Ser Leu Ala Gly Thr 195 200 205

Thr Leu Gly Gly Leu Leu Ser His Val Thr Tyr Phe Ser Asp Gln Thr 210 215 220

Thr Asp Asp Lys Ser Leu Val Ala Asp Phe Lys Thr Thr Met Phe Gly 225 230 235 240

Ser Gly Ser Thr Asn Phe Leu Pro Asn Ile Gly Ser Leu Leu Asp Phe 245 250 255

Asp Pro Leu Phe Leu Gln Asn Asn Ser Ser Val Leu Lys Met Leu Leu

Asp Asn Glu Glu Thr Gln Phe Lys Lys Asn Leu His Asn Ser Gly Ser 275 280 285

Ser Glu Ser Glu Leu Thr Ala Ser Ser Trp Gln Gly His Asn Ser Tyr 290 295 300	
Gly Ser Thr Gly Pro Val Asn Leu Asp Cys Val Trp Lys Phe 305 310 315	
<210> 53 <211> 932 <212> DNA <213> Arabidopsis thaliana	
<220> <221> CDS <222> (43)(759) <223> G350	
<400> 53 ctttcttctc aatttagaac ttagtagcta gtcttcaaga ta atg gca ctt gaa Met Ala Leu Glu 1	54
act ctt act tct cca aga tta tct tct ccg atg ccg act ctg ttt caa Thr Leu Thr Ser Pro Arg Leu Ser Ser Pro Met Pro Thr Leu Phe Gln 5 10 20	102
gat tca gca cta ggg ttt cat gga agc aaa ggc aaa cga tct aag cga Asp Ser Ala Leu Gly Phe His Gly Ser Lys Gly Lys Arg Ser Lys Arg 25 30 35	150
tca aga tct gaa ttc gac cgt cag agt ctc acg gag gat gaa tat atc Ser Arg Ser Glu Phe Asp Arg Gln Ser Leu Thr Glu Asp Glu Tyr Ile 40 45 50	198
gct tta tgt ctc atg ctt ctt gct cgc gac gga gat aga aac cgt gac Ala Leu Cys Leu Met Leu Leu Ala Arg Asp Gly Asp Arg Asn Arg Asp 55 60 65	246
ctt gac ctg cct tct tct tcg tct tca cct cct ctg ctt cct cct ctt Leu Asp Leu Pro Ser Ser Ser Ser Pro Pro Leu Leu Pro Pro Leu 70 75 80	. 294
cct act ccg atc tac aag tgt agc gtc tgt gac aag gcg ttt tcg tct Pro Thr Pro Ile Tyr Lys Cys Ser Val Cys Asp Lys Ala Phe Ser Ser 85 90 95 100	342
tac cag gct ctt ggt gga cac aag gca agt cac cgg aaa agc ttt tcg Tyr Gln Ala Leu Gly Gly His Lys Ala Ser His Arg Lys Ser Phe Ser 105 110 115	390
ctt act caa tct gcc gga gga gat gag ctg tcg aca tcg tcg gcg ata Leu Thr Gln Ser Ala Gly Gly Asp Glu Leu Ser Thr Ser Ser Ala Ile 120 125 130	438
acc acg tct ggt ata tcc ggt ggc ggg gga gga agt gtg aag tcg cac Thr Thr Ser Gly Ile Ser Gly Gly Gly Gly Gly Ser Val Lys Ser His 135 140 145	486
gtt tgc tct atc tgt cat aaa tcg ttc gcc acc ggt caa gct ctc ggc Val Cys Ser Ile Cys His Lys Ser Phe Ala Thr Gly Gln Ala Leu Gly 150 160	534
ggc cac aaa cgg tgc cac tac gaa gga aag aac gga ggc ggt gtg agt Gly His Lys Arg Cys His Tyr Glu Gly Lys Asn Gly Gly Gly Val Ser 165 170 175 180	582
agt agc gtg tcg aat tct gaa gat gtg ggg tct aca agc cac gtc agc Ser Ser Val Ser Asn Ser Glu Asp Val Gly Ser Thr Ser His Val Ser 185 190 195	630

agt ggc cac cgt ggg ttt gac ctc aac ata ccg ccg ata ccg gaa ttc

MBI15 Sequence Listing.ST25 Ser Gly His Arg Gly Phe Asp Leu Asn Ile Pro Pro Ile Pro Glu Phe 200 205 210	
tcg atg gtc aac gga gac gaa gag gtg atg agt cct atg ccg gcg aag Ser Met Val Asn Gly Asp Glu Glu Val Met Ser Pro Met Pro Ala Lys 215 220 225	726
aaa ctc cgg ttt gac ttc ccg gag aaa ccc taa acataaacct aggaaaaact Lys Leu Arg Phe Asp Phe Pro Glu Lys Pro 230 235	779
ttacagaatt cattttatag gaaattgttt tactgtatat acaaatatcg attttgattg	839
atgttcttct tcactgaaaa attatgattc tttgttgtat aattgatgtt tctgaaaaag	899
atataacttt ttattaaaaa aaaaaaaaaa aaa	932
<210> 54 <211> 238 <212> PRT <213> Arabidopsis thaliana	
<400> 54	
Met Ala Leu Glu Thr Leu Thr Ser Pro Arg Leu Ser Ser Pro Met Pro 1 10 15	
Thr Leu Phe Gln Asp Ser Ala Leu Gly Phe His Gly Ser Lys Gly Lys 20 25 30	
Arg Ser Lys Arg Ser Arg Ser Glu Phe Asp Arg Gln Ser Leu Thr Glu 35 40 45	
Asp Glu Tyr Ile Ala Leu Cys Leu Met Leu Leu Ala Arg Asp Gly Asp 50 55 60	
Arg Asn Arg Asp Leu Asp Leu Pro Ser Ser Ser Ser Pro Pro Leu 65 70 75 80	
Leu Pro Pro Leu Pro Thr Pro Ile Tyr Lys Cys Ser Val Cys Asp Lys	
Ala Phe Ser Ser Tyr Gln Ala Leu Gly Gly His Lys Ala Ser His Arg 100 105 110	
Lys Ser Phe Ser Leu Thr Gln Ser Ala Gly Gly Asp Glu Leu Ser Thr 115 120 125	
Ser Ser Ala Ile Thr Thr Ser Gly Ile Ser Gly Gly Gly Gly Gly Ser	
Val Lys Ser His Val Cys Ser Ile Cys His Lys Ser Phe Ala Thr Gly 145 150 160	•
Gln Ala Leu Gly Gly His Lys Arg Cys His Tyr Glu Gly Lys Asn Gly 165 170 175	
Gly Gly Val Ser Ser Ser Val Ser Asn Ser Glu Asp Val Gly Ser Thr	
Ser His Val Ser Ser Gly His Arg Gly Phe Asp Leu Asn Ile Pro Pro 195 200 205	
Page 71	

Ile Pro Glu Phe Ser Met Val Asn Gly Asp Glu Glu Val Met Ser Pro 210 215 220

Met Pro Ala Lys Lys Leu Arg Phe Asp Phe Pro Glu Lys Pro 225 230

<210 <211 <212 <213	> 1 > D	022 NA	dops	is t	hali	ana										
<220 <221 <222 <223	> C > (	DS 31). 986	. (84	6)												
<400 catt	> 5 aaat	5 tgg	ctcc	tgtg	ja ac	ctaa	attt	atg Met	gac Asp	tat Tyr	gat Asp	ccc Pro	aac Asn	acc Thr	aat Asn	54
ccg Pro	ttc Phe 10	gac Asp	ctt Leu	cat His	ttc Phe	tcc Ser 15	ggt Gly	aaa Lys	ctt Leu	ccg Pro	aaa Lys 20	aga Arg	gaa Glu	gtc Val	tcg Ser	102
gct Ala 25	tca Ser	gct Ala	tct Ser	aaa Lys	gtt Val 30	gta Val	gag Glu	aag Lys	aaa Lys	tgg Trp 35	tta Leu	gtg Val	aaa Lys	gat Asp	gag Glu 40	150
aag Lys	aga Arg	aat Asn	atg Met	cta Leu 45	caa Gln	gat Asp	gaa Glu	ata Ile	aac Asn 50	cgg Arg	gtt Val	aat Asn	tcg Ser	gag Glu 55	aac Asn	198
aag Lys	aag Lys	cta Leu	acc Thr 60	gaa Glu	atg Met	tta Leu	gca Ala	aga Arg 65	gtc Val	tgt Cys	gag Glu	aag Lys	tac Tyr 70	tat Tyr	gct Ala	246
ctt Leu	aat Asn	aat Asn 75	ctt Leu	atg Met	gag Glu	gag Glu	ttg Leu 80	GIn	agt Ser	cga Arg	aag Lys	agt Ser 85	cct Pro	gaa Glu	agt Ser	294
gtt Val	aac Asn 90	ttt Phe	cag Gln	aac Asn	aaa Lys	cag Gln 95	cta Leu	acg Thr	999 Gly	aaa Lys	cga Arg 100	aaa Lys	caa Gln	gaa Glu	ctt Leu	342
gat Asp 105	gag Glu	ttt Phe	gtt Val	agc Ser	tcc Ser 110	cca Pro	att Ile	gga Gly	ctc Leu	agt Ser 115	ctc Leu	gga Gly	cca Pro	atc Ile	gag Glu 120	390
aac Asn	atc Ile	acc Thr	aac Asn	gat Asp 125	aaa Lys	gcg Ala	acg Thr	gtt Val	tca Ser 130	acc Thr	gct Ala	tac Tyr	ttt Phe	gct Ala 135	gct Ala	438
gag Glu	aag Lys	tct Ser	gac Asp 140	aca Thr	agc Ser	ttg Leu	act Thr	gtg Val 145	aaa Lys	gat Asp	gga Gly	tat Tyr	caa Gln 150	tgg Trp	agg Arg	486
aaa Lys	tac Tyr	999 Gly 155	caa Gln	aag Lys	att Ile	acg Thr	aga Arg 160	Asp	aat Asn	cca Pro	tct Ser	cct Pro 165	Arg	gct Ala	tac Tyr	534
ttc Phe	aga Arg 170	Cys	tcg Ser	ttt Phe	tca Ser	ccg Pro 175	tct Ser	tgt Cys	cta Leu	gtc Val	aag Lys 180	ьys	aag Lys	gtg Val	caa Gln	582
cga Arg 185	Ser	gca Ala	gaa Glu	gat Asp	cca Pro 190	Ser	ttc Phe	ttg Leu	gta Val	gcc Ala 195	act Thr	tac Tyr	gaa Glu	999 Gly	aca Thr 200	630
cat	aac	cac	acc	gga	cca	cat	gca	agt		tcc age	agg 72	aca	gtg	aaa	ctt	678

His	Asn	His	Thr	Gly 205	Pro	His					List Arg			Lys 215	Leu	
														gag Glu		726
														ttg Leu		774
														99 ⁹ Gly		<b>822</b>
			cat His				tga	aagt	tete	cta g	gaaca	tgta	at a	ttct	gttt	876
tgt	ctat	tt t	gttg	gctca	t to	ctag	jtaaa	aag	gtaa	aaga	tttg	jttt	gat d	ettga	ttagg	936
aggo	ataç	gat ç	gtcaa	ttt	a at	gtgt	gtgt	ata	ataat	tac	atca	aato	cta a	agtat	ccaaa	996
aagg	gtca	acc o	ccat	ttta	t ct	tate	3									1022
<210 <213 <213 <213	l> 2 ?> E	66 271 PRT Arabi	idops	sis t	:hali	lana										
<400	)> 5	66														
Met 1	Asp	Tyr	Asp	Pro 5	Asn	Thr	Asn	Pro	Phe 10	Asp	Leu	His	Phe	Ser 15	Gly	
Lys	Leu	Pro	Lys 20	Arg	Glu	Val	Ser	Ala 25	Ser	Ala	Ser	Lys	Val 30	Val	Glu	
ГЛа	Lys	Trp 35	Leu	Val	Lys	Asp	Glu 40	Lys	Arg	Asn	Met	Leu 45	Gln	Asp	Glu	
Ile	Asn 50	Arg	Val	Asn	Ser	Glu 55	As'n	Lys	Lys	Leu	Thr 60	Glu	Met _.	Leu	Ala	
Arg 65	Val	Сув	Glu	Lys	Tyr 70	Tyr	Ala	Leu	Asn	Asn 75	Leu	Met	Glu	Glu	Leu 80	
Gln	Ser	Arg	Lys	Ser 85	Pro	Glu	Ser	Val	Asn 90	Phe	Gln	Asn	Lys	Gln 95	Leu	
Thr	Gly	Lys	Arg 100	Lys	Gln	Glu	Leu	Asp 105	Glu	Phe	Val	Ser	Ser 110	Pro	Ile	
Gly	Leu	Ser 115	Leu	Gly	Pro	Иe	Glu 120	Asn	Ile	Thr	Asn	Asp 125	Lys	Ala	Thr	
Val	Ser 130	Thr	Ala	Tyr	Phe	Ala 135	Ala	Glu	Lys	Ser	Asp 140	Thr	Ser	Leu	Thr	
Val 145	Lys	Asp	Gly	Tyr	Gln 150	Trp	Arg	Lys	Tyr	Gly 155	Gln	Lys	Ile	Thr	Arg 160	
Asp	Asn	Pro	Ser	Pro 165	Arg	Ala	Tyr	Phe	Arg 170	Сув	Ser	Phe	Ser	Pro 175		

Cys	Leu	Val	Lys 180	Lys	Lys	Val	Gln	Arg 185	Ser	Ala	Glu	Asp	Pro 190	Ser	Phe	
Leu	Val	Ala 195	Thr	Tyr	Glu	Gly	Thr 200	His	Asn	His	Thr	Gly 205	Pro	His	Ala	
Ser	Val 210	Ser	Arg	Thr	Val	Lys 215	Leu	Asp	Leu	Val	Gln 220	Gly	Gly	Leu	Glu	
Pro 225	Val	Glu	Glu	Lys	Lys 230	Glu	Arg	Gly	Thr	Ile 235	Gln	Glu	Val	Leu	Val 240	
Gln	Gln	Met	Ala	Ser 245	Ser	Leu	Thr	Lys	Asp 250	Pro	Lys	Phe	Thr	Ala 255	Ala	
Leu	Ala	Thr	Ala 260	Ile	Ser	Gly	Arg	Leu 265	Ile	Glu	His	Ser	Arg 270	Thr		
<210 <210 <210 <210	L> 1 2> 1	57 1230 DNA Arab:	idops	sis t	:hali	iana							٠			
<220 <220 <220 <220	1> ( 2>	CDS (1). G134!	. (108 9	39)												
<40 atg Met 1	act	57 agt Ser	cgg Arg	aga Arg 5	gaa Glu	gta Val	cgg Arg	tgc Cys	cgg Arg 10	tgc Cys	ggc Gly	aga Arg	cgg Arg	atg Met 15	tgg Trp	48
gtt Val	caa Gln	cca Pro	gac Asp 20	gcc Ala	cgt Arg	acc Thr	gtc Val	caa Gln 25	tgc Cys	tca Ser	acc Thr	Cys	cac His 30	acc Thr	gtc Val	96
acg Thr	cag Gln	ctc Leu 35	tac Tyr	tcg Ser	cta Leu	gtg Val	gac Asp 40	ata Ile	gct Ala	cgc Arg	ggt Gly	gca Ala 45	aac Asn	cgc	ata Ile	144
att Ile	cat His 50	ggg Gly	ttt Phe	caa Gln	cag Gln	cta Leu 55	ctt Leu	aga Arg	caa Gln	cac His	caa Gln 60	ccg Pro	caa Gln	cat His	cat His	192
gaa Glu 65	caa Gln	caa Gln	caa Gln	caa Gln	caa Gln 70	atg Met	atg Met	gct Ala	caa Gln	ccg Pro 75	cca Pro	cca Pro	cgg	ctg Leu	ctt Leu 80	240
gag Glu	cct Pro	ctt Leu	ccc Pro	tcg Ser 85	ccg Pro	ttt Phe	999 Gly	aag Lys	aag Lys 90	aga Arg	gca Ala	gtt Val	tta Leu	tgc Cys 95	ggc Gly	288
gtg Val	aac Asn	tat Tyr	aag Lys 100	gga Gly	aaa Lys	agt Ser	tat Tyr	agc Ser 105	Leu	aaa Lys	ggt Gly	tgc Cys	atc Ile 110	agt Ser	gat Asp	336
Āla	Lys	Ser 115		Arg	Ser	Leu	Leu 120	Val	Gln	Gln	Met	Gly 125	Phe	Pro	Ile	384
gac Asp	Ser 130	Ile	ctc Leu	atg Met	ctc Leu	aca Thr 135	gaa Glu	gat Asp	gaa Glu	gcc Ala	agc Ser 140	Pro	cag Gln	aga Arg	ata Ile	432
ccg	acg	aag	aga	aac	att	agg	aag	gcg		aga Page		tta	gtt	gaa	999	480

PCT/US00/31418 WO 01/35726

Pro	Thr	Lvs	Ara	Asn	Tle	Ara					List Trp				Glv	
145	•	3	3		150		-,-			155					160	
											tct Ser					528
cag Gln	cag Gln	aat Asn	gac Asp 180	tac Tyr	aac Asn	gga Gly	gac Asp	gag Glu 185	atc Ile	gat Asp	ggt Gly	caa Gln	gat Asp 190	gaa Glu	gcc Ala	<b>576</b>
ttg Leu	tgc Cys	cct Pro 195	tta Leu	gac Asp	cat His	gaa Glu	aca Thr 200	gaa Glu	gga Gly	aaa Lys	atc Ile	att Ile 205	gat Asp	gac Asp	gag Glu	624
att Ile	aac Asn 210	cgg Arg	ata Ile	ctc Leu	gtg Val	agg Arg 215	cct Pro	ctc Leu	gtc Val	cat His	gga Gly 220	gct Ala	aag Lys	ctt Leu	cac His	672
gct Ala 225	gtc Val	atc Ile	gac Asp	gcc Ala	tgt Cys 230	aac Asn	agc Ser	999 Gly	act Thr	gtc Val 235	ctt Leu	gat Asp	tta Leu	ccc Pro	ttc Phe 240	720
att Ile	Cys Cys	agg Arg	atg Met	gag Glu 245	agg Arg	aat Asn	ggt Gly	tct Ser	tat Tyr 250	gaa Glu	tgg Trp	gaa Glu	gac Asp	cat His 255	aga Arg	768
tca Ser	gtc Val	aga Arg	gct Ala 260	tac Tyr	aaa Lys	gga Gly	aca Thr	gat Asp 265	ggt Gly	gga Gly	gca Ala	gct Ala	ttc Phe 270	tgt Cys	ttc Phe	816
agt Ser	gct Ala	tgt Cys 275	gac Asp	gat Asp	gat Asp	gaa Glu	tcc Ser 280	agt Ser	ggt Gly	tac Tyr	act Thr	cct Pro 285	gtg Val	ttc Phe	acg Thr	864
											ata Ile 300					912
aca Thr 305	gct Ala	gga Gly	cca Pro	gca Ala	ccc Pro 310	acg Thr	tat Tyr	ggc Gly	cac His	ctg Leu 315	ctt Leu	aac Asn	ctt Leu	atg Met	tgt Cys 320	960
tct Ser	gca Ala	ata Ile	cga Arg	gag Glu 325	gcc Ala	cag Gln	.tct Ser	cgc Arg	ctc Leu 330	Ala	ttt Phe	aac Asn	Gly	gac Asp 335	tac Tyr	1008
aca Thr	agc Ser	tct Ser	gat Asp 340	gca Ala	tcc Ser	gcg Ala	gag Glu	cca Pro 345	ctg Leu	cta Leu	aca Thr	tca Ser	tct Ser 350	gag Glu	gaa Glu	1056
ttt Phe	gac Asp	gtg Val 355	tac Tyr	gcg Ala	aca Thr	aag Lys	ttt Phe 360	gta Val	ctc Leu	tga	atgo	etgt	aca 1	tatg	atgctg	1109
caaa	tago	etc 9	ggaa	acgti	tt c	tatgi	tgtai	t gta	atca	tgta	atga	atta	tgt 1	tgca	tagcct	1169
ctct	ctt	ett a	acga	gcaa	ta ag	gcta	tgaaa	a taa	attga	attc	gcta	aga	aat	ttaa	aatgaa	1229
a		-														1230

<400> 58

Met Ala Ser Arg Arg Glu Val Arg Cys Arg Cys Gly Arg Arg Met Trp 1 5 10 15

Val Gln Pro Asp Ala Arg Thr Val Gln Cys Ser Thr Cys His Thr Val 20 30 30

<210> 58 <211> 362 <212> PRT

<213> Arabidopsis thaliana

#### MBI15 Sequence Listing.ST25

Thr Gln Leu Tyr Ser Leu Val Asp Ile Ala Arg Gly Ala Asn Arg Ile 35 40 45

Ile His Gly Phe Gln Gln Leu Leu Arg Gln His Gln Pro Gln His His 50 55 60

Glu Gln Gln Gln Gln Met Met Ala Gln Pro Pro Pro Arg Leu Leu 65 70 75 80

Glu Pro Leu Pro Ser Pro Phe Gly Lys Lys Arg Ala Val Leu Cys Gly 85 90 95

Val Asn Tyr Lys Gly Lys Ser Tyr Ser Leu Lys Gly Cys Ile Ser Asp 100 105 110

Ala Lys Ser Met Arg Ser Leu Leu Val Gln Gln Met Gly Phe Pro Ile 115 120 125

Asp Ser Ile Leu Met Leu Thr Glu Asp Glu Ala Ser Pro Gln Arg Ile 130 135 140

Pro Thr Lys Arg Asn Ile Arg Lys Ala Met Arg Trp Leu Val Glu Gly 145 150 155 160

Asn Arg Ala Arg Asp Ser Leu Val Phe His Phe Ser Gly His Gly Ser 165 170 175

Gln Gln Asn Asp Tyr Asn Gly Asp Glu Ile Asp Gly Gln Asp Glu Ala 180 185 190

Leu Cys Pro Leu Asp His Glu Thr Glu Gly Lys Ile Ile Asp Asp Glu
195 200 205

Ile Asn Arg Ile Leu Val Arg Pro Leu Val His Gly Ala Lys Leu His 210 215 220

Ala Val Ile Asp Ala Cys Asn Ser Gly Thr Val Leu Asp Leu Pro Phe 225 230 235

Ile Cys Arg Met Glu Arg Asn Gly Ser Tyr Glu Trp Glu Asp His Arg 245 250 255

Ser Val Arg Ala Tyr Lys Gly Thr Asp Gly Gly Ala Ala Phe Cys Phe 260 265 270

Ser Ala Cys Asp Asp Asp Glu Ser Ser Gly Tyr Thr Pro Val Phe Thr 275 280 285

Gly Lys Asn Thr Gly Ala Met Thr Tyr Ser Phe Ile Lys Ala Val Lys 290 295 300

Thr Ala Gly Pro Ala Pro Thr Tyr Gly His Leu Leu Asn Leu Met Cys 305 310 315

Ser Ala Ile Arg Glu Ala Gln Ser Arg Leu Ala Phe Asn Gly Asp Tyr Page 76

MBI15 Sequence Listing.ST25 330 335

Thr Ser Ser Asp Ala Ser Ala Glu Pro Leu Leu Thr Ser Ser Glu Glu 340 345 350

Phe Asp Val Tyr Ala Thr Lys Phe Val Leu 355

. 325

# INTERNATIONAL SEARCH REPORT

Interna al application No.
PCT/US00/31418

A. CLAS	SIFICATION OF SUBJECT MATTER		··· · · ·	Į.				
IPC(7)	· A01H 1/00 5/00: C12N 5/14, 15/82			Ì				
US CI. 435/320.1, 419, 468, 800/278, 279, 287, 301, 305-310, 312, 314, 317, 320, 322								
According to	International Patent Classification (IPC) or to both nat	ional clas	sification and IPC					
B. FIEL	DS SEARCHED							
Minimum do	cumentation searched (classification system followed b	y classific	ation symbols)					
U.S. : 43	35/320.1, 419, 468; 800/278, 279, 287, 301, 305-310,	312, 314	, 317, 320, 322					
Documentation	on searched other than minimum documentation to the	extent tha	t such documents are included	l in the fields searched				
Electronic da EAST, USPA	ata base consulted during the international search (name AT, STN, Agricola, CaPlus, Biosis, Embase	e of data b	pase and, where practicable, so	earch terms used)				
C DOC	UMENTS CONSIDERED TO BE RELEVANT							
	Citation of document, with indication, where app	ropriate.	of the relevant passages	Relevant to claim No.				
Category *	WO 97/47183 A1 (PURDUE RESEARCH FOUNDA	TION) 1	December 1997	1-9, 12, 13, 25				
	(18.12.1997), entire reference.	·		10, 11, 26, 27				
Y				1 0 12 12 25				
X	US 5,939,601 (KLESSIG et al) 17 August 1999 (17.08.1999), entire reference.							
Y	10, 11,							
A	Database Genbank on NCBI, US National Library on No. AB009055, SATO, S. et al 'Strucural analysis on IV. Sequence features of the regions of 1,456,315 by assigned P1 and TAC clones. 27 December 2000, D 41-54, see bases 16,003-16,490, 16,571-16,683 and	f Arabido covered NA RES.	psis thaliana chromosome 5. by nineteen physically 1998, Vol. 5, No. 1, pages	1-13, 25-27				
			Con motors family annex					
Furthe	er documents are listed in the continuation of Box C.	<u> </u>	See patent family annex.					
1	Special categories of cited documents:	"T"	later document published after the in- date and not in conflict with the appli principle or theory underlying the in-	ication but cited to understand the				
"A" docume of parti	nt defining the general state of the art which is not considered to be rular relevance		• •					
, ·	application or patent published on or after the international filing date	"X"	document of particular relevance; the considered novel or cannot be considered movel or cannot be considered when the document is taken alone	ered to involve an inventive step				
"L" docume establis specific	nt which may throw doubts on priority claim(s) or which is cited to h the publication date of another citation or other special reason (as d)	-Y-	document of particular relevance; the considered to involve an inventive st combined with one or more other su	ep when the document is ch documents, such combination				
1	ent referring to an oral disclosure, use, exhibition or other means		being obvious to a person skilled in	the art				
	ent published prior to the international filing date but later than the date claimed	"&"	document member of the same pater					
Date of the	actual completion of the international search	Date of	mailing of the international se	earch report				
23 Februar	y 2001 (23.02.2001)		UY MAKE DO	20				
Name and	mailing address of the ISA/US	Authori	zed officer	RY J. DEY				
C	ornmissioner of Patents and Trademarks	David !		AL SPECIALIST				
	ox PCT /ashington, D.C. 20231		THE WILLS					
	May (703)305-3330	Telepho	ne No. 703-308- <b>TECHNOLO</b>	RA CENIER 1800				

Facsimile No. (703)305-3230
Form PCT/ISA/210 (second sheet) (July 1998)

## INTERNATIONAL SEARCH REPORT

Inter nal application No.

PCT/US00/31418

Box	I Obse	rvations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This	internat	ional report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1.		Claim Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
2.		Claim Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3.	6.4(a).	Claim Nos.: 14 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule
Box	п Ор	servations where unity of invention is lacking (Continuation of Item 2 of first sheet)
		ional Searching Authority found multiple inventions in this international application, as follows: ontinuation Sheet
1.		As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.		As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.		As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4.	$\boxtimes$	No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-13 & 25-27 and SEQ ID NOs 1&2
Rem	ark on l	Protest The additional search fees were accompanied by the applicant's protest.
		No protest accompanied the payment of additional search fees.

#### INTERNATIONAL SEARCH REPORT

Inte.......nal application No.

PCT/US00/31418

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Groups 1-XXIX, claim(s) 1-14 and 25-27, drawn to a transgenic plant having modified seed characteristics, polynucleotides and vectors for producing said transgenic plant and a method of making said transgenic plant. Applicant must elect one pair of sequences (one nucleic acid and the corresponding amino acid translation) to be examined, i.e. SEQ ID NO: 1 and 2 in Group I, SEQ ID NO: 3 and 4 in Group II, SEQ ID NO: 5 and 6 in Group III, etc.

Group XXX, claim(s) 15-17, drawn to a method of identifying a factor that is modulated.

Group XXXI, claims(s) 18, drawn to a method of identifying a molecule that modulates activity or expression of a polynucleotide or polypeptide.

Group XXXII, claims(s) 19 and 20, drawn to an integrated computer system.

Group XXXIII, claim(s) 21-24, drawn to a method for identifying a polynucleotide sequence comprising selecting a nucleic acid sequence from a database that meets a selected sequence criteria.

The inventions listed as Groups I-XXXIII do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The inventions listed as Groups I-XXXIII do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Groups I-XXIX are drawn to a transgenic plant and a method of producing said plant with a nucleic acid sequence. The methods of Groups I-XXIX differ from each other in that they are directed to a plant transformation method and transgenic plant with a structurally and functionally distinct nucleic acid sequence which encodes a structurally and functionally distinct amino acid sequence. In addition, Groups XXX, XXXI and XXXIII are different methods from any of Groups I-XXIX in that they have different method steps and different end products, and Group XXXII requires a computer system. Thus, there is no single special technical feature, which links the inventions of Groups I-XXXIII under PCT Rule 13.2.